Published in final edited form as:

J Am Chem Soc. 2008 December 3; 130(48): 16407–16416. doi:10.1021/ja8063205.

Total Syntheses of (+)-Tedanolide and (+)-13-Deoxytedanolide

Joshua R. Dunetz[†], Lisa D. Julian[‡], Jason S. Newcom[‡], and William R. Roush[†]

Department of Chemistry, Scripps Florida, Jupiter, Florida 33458, and Department of Chemistry, University of Michigan, Ann Arbor, Michigan 41809, E-mail: roush@scripps.edu

Abstract

Convergent total syntheses of the potent cytotoxins (+)-tedanolide (**1**) and (+)-13-deoxytedanolide (**2**) are described. The carbon framework of these compounds was assembled via a stereoselective aldol reaction that unifies the C(1)–C(12) ketone fragment **5** with a C(13)–C(23) aldehyde fragment **6** (for 13-deoxytedanolide) or **52** (for tedanolide). Multiple obstacles were encountered en route to (+)-**1** and (+)-**2** that required very careful selection and orchestration of the stereochemistry and functionality of key intermediates. Chief among these issues was the remarkable stability and lack of reactivity of hemiketals **33b** and **34** that prevented the tedanolide synthesis from being completed from aldol **4**. Key to the successful completion of the tedanolide synthesis was the observation that the 13-deoxy hemiketal **36** could be oxidized to C(11,15)-diketone **38** en route to 13-deoxytedanolide. This led to the decision to pursue the tedanolide synthesis via C(15)-(*S*)-epimers, since this stereochemical change would destabilize the hemiketal that plagued the attempted synthesis of tedanolide via C(15)-(*R*) intermediates. However, use of C(15)-(*S*) configured intermediates required that the side chain epoxide be introduced very late in the synthesis, owing to the ease with which the C(15)-(*S*)-OH cyclized onto the epoxide of intermediate **50**.

Introduction

The tedanolides are a group of 18-membered macrolactones that display remarkable biological activity. (+)-Tedanolide (**1**), reported by Schmitz and co-workers in 1984 after isolation from the Caribbean fire sponge *Tedania ignis*, is highly cytotoxic against human nasopharynx carcinoma (ED₅₀ = 250 pg/mL) and lymphocytic leukemia (ED₅₀ = 16 pg/mL) cell lines and causes cell accumulation in the S phase at concentrations as low as 10 ng/mL.¹ In addition, tedanolide has been shown to increase the lifespan of mice implanted with lymphocytic leukemia by 23% at 1.5 µg/kg of body weight.² The closely related macrolide (+)-13-deoxytedanolide (**2**) was isolated by Fusetani and co-workers in 1991 from the Japanese sponge *Mycale adhaerens* and demonstrates potent cytotoxicity against P388 murine leukemia cells (IC₅₀ = 94 pg/mL).³ In an elegant study designed to probe the mechanism of 13-deoxytedanolide activity, Fusetani and co-workers demonstrated that the macrolide binds to the eukaryotic 60S ribosomal subunit^{4a} and identified the C(11)–C(23) region of the natural product as the pharmacophore.^{4b,5} A third member of this macrolide family, (+)-tedanolide C (**3**), was reported by Ireland and co-workers in 2006 from the extract of Papua New Guinea marine sponge *Ircinia* sp.⁶ The reported structure of tedanolide C has different stereochemistry and a different methylation and oxygenation pattern than the two older congeners. Nevertheless, tedanolide C exhibits strong activity (IC₅₀ = 57 ng/mL) against the HTC-116 colorectal cancer cell line.

[†]Scripps Florida.

[‡]University of Michigan.

The impressive biological profiles and the synthetic challenges presented by the tedanolides have encouraged our laboratory^{7,8} and others to pursue their synthesis.^{9–17} The complex architecture of tedanolide (**1**) is underscored by a heavily oxygenated and methylated carbon skeleton containing 13 stereocenters and two polysubstituted olefins—one of which is incorporated into a β,γ -unsaturated ketone with the potential to migrate into conjugation with the carbonyl. In addition, tedanolide possesses a sensitive α -hydroxy trisubstituted epoxide and four β -hydroxy ketone units that are potentially susceptible to retro aldol decomposition. Total syntheses of tedanolide were recently completed by Kalesse^{9a,b} (2006) and Smith^{10a} (2007); a few years earlier, Smith^{10c,d} (2003) and our laboratory⁷ (2005) reported total syntheses of 13-deoxytedanolide.

Results and Discussions

In planning our synthetic approach to tedanolide and 13-deoxytedanolide, we initially envisioned that **4** would serve as a common intermediate that would permit entry into both macrolides (Scheme 1). We planned to assemble **4** from the convergent aldol coupling of methyl ketone **5** and aldehyde **6**, which would establish the C(13)-alcohol stereochemistry of tedanolide via Felkin addition of **5** to **6**. We chose the (*R*)-configuration for the C(15)-position of **6** after previous studies from our group demonstrated that the lithium enolates of methyl ketone models of **5** add to 2,3-*anti* aldehydes (e.g., **6**) with greater Felkin selectivity than additions to 2,3-*syn* aldehydes (e.g., **39**, vide infra).¹⁸ We envisaged that aldol **4** could be converted to tedanolide via a sequence involving C(15)-oxidation to the requisite carbonyl and macrolactonization. After considerable experimentation,^{8b} our decision to mask the C(1)-acid in **5** as an allyl ester and the C(16)-hydroxymethyl group in **6** as an allyl carbonate (Alloc) was motivated by our desire to remove both protection groups simultaneously prior to macrolactonization. We envisioned that aldol adduct **4** could also be converted to 13-deoxytedanolide by using a similar sequence that also included deoxygenation of the C(13)-alcohol.

We developed a convergent route that assembles the C(1)–C(12) ketone **5** via the aldol reaction of ethyl ketone **7** and aldehyde **8**. The synthesis of ethyl ketone **7** began with asymmetric hydrogenation¹⁹ of β -keto ester **9** followed by alkylation of the β -hydroxy ester under Frater conditions²⁰ to access the 2,3-*anti* ester **10** (Scheme 2). The hydroxyl group of **10** was protected as a DMPM ether in **11** through a sequence involving ester reduction with LiAlH_4 , conversion of the 1,3-diol to the 3,4-dimethoxybenzylidene acetal, and regioselective acetal reductive opening with DIBAL. The primary alcohol of **11** was oxidized under Swern conditions,²¹ and the resulting aldehyde was treated with 3 equiv of γ -siloxy allylstannane **12**^{22a} and $\text{BF}_3\text{-OEt}_2$ in CH_2Cl_2 at -78°C ^{22b} to afford the 3,4-*syn*-4,5-*syn*-homoallylic alcohol with >95:5 dr. Methylation of the alcohol (MeOTf , 2,6-di-*tert*-butyl-4-methylpyridine)²³ and oxidative cleavage of the olefin provided aldehyde **14**, which in turn was converted to allyl ester **15** via chlorite oxidation to the carboxylic acid²⁴ and esterification with allyl alcohol under Mitsunobu conditions.²⁵

The conversion of allyl ester **15** to the ethyl ketone **7** was complicated by the competitive formation of lactone **16** when **15** was treated with DDQ²⁶ followed by oxidation of the resulting secondary alcohol via the Parikh-Doering method (eq 1).²¹ Lactone **16** is produced during isolation of the C(5)-alcohol after DDQ-mediated benzyl ether cleavage. This side reaction became more prominent when the debenzylation-oxidation sequence was performed on gram scale. The alcohol intermediate proved very sensitive to acid,²⁷ and great care was taken to minimize the formation of lactone **16** by removing the mildly acidic dihydroquinone (DDQ byproduct) with aqueous sodium bicarbonate extractions.

(1)

Aldehyde **8** (to be coupled with ethyl ketone **7** en route to C(1)–C(12) ketone **5**) was synthesized from the readily available *anti*- β -hydroxy- α -methylbutyrate **17**²⁸ as shown in Scheme 3. Ester **17** was converted to aldehyde **18** via ester reduction with LiAlH₄, dimethoxybenzylidene acetal formation, regioselective acetal reductive opening, and Swern oxidation of the resulting primary alcohol. Wittig olefination of **18** with the stabilized ylide Ph₃P=C(Me)CO₂Et provided the α,β -unsaturated ester **19** with excellent selectivity (>95:5 *E/Z*). Reduction of **19** to the allylic alcohol followed by Parikh-Doering oxidation²¹ of the primary alcohol furnished aldehyde **8** in 70% yield over three steps.

The stereoselective aldol coupling of ethyl ketone **7** and aldehyde **8** was effected by using TiCl₄ and *i*-Pr₂NEt and provided **20** in 73% yield with 10:1 diastereoselectivity (Scheme 4).²⁹ The selectivity of this transformation stems from a transition state **TS** in which the C–O bond of the titanium enolate eclipses the α' -C–H bond, allowing the aldehyde to approach from the less-hindered face of the enolate.²⁹ Extensive optimization was required to minimize the formation of **21** via intramolecular cyclization of the titanium enolate onto the allyl ester. This competitive pathway was suppressed by treating ketone **7** with TiCl₄ and *i*-Pr₂NEt for a maximum of 8 min; longer exposure times led to substantially lower yields (<30%) of **20** and correspondingly increased amounts of **21**. The synthesis of C(1)–C(12) ketone **5** was completed in three steps with the silylation of the C(7)-alcohol of **20**, DDQ cleavage of the dimethoxybenzyl ether,²⁶ and TPAP oxidation³⁰ of the liberated secondary alcohol.

The synthesis of C(13)–C(23) aldehyde **6** is summarized in Scheme 5. Wittig olefination of the known aldehyde **22**³¹ with the stabilized ylide Ph₃P=C(Me)CO₂Et furnished the α,β -unsaturated ester in excellent selectivity (97:3 *E/Z*), and DIBAL reduction provided the allylic alcohol. Subsequent Sharpless epoxidation³² of the allylic alcohol (>18:1 dr) and Parikh-Doering oxidation of the hydroxyl group afforded epoxyaldehyde **23**. Evans aldol reaction^{33a} of **23** with the chiral crotonate imide **24**,^{33b} silylation of the newly-formed alcohol and reduction of the acyl oxazolidinone with LiBH₄ in aqueous THF³⁴ provided homoallylic alcohol **25a**. The primary alcohol of **25a** was temporarily protected as a 2-bromoethyl carbonate (BEC); unfortunately, the Alloc group we desired at this position in **6** could not be directly installed due to its incompatibility with subsequent olefin oxidative cleavage steps. Standard oxidative cleavage of the vinyl group of BEC-protected **25b** furnished an aldehyde that was converted to **27** via asymmetric crotylboration with (*S,S*)-**26**.³⁵

The conversion of **27** to the targeted aldehyde **6** required the installation of the C(21)–C(22) (*Z*)-olefin and conversion of the terminal C(13)-alkene to the aldehyde. Our initial efforts to chemoselectively functionalize the C(13)-alkene in the presence of the C(21)–C(22) olefin were unsuccessful; as a result, it was necessary to functionalize the terminal alkene before introducing the C(21)–C(22) olefin. This was accomplished by silylation of the secondary alcohol of **27**, dihydroxylation of the terminal alkene, and protection of the resulting diol as two α -methoxyacetate esters in **28**. The C(21)–C(22) (*Z*)-olefin was installed upon DDQ oxidative cleavage of the PMB ether of **28**, Dess-Martin oxidation³⁶ of the primary alcohol,

and Wittig olefination of the resulting aldehyde. Treatment of **29** with allyl alcohol and MeMgBr, which generates the magnesium alkoxide in situ, converted the bromoethyl carbonate (BEC) to the allyl carbonate (Alloc) and also cleaved the two α -methoxyacetate esters. Treatment of the liberated diol with Pb(OAc)₄ then furnished the targeted C(13)–C(23) aldehyde **6**.

Completion of the synthesis of the tedanolide backbone was accomplished by converting methyl ketone **5** to its lithium enolate with LHMDs followed by the addition of aldehyde **6** (Scheme 6). This aldol reaction provided adduct **4** as a single diastereomer³⁷ in 59% yield, along with recovered ketone **5** (27%) and aldehyde **6** (11%). Surprisingly, our efforts to protect the newly-formed C(13)-alcohol as a silyl ether (TMS or TES) or acetate ester (Ac or AcOMe) were unsuccessful, and we carried the unprotected aldol **4** forward. Cleavage of the allyl ester and allyl carbonate units by treatment with Pd(PPh₃)₄ and *n*-Bu₃SnH³⁸ liberated the seco-acid for macrolactonization;³⁹ however, efforts to obtain macrolactone **30** were complicated by the formation of 14-membered macrolactone **31** via competitive cyclization onto the unprotected C(13)-alcohol. We did not observe equilibration of **30** and **31** upon treatment of these macrolactones with titanium isopropoxide.

In an attempt to circumvent the competitive formation of **31**, we tried to alter the conformation of the seco-acid by installing the C(15)-ketone prior to macrolactonization. Selective cleavage of the C(15)-TES ether of **4** with aqueous AcOH in THF led to secondary alcohol **32a** that cyclized onto the C(11)-carbonyl to form hemiketal **33a** (Scheme 7). Silylation of the C(13)-alcohol of **33a** with TESCl provided the hemiketal **33b** which proved to be remarkably stable and unreactive toward oxidants. Extensive efforts to intercept the ring-opened isomer **32b** with oxidants (Dess-Martin periodinane,³⁶ SO₃-pyridine in DMSO,²¹ H₂CrO₆ (Jones oxidation),⁴⁰ PDC,⁴¹ TPAP/NMO³⁰) en route to the C(15)-ketone failed. We postulated that converting these intermediates to the macrolactone **34** might shift the hemiketal/hydroxy ketone equilibrium to the hydroxy ketone isomer **35** and permit C(15)-oxidation. Therefore, conversion of **33b** to the seco-acid and then macrocyclization as described for **4** provided macrocyclic hemiketal **34** (Scheme 8). However, the hydroxy ketone tautomer **35** could not be intercepted with oxidants to afford the targeted C(11,15)-dione, and under various oxidation conditions the hemiketal **34** was recovered intact. As a result, we were unable to elaborate **34/35** to tedanolide even though these intermediates are only two synthetic steps from the natural product (i.e., C(15)-oxidation and global desilylation).

Although it was not possible to convert hemiketal **33b** or **34** to tedanolide, we were able to use **33a** for the synthesis of 13-deoxytedanolide (Scheme 9). By analogy to the principles of the Thorpe-Ingold effect,^{42,43} which stipulates that increased numbers of substituents (typically in a geminal arrangement) between two reactive termini lead to an increased rate of ring formation, we speculated that removal of the C(13)-alcohol would decrease the stability of the hemiketal relative to the hydroxy ketone tautomer and enable oxidation of C(15)-OH to the C(11,15)-dione. Excision of C(13)-OH of **33a** was accomplished via conversion of the alcohol to the pentafluorophenylthiocarbonate⁴⁴ and subsequent treatment with Et₃B and *n*-Bu₃SnH.⁴⁵ Although the resulting hemiketal **36** did not appear to be in equilibrium with hydroxy ketone isomer **37** according to ¹H NMR analysis, we were gratified that treatment of **36** with Dess-Martin periodinane furnished the triketone **38** in 71% yield. Ketone **38** was converted to 13-deoxytedanolide (**2**) in three steps via seco-acid formation and Yonemitsu-modified Yamaguchi lactonization,³⁹ followed by removal of the three TBS ethers with Et₃N-buffered Et₃N-3HF (thereby generating Et₃N-2HF in situ).⁴⁶ Synthetic 13-deoxytedanolide was identical in all respects (¹H NMR, ¹³C NMR, IR, optical rotation, HRMS) to an authentic sample.

Having completed the synthesis of the 13-deoxy congener, we continued our efforts to synthesize tedanolide which had been thwarted by the stability of hemiketal intermediates **33b** and **34**. We hypothesized that inversion of the C(15)-(*R*)-alcohol stereochemistry in our first generation intermediates to the C(15)-(*S*)-configuration would destabilize the corresponding hemiketal **B** via increased 1,3-diaxial interactions and permit the requisite C(15)-oxidation (Scheme 10).^{47,48} Thus, we targeted aldehyde **39** containing the (*S*)-configuration at the C(15)-position.

We were unable to install the C(15)-(*S*)-OH of **39** via crotylboration (Scheme 11) of **40** (the immediate precursor to **27**, Scheme 6), which was surprising since a similar crotylboration reaction set the C(15)-(*R*)-OH of aldehyde **6** (vide supra). The reactions of (*R,R*)-tartrate crotylboronate³⁵ or (–)-Ipc₂-(*Z*)-crotylboronane⁴⁹ with aldehyde **40** did not provide **41**, but rather gave either the undesired C(15)-(*R*)-OH diastereomer (i.e., **27**) or unidentified products lacking the diagnostic olefin protons in the ¹H NMR spectrum. Alternatively, we pursued this key stereocenter using an Evans aldol approach.^{33a} Reaction of aldehyde **40** with chiral imide **42** provided the aldol adduct **43** with excellent yield and selectivity. Treatment of the alcohol with TESCl afforded the silyl ether. However, efforts to cleave the acyl oxazolidinone unit of **43** led to competitive reduction of the 2-bromoethyl carbonate, and so we looked to replace the carbonate protection group with a silyl ether as shown in Scheme 12. Aldol reaction of chiral imide **42** with aldehyde **45** and subsequent reduction of the acyl oxazolidinone provided 1,3-diol **47**. The primary alcohol of **47** was temporarily protected as a methoxyacetate ester before silylation of the secondary alcohol. Subsequent cleavage of the methoxyacetate ester followed by oxidation of the resulting alcohol completed the synthesis of aldehyde **49** with the desired (*S*)-configuration at the C(15)-position.

The aldol reaction of the lithium enolate generated from C(1)–C(12) ketone **5** and C(13)–C(23) aldehyde **49** led to the Felkin adduct **50** as the major product³⁷ with 78:22 diastereoselectivity (Scheme 13). The diminished diastereoselectivity (compared to the aldol coupling of **5** and 2,3-*anti* aldehyde **6**) was expected due to the mismatched 1,3-induction imparted by the 2,3-*syn* aldehyde **49**.^{18,50} Specifically, the expected Felkin chair-like transition state (**TS-1**) for the aldol reaction of **5** with 2,3-*syn* aldehyde **49** is destabilized relative to the Felkin chair-like transition state (**TS-2**) for the 2,3-*anti* aldehyde **6** due to the stereoelectronic repulsion of interacting dipoles of the aldehyde carbonyl and C(15)-hydroxy TES ether.⁵⁰ In an attempt to move the synthesis forward, aldol product **50** was treated with aqueous AcOH in THF to cleave the TES ether. Unfortunately, these desilylation conditions led to the 5-*exo*-cyclization of the liberated C(15)-alcohol onto the epoxide⁵¹ to form tetrahydrofuran **51**. *The analogous cyclization was not observed during manipulation of aldol 4 deriving from the 2, 3-anti-aldehyde 6* (Scheme 7). We hoped that cleavage of the C(15)-TES ether under buffered or basic conditions (Et₃N-3HF, TBAF, TAS-F) might avoid the epoxide-opening, but these efforts led to competitive desilylation of the various TBS ethers. Nevertheless, even by using these neutral or weakly basic conditions, evidence for cyclization of C(15)-OH onto the epoxide was observed (e.g., **51** and desilylated versions of **51** were detected). Thus, the change in stereochemistry of C(15)-OH significantly affected the reactivity of intermediate **50** (Scheme 13). As a result, we were unable to elaborate these intermediates to tedanolide.

At this point, we targeted aldehyde **52** as a third-generation coupling partner for ketone **5** en route to tedanolide (eq 2). The C(13)–C(23) fragment **52** maintains the (*S*)-configuration at the C(15)-position but lacks the epoxide. This revision to our approach to tedanolide would now require a late-stage, C(17)-OH-directed epoxidation of the C(18)–C(19) olefin,^{12c,52} a strategy utilized by Smith^{10c,d} in the synthesis of 13-deoxytedanolide and by Kalesse^{9a,b} and Smith^{10a} in their syntheses of tedanolide. Aldehyde **52** was designed with protecting groups for the C(15)- and C(17)-alcohols that could be selectively removed prior to C(15)-alcohol

oxidation and C(17)-directed epoxidation, respectively, at appropriate junctions of the synthesis following the aldol coupling with **5**.

(2)

We sought to construct the C(13)–C(23) framework of aldehyde **52** via the anti-aldol reaction of aldehyde **53** and ketone **54** (eq 2) using conditions developed by Paterson (enol borinate formation with *c*-Hex₂BCl, Et₃N in Et₂O at 0 °C followed by aldehyde addition).⁵³ This convergent assembly would set the C(16)- and C(17)-stereocenters in a single operation through a transition state believed to minimize electronic repulsion between the oxygen lone pairs of the enol borinate and the benzyl ether.^{53b} We were encouraged by literature precedent showing that β-siloxy ethyl ketones participate in the anti-aldol reaction under Paterson's conditions without β-elimination of the siloxy substituent.^{12c} Furthermore, after the completion of this work,^{8a} Kalesse and co-workers reported the successful anti-aldol coupling of aldehyde **53** with the C(13)-OPMB analog of ketone **54** under similar conditions.^{9a}

Fragments **53** and **54** were synthesized by using easily scalable routes (Scheme 14). The synthesis of **53**^{9a,d} commenced with the Wittig olefination of trityl-protected aldehyde **55** followed by cleavage of the trityl ether under acidic conditions. Swern oxidation of alcohol **56** afforded β,γ-unsaturated aldehyde **57**. The stabilized ylide Ph₃P=C(Me)CO₂Et was added directly to the Swern reaction mixture to generate α,β-unsaturated ester **58** in a one-pot oxidation-olefination sequence. Reduction of the ester and subsequent MnO₂ oxidation of the allylic alcohol then provided aldehyde **53**. Ketone **54** was prepared from the DMPM-protected **60**⁵⁴ in four synthetic steps. Addition of the lithium enolate of *tert*-butyl acetate to **60** resulted in a β-hydroxy ester that was reduced to a 1,3-diol with LiAlH₄. Selective silylation of the primary alcohol and oxidation of the secondary alcohol furnished the requisite ketone **54**.

Our early efforts to unify aldehyde **53** and ketone **54** using Paterson's anti-aldol conditions led to **61** as the major of two diastereomers with 80:20 selectivity (Scheme 15). Literature precedent^{9a,12c} suggested that **62** would be the minor product; however, Mosher ester analysis⁵⁵ revealed that the C(17)-alcohol of the major and minor aldols both have identical (*S*)-configurations. Independent conversion of the two aldol products to their corresponding benzylidene acetals (of type **65**) and acetonides (of type **66**) revealed through spectroscopic analysis⁵⁶ that both aldol adducts have identical stereochemistry within the C(14)–C(17) positions; that is, both aldol diastereomers are anti-aldol products with (*S*)-configurations at C(17)-OH.

We speculated that **63** might be the correct structure of the minor aldol product, and that it might arise from epimerization of the bisallylic C(20)-stereocenter during the synthesis of aldehyde **53**. This hypothesis was supported by our observation that only one diastereomer is formed from the aldol coupling of ketones **54** or **67** with methacrolein under identical conditions (eq 3). Furthermore, Mosher ester analysis of the primary alcohol **59** (precursor to **53**, Scheme 14) revealed an 80:20 ratio of diastereomers at the C(20)-bisallylic stereocenter. We concluded that C(20)-epimerization was occurring during the oxidation-olefination sequence used in the conversion of alcohol **56** (>95:5 er as determined by Mosher ester analysis) to α,β-unsaturated ester **58**. After vigorously investigating different oxidation and olefination conditions,^{8a} we found that treatment of **56** with Dess-Martin periodinane at

0 °C, followed by ylide addition and olefination at 0 °C provided ester **58** (and subsequently aldehyde **53**) with 94:6 enantiomeric ratio. When aldehyde **53** (of 94:6 er) was used in the anti-aldol reaction with **54**, we obtained a 94:6 mixture of aldol diastereomers **61** and **63** (eq 4). It is remarkable that the aldol selectivity perfectly matched the enantiomeric purity of aldehyde **53**. These experiments were also crucial to our ability to bring sufficient material forward en route to tedanolide.

(3)

(4)

The synthesis of C(13)–C(23) aldehyde **52** from **61** was completed in five steps (Scheme 16). Reduction of the β -hydroxy ketone **61** to the *syn*-1,3-diol with DIBAL and selective silylation⁵⁷ of the allylic alcohol provided TES ether **70**. The dimethoxybenzyl ether of **70** was transferred to the secondary alcohol via benzylidene acetal formation followed by regioselective acetal reductive opening with DIBAL. Finally, oxidation of primary alcohol **71** using the Dess-Martin periodinane provided aldehyde **52** in excellent yield.

Addition of the lithium enolate of ketone **5** to aldehyde **52** provided Felkin adduct **72** (as determined by NMR analysis³⁷ and Mosher ester analysis⁵⁵) in 77% yield with 75:25 selectivity (Scheme 17). The diastereomeric ratio for this transformation is similar to that observed previously for the aldol reaction of 2,3-*syn* aldehyde **49** (vide supra), and variations to the stoichiometry of aldehyde **52** (0.8–1.4 equiv) relative to ketone did not affect the yield (based on limiting reagent) or selectivity. At this point, we looked to protect the C(13)-alcohol of **72** before further elaborating the intermediate to the natural product. Treatment of **72** with TBSOTf and 2,6-lutidine in CH_2Cl_2 led to the product **73** as a mixture of TES and TBS ethers at the C(17)-position. It seemed unlikely that this substitution is due to the direct silylation of the C(17)-OTES oxygen, as the TES:TBS ratio at this position was unchanged when a mixture of product **73**-TES and **73**-TBS was resubjected to the silylation reaction conditions. However, attempts to suppress the scrambling of the silyl ether protecting group at this position were unsuccessful.

Instead, protection of the C(13)-OH of **72** as a SEM ether proceeded without complication in quantitative yield (Scheme 18). At this point, we planned to convert the C(15)-dimethoxybenzyl ether to the C(15)-ketone via cleavage of the DMPM group and subsequent oxidation. Treatment of SEM-protected **72** with DDQ liberated the C(15)-(*S*)-OH of **74**; however, hemiketal **75** was observed as the exclusive product by spectroscopic analysis.

Furthermore, efforts to intercept the hydroxy ketone isomer **74** with oxidants (e.g., DMP, SO₃-pyridine in DMSO) were unsuccessful, and we were unable to obtain the desired C(15)-ketone **76**.

Our next course of action was to form the macrolactone prior to C(15)-oxidation (Scheme 19). We postulated that conformational constraints imposed by the macrocycle might destabilize the hemiketal, or even disfavor hemiketal formation from the hydroxy ketone. Therefore, the C(13)-alcohol of aldol adduct **72** was protected as a SEM ether, and the seco-acid was liberated via stepwise cleavage of the primary TBS ether (along with the C(17)-TES ether⁵⁸) to provide **77**, followed by removal of the allyl ester. Macrolactonization of this seco-acid was effected at room temperature using the Yonemitsu modification of the Yamaguchi protocol.³⁹ The cyclization was completely selective for the primary alcohol (no reaction at the secondary C(17)-OH was observed)⁵⁹ and the macrolactone was obtained in 32% overall yield from **72** (4 steps). Finally, the C(17)-OH was protected as a SEM ether to provide **78**.

Treatment of **78** with DDQ liberated the C(15)-alcohol. *Spectroscopic analysis revealed that **79** exists as the open chain hydroxy ketone, and we did not detect the hemiketal isomer.* We were gratified that treatment of **79** with Dess-Martin periodinane afforded the C(11,15)-dione **80** in 71% yield over two steps from DMPM ether **78**. The synthesis of (+)-tedanolide (**1**) was then completed in three steps. Removal of the two SEM ethers was accomplished by using MgBr₂ and EtSH in Et₂O⁶⁰ (as previously demonstrated by Smith^{10a}). C(17)-Hydroxyl-directed epoxidation of the C(18)–C(19) olefin with *m*-CPBA^{12c} gave a single epoxide diastereomer in 54% yield. Finally, global desilylation of the penultimate intermediate with Et₃N-buffered Et₃N-3HF⁴⁶ provided synthetic (+)-tedanolide. The spectroscopic data (¹H NMR, ¹³C NMR, IR, optical rotation, HRMS) from our synthetic tedanolide were in excellent agreement with data obtained for an authentic sample,¹ as well as with data reported by Kalesse^{9a,b} and Smith^{10a} for synthetic tedanolide.

Summary

Convergent, stereocontrolled total syntheses of (+)-tedanolide and (+)-13-deoxytedanolide have been accomplished. A key transformation for the assembly of both natural products is the aldol coupling of the C(1)–C(12) methyl ketone **5** with a C(13)–C(23) aldehyde partner. Aldehyde **6** with 2,3-*anti* stereochemistry was initially employed, and its aldol reaction with the lithium enolate generated from **5** proceeded with >95:5 selectivity for the Felkin aldol **4**. However, it proved impossible to elaborate **4** to tedanolide owing to the remarkable stability of hemiketals **33b** and **34** and especially their lack of reactivity toward oxidants. Deoxygenation of hemiketal **33a** provided the 13-deoxy hemiketal **36** which was successfully oxidized to the C(11,15)-diketone **38** by using the Dess-Martin periodinane reagent. The later intermediate was smoothly elaborated to (+)-13-deoxytedanolide.

Recognition that 13-deoxy hemiketal **36** could be oxidized by way of its hydroxy ketone tautomer **37** suggested that the remarkable stability and lack of reactivity of hemiketals **33b** and **34** might be due to the equatorial conformation of ring substituents in **33b** and **44** which stabilizes the hemiketal relative to the acyclic hydroxy ketone isomer. This then led to the decision to redesign the synthesis of tedanolide to proceed by way of the stereochemically inverted C(15)-(S)-alcohol intermediates, since hemiketal intermediates with C(15)-(S) series would be destabilized by having at least one large substituent in an axial position (Scheme 10). However, use of C(15)-(S)-configured intermediates also required that the side chain epoxide unit be introduced very late in the synthesis, owing to the ease with which C(15)-(S)-OH cyclized onto the epoxide in intermediate **50** (Scheme 13).

Therefore, 2,3-*syn* aldehyde **52** was employed in the aldol coupling with methyl ketone **5** which gave the Felkin aldol **72** with 75:25 selectivity. Elaboration of Felkin aldol **72** to macrolactone **78** set the stage for removal of the C(15)-DMPM ether and oxidation of the C(15)-OH to the required ketone in **80**. Fortunately, the hemiketal that plagued the unsuccessful tedanolide syntheses via C(15)-(*R*)-configured intermediates **33b** and **34** was not observed for the C(15)-(*S*)-hydroxy ketone **79** derived from **78**. Intermediate **80** was smoothly elaborated to tedanolide via deprotection of C(17)-OH, which was used to direct the epoxidation of the adjacent C (18,19)-olefin. Final, global cleavage of all silyl ethers then completed the total synthesis of (+)-tedanolide.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgements

Support from the National Institutes of Health (GM 038436) and a Bristol-Myers Squibb Graduate Fellowship to L.D.J. is gratefully acknowledged. We thank Gregory C. Lane for preliminary work^{8c} involving the synthesis of alcohol **25a** from aldehyde **22** (Scheme 5). We also thank Professor Schmitz for supplying an authentic sample of (+)-tedanolide, and Professors Fusetani and Matsunaga for supplying an authentic sample of (+)-13-deoxytedanolide.

References

1. Schmitz FJ, Gunasekera SP, Yalamanchili G, Hossain MB, van der Helm D. *J Am Chem Soc* 1984;106:7251.
2. Schmitz, F. J.; Gunasekera, S. P.; Hossain, M. B.; van der Helm, D.; Yalamanchili, G. (1988) U.S. Patent Application 87-7347.
3. Fusetani N, Sugawara T, Matsunaga S. *J Org Chem* 1991;56:4971.
4. (a) Nishimura S, Matsunaga S, Yoshida M, Hirota H, Yokoyama S, Fusetani N. *Bioorg Med Chem* 2005;13:449. [PubMed: 15598566] (b) Nishimura S, Matsunaga S, Yoshida S, Nakao Y, Hirota H, Fusetani N. *Bioorg Med Chem* 2005;13:455. [PubMed: 15598567]
5. For structure-activity relationship studies of the structurally-related myriaporones 3 and 4: Hines J, Roy M, Cheng H, Agapakis CM, Taylor R, Crews CM. *Mol BioSyst* 2006;2:371. [PubMed: 16880957]
6. Chevallier C, Bugni TS, Feng X, Harper MK, Orendt AM, Ireland CM. *J Org Chem* 2006;71:2510. [PubMed: 16526806]
7. Total synthesis of (+)-13-deoxytedanolide: Julian LD, Newcom JS, Roush WR. *J Am Chem Soc* 2005;127:6186. [PubMed: 15853320]
8. Previous efforts toward tedanolide: (a) Dunetz JR, Roush WR. *Org Lett* 2008;10:2059. [PubMed: 18422319] (b) Roush WR, Newcom JS. *Org Lett* 2002;4:4739. [PubMed: 12489975] (c) Roush WR, Lane GC. *Org Lett* 1999;1:95. [PubMed: 10822542]
9. (a) Ehrlich G, Hassfeld J, Eggert U, Kalesse M. *Chem Eur J* 2008;14:2232. (b) Ehrlich G, Hassfeld J, Eggert U, Kalesse M. *J Am Chem Soc* 2006;128:14038. [PubMed: 17061881] (c) Hassfeld J, Eggert U, Kalesse M. *Synthesis* 2005;1183. (d) Ehrlich G, Kalesse M. *Synlett* 2005:655. (e) Hassfeld J, Kalesse M. *Synlett* 2002:2007.
10. (a) Smith AB III, Lee D. *J Am Chem Soc* 2007;129:10957. [PubMed: 17691787] (b) Cho CG, Kim WS, Smith AB III. *Org Lett* 2005;7:3569. [PubMed: 16048344] (c) Smith AB III, Adams CM, Lodise Barbosa SA, Degnan AP. *Proc Natl Acad Sci USA* 2004;101:12042. [PubMed: 15163795] (d) Smith AB III, Adams CM, Lodise Barbosa SA, Degnan AP. *J Am Chem Soc* 2003;125:350. [PubMed: 12517144] (e) Smith AB III, Lodise SA. *Org Lett* 1999;1:1249. [PubMed: 10825976]
11. (a) Jung ME, Zhang T-h. *Org Lett* 2008;10:137. [PubMed: 18067310] (b) Jung ME, Yoo D. *Tetrahedron Lett* 2008;49:816. (c) Jung ME, Yoo D. *Org Lett* 2007;9:3543. [PubMed: 17676748] (d) Jung ME, Lee CP. *Org Lett* 2001;3:333. [PubMed: 11428007] (e) Jung ME, Lee CP. *Tetrahedron Lett* 2000;41:9719. (f) Jung ME, Marquez R. *Org Lett* 2000;2:1669. [PubMed: 10880197] (g) Jung ME, Marquez R. *Tetrahedron Lett* 1999;40:3129.

12. (a) Wong CM, Loh TP. *Tetrahedron Lett* 2006;47:4485. (b) Loh TP, Feng LC. *Tetrahedron Lett* 2001;42:6001. (c) Loh TP, Feng LC. *Tetrahedron Lett* 2001;42:3223.
13. (a) Matsui K, Zheng B-Z, Kusaka S-i, Kuroda M, Yoshimoto K, Yamada H, Yonemitsu O. *Eur J Org Chem* 2001:3615. (b) Zheng B-Z, Yamauchi M, Dei H, Kusaka S-i, Matsui K, Yonemitsu O. *Tetrahedron Lett* 2000;41:6441. (c) Matsushima T, Nakajima N, Zheng BZ, Yonemitsu O. *Chem Pharm Bull* 2000;48:855. [PubMed: 10866148] (d) Zheng B-Z, Maeda H, Mori M, Kusaka S-i, Yonemitsu O, Matsushima T, Nakajima N, Uenishi J-i. *Chem Pharm Bull* 1999;47:1288. [PubMed: 10517009] (e) Matsushima T, Mori M, Zheng B-Z, Maeda H, Nakajima N, Uenishi J-i, Yonemitsu O. *Chem Pharm Bull* 1999;47:308. [PubMed: 10212381] (f) Matsushima T, Zheng B-Z, Maeda H, Nakajima N, Uenishi J-i, Yonemitsu O. *Synlett* 1999:780. (g) Matsushima T, Mori M, Nakajima N, Maeda H, Uenishi J-i, Yonemitsu O. *Chem Pharm Bull* 1998;46:1335. (h) Matsushima T, Horita K, Nakajima N, Yonemitsu O. *Tetrahedron Lett* 1996;37:385.
14. Liu JF, Abiko A, Pei Z, Buske DC, Masamune S. *Tetrahedron Lett* 1998;39:1873.
15. Iwata Y, Tanino K, Miyashita M. *Org Lett* 2005;7:2341. [PubMed: 15932193]
16. (a) Taylor RE, Hearn BR, Ciavarrri JP. *Org Lett* 2002;4:2953. [PubMed: 12182597] (b) Taylor RE, Ciavarrri JP, Hearn BR. *Tetrahedron Lett* 1998;39:9361.
17. (a) Nyavanandi VK, Nadipalli P, Nanduri S, Naidu A, Iqbal J. *Tetrahedron Lett* 2007;48:6905. (b) Nyavanandi VK, Nanduri S, Dev RV, Naidu A, Iqbal J. *Tetrahedron Lett* 2006;47:6667.
18. Gustin DJ, VanNieuwenhze MS, Roush WR. *Tetrahedron Lett* 1995;36:3443.
19. (a) Taber DF, Silverberg LJ. *Tetrahedron Lett* 1991;32:4227. For a review: (b) Noyori R, Ohkuma T. *Angew Chem Int Ed* 2001;40:40.
20. Fráter G, Müller U, Günther W. *Tetrahedron* 1984;40:1269.
21. For a review: Mancuso AJ, Swern D. *Synthesis* 1981:165.
22. (a) Keck GE, Abbott DE, Wiley MR. *Tetrahedron Lett* 1987;28:139. (b) Evans DA, Dart MJ, Duffy JL, Yang MG. *J Am Chem Soc* 1996;118:4322.
23. Evans DA, Ratz AM, Huff BE, Sheppard GS. *Tetrahedron Lett* 1994;35:7171.
24. (a) Lindgren BO, Nilsson T. *Acta Chem Scand* 1973;27:888. For a modification using 2-methyl-2-butene as chlorine scavenger: (b) Kraus GA, Taschner MJ. *J Org Chem* 1980;45:1175.
25. For a review: Mitsunobu O. *Synthesis* 1981:1.
26. Oikawa Y, Tanaka T, Horita K, Yoshioka T, Yonemitsu O. *Tetrahedron Lett* 1984;25:5393.
27. A sample of secondary alcohol in CDCl_3 (treated with K_2CO_3) completely converted to lactone 16 within 20 minutes, as observed by ^1H NMR analysis.
28. Available from methyl acetoacetate (see ref. 20) via a sequence of asymmetric hydrogenation and alkylation that parallels the conversion of 9 to 10 (Scheme 2).
29. Evans DA, Rieger DL, Bilodeau MT, Urpi F. *J Am Chem Soc* 1991;113:1047.
30. (a) Griffith WP, Ley SV, Whitcombe GP, White AD. *J Chem Soc, Chem Commun* 1987:1625. For a review: (b) Ley SV, Norman J, Griffith WP, Marsden SP. *Synthesis* 1994:639.
31. Harried SS, Lee CP, Yang G, Lee TIH, Myles DC. *J Org Chem* 2003;68:6646. [PubMed: 12919029]
32. For a review: Katsuki T, Martin VS. *Org React* 1996;48:1.
33. (a) Evans DA, Bartroli J, Shih TL. *J Am Chem Soc* 1981;103:2127. (b) Evans DA, Sjogren EB, Bartroli J, Dow RL. *Tetrahedron Lett* 1986;27:4957.
34. Penning TD, Djuric SW, Haack RA, Kalish VJ, Miyashiro JM, Rowell BW, Yu SS. *Synth Commun* 1990;20:307.
35. Roush WR, Palkowitz AD, Ando K. *J Am Chem Soc* 1990;112:6348.
36. Dess DB, Martin JC. *J Org Chem* 1983;48:4155.
37. The stereochemistry was assigned by using NMR methods: Roush WR, Bannister TD, Wendt MD, VanNieuwenhze MS, Gustin DJ, Dilley GJ, Lane GC, Scheidt KA, Smith WJ III. *J Org Chem* 2002;67:4284. [PubMed: 12054964]
38. Dangles O, Guibé F, Balavoine G, Lavielle S, Marquet A. *J Org Chem* 1987;52:4984.
39. For a macrolactonization review: Parenty A, Moreau X, Campagne J-M. *Chem Rev* 2006;106:911. [PubMed: 16522013]
40. For a review: Luzzio FA. *Org React* 1998;53:1.

41. Corey EJ, Schmidt G. *Tetrahedron Lett* 1979;20:399.
42. Beesley RM, Ingold CK, Thorpe JF. *J Chem Soc* 1915;107:1080.
43. Implicit to this assumption is the recognition that the equilibrium constant for cyclization of the hydroxy ketone to the hemiketal is determined by the ratio of the rates of formation and breakdown of the hemiketal intermediate. Thus, while the Thorpe-Ingold effect is kinetic in nature, it may also impact the equilibrium constant of reversible processes.
44. Barton DHR, Jaszberenyi JC. *Tetrahedron Lett* 1989;30:2619.
45. Nozaki K, Oshima K, Utimoto K. *Tetrahedron Lett* 1988;29:6125.
46. (a) Giudicelli MB, Picq D, Veyron B. *Tetrahedron Lett* 1990;31:6527. (b) McClinton MA. *Aldrichimica Acta* 1995;28:31.
47. Julian, LD. Studies toward the total synthesis of the tedanolides: Total synthesis of 13-deoxytedanolide. University of Michigan; 2005. Ph.D. thesis
48. While this work was in progress, Kalesse reported experimental evidence for this hypothesis (refs 9a and 9b). Kalesse credits us for this analysis (ref 9b).
49. Brown HC, Jadhav PK. *J Am Chem Soc* 1983;105:2092.
50. (a) Evans DA, Duffy JL, Dart MJ. *Tetrahedron Lett* 1994;35:8537. (b) Evans DA, Dart MJ, Duffy JL, Yang MG. *J Am Chem Soc* 1996;118:4322.
51. Loh and co-workers induced a similar cyclization to analyze epoxide stereochemistry in their tedanolide fragment synthesis (ref 12c).
52. For a review of hydroxy directed epoxidations of allylic alcohols: Adam W, Wirth T. *Acc Chem Res* 1999;32:703.
53. (a) Paterson I, Goodman JM, Isaka M. *Tetrahedron Lett* 1989;30:7121. (b) Vulpetti A, Bernardi A, Gennari C, Goodman JM, Paterson I. *Tetrahedron* 1993;49:685. Mechanistic discussion:
54. Prepared in analogous fashion to the known PMB-protected aldehyde: Organ MG, Wang J. *J Org Chem* 2003;68:5568. [PubMed: 12839448]
55. (a) Dale JA, Mosher HS. *J Am Chem Soc* 1973;95:512. For a useful explanation and examples of Mosher ester analysis: (b) Ohtani I, Kusumi T, Kashman Y, Kakisawa H. *J Am Chem Soc* 1991;113:4092.
56. Acetonide analysis: Rychnovsky SD, Rogers BN, Richardson TI. *Acc Chem Res* 1998;31:9.
57. Hicks JD, Huh CW, Legg AD, Roush WR. *Org Lett* 2007;9:5621. [PubMed: 18027963]
58. We were unable to remove the primary TBS ether without also cleaving the TES ether.
59. Similar selectivity was also observed by Smith and co-workers in their syntheses of tedanolide and 13-deoxytedanolide (refs 10a, c, d).
60. Kim S, Kee IS, Park YH, Park JH. *Synlett* 1991:183. SYNOPSIS TOC (Word Style "SN_Synopsis_TOC")

Figure 1.
The tedanolide natural products

Scheme 1.
Retrosynthetic analysis of the tedanolides

Scheme 2.
Preparation of allyl ester **15**

Scheme 3.
Synthesis of unsaturated aldehyde **8**

Scheme 4.
Completion of the synthesis of C(1)–C(12) ketone **5**

Scheme 5.
 Synthesis of C(13)–C(23) aldehyde **6**

Scheme 6.
Aldol coupling and macrolactonization

Scheme 7.
Hemiketal formation of C(11, 15)-hydroxy ketone

Scheme 8.
Hemiketal stability in the macrolactone

Scheme 9.
Synthesis of (+)-13-deoxytedanolide (2)

C(15)-(R)-OH leads to stable hemiketal with C(16)-C(23) in equatorial position

C(15)-(S)-OH leads to destabilized hemiketal with increased 1,3-diaxial interactions

Scheme 10.
Proposed effect of C(15)-configuration on hemiketal stability

Scheme 11.
 Unsuccessful efforts toward aldehyde **39**

Scheme 12.
Synthesis of C(13)–C(23) aldehyde **49**

Scheme 13.
Aldol coupling of ketone **5** and aldehyde **49**

Scheme 14.
 Synthesis of aldehyde **53** and ketone **54**

Scheme 15.
Aldol coupling of **53** and **54**

Scheme 16.
Completion of the synthesis of aldehyde **52**

Scheme 17.
 Aldol coupling of ketone **5** and aldehyde **52**

Scheme 18.
Attempted C(15)-oxidation of hemiketal **75**

Scheme 19.

Completion of the total synthesis of (+)-tedanolide (1)