urge the physician to initiate steroid and antibiotic treatment, in a “preemptive” strike to avoid more severe exacerbation and/or disease progression. Such an approach is obviously of interest but has to be prospectively examined in a randomized study to prove that “early treatment” improves long-term outcome.

The value of CRP for the diagnosis of exacerbations could be judged through a different approach than only looking at sensitivity and specificity. Exacerbations are triggered mainly by viruses. Around 20% of patients with exacerbations are estimated to benefit from antibiotics because only the acquisition of new strains of bacteria is believed to cause bacterial exacerbations (8). CRP levels are highest in bacterial infection (6, 9). Therefore, the fact that a considerable number of patients show normal levels of CRP during exacerbations does not necessarily mean that CRP is a poor biomarker. Low levels of CRP might just reflect that some patients have no evidence of bacterial infection, and exacerbations might be caused by viruses or concomitant heart failure requiring different treatment. Weis and Almdal recently showed that 30% of 166 patients with an exacerbation of COPD showed normal CRP values, whereas median CRP was 45 mg/L in 51 patients reporting increased sputum purulence and 97 mg/L in 51 patients with a concomitant infiltrate (9). The clinical usefulness of a cutoff value of a biomarker depends on the underlying disease and the potential decision for treatment (10). In 243 patients with lower respiratory tract infection, we found that the sensitivity of CRP at a cutoff of 50 mg/L was 93.8% to detect bacterial infection requiring antibiotic treatment (11). Thus, among the blind—despite a moderate sensitivity and specificity—CRP seems to be a one-eyed biomarker for defining an exacerbation. Using one eye in the right direction may still help us to make correct decisions for the treatment of patients with COPD exacerbations.

Any observational study is biased by the choice of the “gold standard,” and for COPD exacerbation this gold standard does not exist. Accordingly, observational studies might overestimate the diagnostic value of a marker. Interventional studies, in which therapy is guided by cutoff ranges of a biomarker and in which efficacy is the outcome measure, have the potential to provide more clinically relevant data. Conceptually, the level of a surrogate marker like CRP mirrors only the likelihood for a disease. Regrettably, Hurst and colleagues did not use their observational experience to suggest specific cutoff ranges for future intervention studies. For procalcitonin, this has been done, and attempts to guide antibiotic therapy in lower respiratory tract infections have shown promising results (12, 13). Results of a recently completed intervention study performed in COPD exacerbations show that antibiotic therapy can safely be guided using a defined cutoff for procalcitonin (14). The time has arrived to move beyond the observational reporting of “promising” biomarkers. Specific cutoff ranges must be proposed and intervention studies conducted. Only this will open both our eyes, and reveal if CRP or other biomarkers can really help us in the routine care of patients with COPD.

Conflict of Interest Statement: Neither author has a financial relationship with a commercial entity that has an interest in the subject of this manuscript.

BEAT MÜLLER, M.D.
MICHAEL TAMM, M.D.
University Hospital, Basel
Basel, Switzerland

References

1. Barnes P, Chowdhury B, Kharitonov SA, Magnussen H, Page CP, Postma D, Saetta M. Pulmonary biomarkers in chronic obstructive pulmonary disease. *Am J Respir Crit Care Med* 2006;174:6–14.
2. Hurst JR, Donaldson GC, Perera WR, Wilkinson TMA, Bilello JA, Hagan GW, Vessey RS, Wedzicha JA. Use of plasma biomarkers at exacerbation of chronic obstructive pulmonary disease. *Am J Respir Crit Care Med* 2006;174:867–874.
3. Man P, Connett JE, Anthonisen NR, Wise RA, Tashkin DP, Sin DD. C-reactive protein and mortality in mild to moderate chronic obstructive pulmonary disease. *Thorax* (In press).
4. Pepys MB, Hirschfield GM, Tennent GA, Gallimore JR, Kahan MC, Bellotti V, Hawkins PN, Myers RM, Smith MD, Polara A, et al. P. Targeting C-reactive protein for the treatment of cardiovascular disease. *Nature* 2006;440:1217–1221.
5. Muller B, Peri G, Doni A, Perruchoud AP, Landmann R, Pasqualini F, Mantovani A. High circulating levels of the IL-1 type II decoy receptor in critically ill patients with sepsis: association of high decoy receptor levels with glucocorticoid administration. *J Leukoc Biol* 2002;72:643–649.
6. Dev D, Wallace E, Sankaran R, Cunniffe J, Govan JR, Wathen CG, Emmanuel FX. Value of C-reactive protein measurements in exacerbations of chronic obstructive pulmonary disease. *Respir Med* 1998;92:664–667.
7. Blum CA, Muller B, Huber P, Kraenzlin M, Schindler C, DeGeyter C, Keller U, Puder DD. Low-grade inflammation and estimates of insulin resistance during the menstrual cycle in lean and overweight women. *J Clin Endocrinol Metab* 2005;90:3230–3235.
8. Sethi S, Evans N, Grant BJ, Murphy TF. New strains of bacteria and exacerbations of chronic obstructive pulmonary disease. *N Engl J Med* 2002;347:465–471.
9. Weis N, Almdal T. C reactive protein can it be used as a marker of infection in patients with exacerbation of chronic obstructive pulmonary diseases. *Eur J Intern Med* 2006;17:88–91.
10. Christ-Crain M, Muller B. Procalcitonin in bacterial infections: hype, hope, more or less? *Swiss Med Wkly* 2005;135:451–460.
11. Stolz D, Christ-Crain M, Gencay M, Bingisser R, Huber PR, Müller B, Tamm M. Diagnostic value of signs, symptoms and laboratory values in lower respiratory tract infection. *Swiss Med Wkly* 2006;136:434–440.
12. Christ-Crain M, Jaccard-Stolz D, Bingisser R, Gencay M, Huber P, Tamm M, Müller B. Effect of procalcitonin-guided treatment on antibiotic use and outcome in lower respiratory tract infections: cluster-randomised, single-blinded intervention trial. *Lancet* 2004;363:600–607.
13. Christ-Crain M, Stolz D, Bingisser R, Muller C, Miedinger D, Huber PR, Zimmerli W, Harbarth S, Tamm M, Muller B. Procalcitonin-guidance of antibiotic therapy in community-acquired pneumonia: a randomized trial. *Am J Respir Crit Care Med* 2006;174:84–93.
14. Stolz D, Christ-Crain M, Bingisser R, Leuppi J, Miedinger D, Muller C, Huber PR, Muller B, Tamm M. Antibiotic treatment of exacerbations of COPD: a randomized controlled trial comparing procalcitonin-guidance with standard therapy. *Chest* (In press)

DOI: 10.1164/rccm.200607-922ED

Just a Spoonful of Technology Makes the Protocol Go Down

Important therapeutic questions in critical care, including questions about fluid management, discontinuation of mechanical ventilation, management of acute lung injury, and nutritional support, do not involve testing a single drug at a fixed dose

against a placebo (1–4). We are interested in evaluating hypotheses such as “routine screening for extubation readiness reduces duration of mechanical ventilation” or “minimizing alveolar distention in acute lung injury reduces mortality.” We do this by

transforming these mechanistic principles into testable, reproducible protocols. Protocols do more than make vague recommendations—they provide precise and reliable instructions that are modified over time in response to data measured from the patient.

Unfortunately, the very aspects of protocols that allow them to test complex treatment strategies make them difficult to evaluate in randomized trials. First, investigators must make multiple, often arbitrary, choices in constructing the protocol. Evidence is rarely available to inform each of the decisions necessary to build even the simplest protocol. When bedside clinicians try to implement the results of these trials or when consensus groups try to make recommendations based on them, they are often faced with questions about which aspects of the protocol evaluated in the study can be modified without sacrificing its efficacy. Second, for feasibility reasons, trials can only test a limited number of different protocols. A winning strategy can only be said to be better than its comparator. Therefore, the control arms of trials of protocols are now scrutinized as carefully as the intervention protocol (5). Third, it is rarely possible to blind clinicians in the study to treatment assignment. Therefore, bias from differential use of co-interventions or assessment of outcomes can never be completely excluded. Given these design limitations, it is not surprising that clinical trials of protocols generate a great deal of debate.

This issue of the *AJRCCM* (pp. 894–900) presents the results of a multicenter, multinational, randomized trial of computer-driven protocolized weaning that reduced median duration of mechanical ventilation by 4.5 d (6). The idea of substituting a computer for a physician in the feedback loop between patient and ventilator is not novel; however, clinical trials demonstrating superiority of this approach are limited (7, 8). This study by Lellouche and colleagues represents the culmination of a research program directed at developing a closed-loop knowledge based system for weaning patients (9). The device incorporates clinical data including end-tidal CO₂, body weight, type of humidification, and a history of the patient's respiratory rate response to varying pressure support levels to gradually lower ventilatory support while keeping patients in a comfort zone.

Control arm aficionados will note that these investigators chose a “usual care” control arm in which individual sites relied on their own written guidelines, principles, and protocols (the authors use these terms almost interchangeably) to discontinue mechanical ventilation. Unfortunately, it is difficult, based on the information provided, to tell exactly how the care differed between the arms. In fact, the computer algorithm and the guidelines at the participating hospitals appear quite similar. Patients in both arms of the trial received nearly identical modes of mechanical ventilation on most days, were *supposed* to receive at least daily spontaneous breathing trials, and received similar (and quite limited) use of sedation after initiation of the weaning phase. However, we do not know how many successful spontaneous breathing trials lead to extubation or how long it took to act on this information in either arm. We don't really know how carefully the sites followed their own ventilator guidelines or how rapidly pressure support levels were reduced. The challenging part of this study and, in fact, the challenging part of any study with a control arm in which usual care varies considerably among sites and patients is to figure out exactly what care was superseded. We know more about the type of care the patients in the control arm were supposed to have received than the care they actually received.

Should we be surprised that a protocolized approach to discontinuing mechanical ventilation performed by a non-physician (in this case a computer) that incorporates limited sedation, gradual reduction in pressure support that maintains patient

comfort, and a prompt to extubate after a successful spontaneous breathing trial reduces duration of mechanical ventilation? Since this is essentially what current guidelines recommend, I suspect few readers will find these results surprising (3). It would appear that the computer was simply more effective at implementing current guidelines than whatever strategies were being used to implement similar strategies at the participating hospitals. Again, unfortunately, we are provided with limited information on whether these hospitals used non-physicians empowered to independently implement their weaning guidelines as recommended by the current literature. I suspect they did not. In their pilot study, the authors suggest that one of the advantages of this device in their hospitals was that it did not require the “presence of a physician at the bedside 24 h a day to apply a protocol” (10). Of course, this ignores the fact that much of the literature on protocolized weaning does not rely on the continuous presence of a physician at the bedside (11). Written guidelines for physicians, as implemented in the study hospitals in the absence of protocolized orders, feedback, or other implementation tools, are known to have little effect on practice (12).

Every year in industrialized nations, hundreds of thousands of people require mechanical ventilation for acute respiratory failure. We have evidence to improve outcomes in these areas with protocols for acute lung injury and weaning, but need strategies to make sure every patient has access to this care. Some physicians may find computer-based protocols more acceptable than sharing responsibility with a respiratory therapist or a nurse (13). Computer-based protocols may be more effective, less expensive, or more reliable than implementation by non-physician clinicians. Hopefully, future studies of this technology will explore these questions with appropriate comparisons and measurements. In the meantime, it is refreshing to see the debate move from *whether* we should have non-physician-driven protocolized mechanical ventilation to *how* we might disseminate and implement these protocols.

Conflict of Interest Statement: G.D.R. has NIH funding to test strategies to implement protocolized ventilation management in the ICU.

GORDON D. RUBENFELD, M.D., M.Sc.
Harborview Medical Center
Seattle, Washington

References

1. Wiedemann HP, Wheeler AP, Bernard GR, Thompson BT, Hayden D, deBoisblanc B, Connors AF Jr, Hite RD, Harabin AL. Comparison of two fluid-management strategies in acute lung injury. *N Engl J Med* 2006;354:2564–2575.
2. The Acute Respiratory Distress Syndrome Network. Ventilation with lower tidal volumes as compared with traditional tidal volumes for acute lung injury and the acute respiratory distress syndrome. *N Engl J Med* 2000;342:1301–1308.
3. MacIntyre NR, Cook DJ, Ely EW Jr, Epstein SK, Fink JB, Heffner JE, Hess D, Hubmayer RD, Scheinhorn DJ. Evidence-based guidelines for weaning and discontinuing ventilatory support: a collective task force facilitated by the American College of Chest Physicians; the American Association for Respiratory Care; and the American College of Critical Care Medicine. *Chest* 2001;120:375S–395S.
4. Martin CM, Doig GS, Heyland DK, Morrison T, Sibbald WJ. Multicentre, cluster-randomized clinical trial of algorithms for critical-care enteral and parenteral therapy (ACCEPT). *CMAJ* 2004;170:197–204.
5. Parshuram CS, Kavanagh BP. Positive clinical trials: understand the control group before implementing the result. *Am J Respir Crit Care Med* 2004;170:223–226.
6. Lellouche F, Mancebo J, Joliet P, Roesler J, Schortgen F, Dojat M, Cabello B, Bouadma L, Rodriguez P, Maggiore S, et al. A multicenter randomized trial of computer-driven protocolized weaning from mechanical ventilation. *Am J Respir Crit Care Med* 2006;174:894–900.

7. Ohlson KB, Westenskow DR, Jordan WS. A microprocessor based feed-back controller for mechanical ventilation. *Ann Biomed Eng* 1982;10:35–48.
8. East TD, Heermann LK, Bradshaw RL, Lugo A, Sailors RM, Ershler L, Wallace CJ, Morris AH, McKinley B, Marquez A, *et al.* Efficacy of computerized decision support for mechanical ventilation: results of a prospective multi-center randomized trial. *Proc AMIA Symp* 1999;251–255.
9. Dojat M, Brochard L, Lemaire F, Harf A. A knowledge-based system for assisted ventilation of patients in intensive care units. *Int J Clin Monit Comput* 1992;9:239–250.
10. Bouadma L, Lellouche F, Cabello B, Taille S, Mancebo J, Dojat M, Brochard L, Lemaire F, Harf A. Computer-driven management of prolonged mechanical ventilation and weaning: a pilot study. A knowledge-based system for assisted ventilation of patients in intensive care units. *Intensive Care Med* 2005;31:1446–1450.
11. Ely EW, Meade MO, Haponik EF, Kollef MH, Cook DJ, Guyatt GH, Stoller JK. Mechanical ventilator weaning protocols driven by non-physician health-care professionals: evidence-based clinical practice guidelines. *Chest* 2001;120:454S–463S.
12. Grimshaw JM, Thomas RE, MacLennan G, Fraser C, Ramsay CR, Vale L, Whitty P, Eccles MP, Matowe L, Shirran L, *et al.* Effectiveness and efficiency of guideline dissemination and implementation strategies. *Health Technol Assess* 2004;8:1–72.
13. Rubenfeld GD, Cooper C, Carter G, Thompson BT, Hudson LD. Barriers to providing lung-protective ventilation to patients with acute lung injury. *Crit Care Med* 2004;32:1289–1293.

DOI: 10.1164/rccm.200607-933ED

Biomass Fuels and Health

The Gap between Global Relevance and Research Activity

The greatest burden on health related to biomass fuels affects the world's poorest and most vulnerable populations. Approximately half of the world's population, and up to 90% of households in rural areas of developing countries, still depend on unprocessed biomass fuels, mainly wood, dung, and crop residues, for cooking and heating (1, 2). These are burned indoors in open fires and stoves that often are not vented. Combustion is incomplete and leads to high indoor concentrations of substances harmful to health, such as respirable particles (PM), carbon monoxide, nitrogen oxides, polycyclic organic hydrocarbons, and other toxic compounds (3). In developing countries, women are traditionally responsible for cooking, and consequently the exposures are highest for them and for their infants and young children, who are often carried on their mother's back. Typically, exposure to high levels of pollutants lasts for 3 to 7 hours daily over many years (2). In global perspective, the magnitude of the health consequences of indoor air pollution, biomass fuels being the major source, is comparable to that of tobacco use, exceeded only by malnutrition, poor water/sanitation, and unsafe sex (4).

What do we know about the health effects of biomass fuels? The known or suggested adverse effects include common respiratory diseases, along with low birth weight and increased infant mortality (2, 3), and even DNA damage (5). Considering the extent of biomass fuel exposure, it is surprising how little research there is to date on its health consequences. At present, the strongest evidence exists for a relation between biomass fuel exposure and acute respiratory infections in children, with effect estimates for incidence and mortality ranging from two- to three-fold (6). Other major respiratory effects that have been linked to biomass fuels more or less consistently include chronic obstructive pulmonary disease, asthma, nasopharyngeal and laryngeal cancer, and susceptibility to pulmonary tuberculosis (2, 3).

In this issue of the *Journal* (pp. 901–905), a study of more than 800 women from a rural community in Mexico investigated the relations between biomass fuel cooking and respiratory symptoms and lung function (7). The study demonstrates that it is feasible to carry out studies applying high-quality research methods in the often-difficult conditions of developing countries. This is important for getting reliable effect estimates and also for being able to assess adequately the benefits from any interventions aiming at solving these problems.

Earlier studies have shown an increase in respiratory symptoms and lung function impairment related to biomass fuel burning in children and in adults (2, 8–13), but there have been some

methodologic issues, such as compromised validity of exposure assessment and limited control for confounding. To address these problems, the Mexican study (7) measured particle concentrations while cooking was in progress with the usual fuel, thus also capturing peak concentrations. In addition, it estimated a cumulative exposure index combining reported daily exposure in hours with years of exposure (hour-years) to assess lifetime biomass fuel exposure. The results showed high PM₁₀ peak concentrations of the order of 2,600 µg/m³ during biomass combustion and large lifetime exposures with the mean cumulative index of 114 hour-years. High PM levels of 500 to 2,500 µg/m³ have been reported also from other developing countries, including Guatemala (14), India (15), and Nepal (13). Peak PM₁₀ over 2,600 µg/m³ among those using biomass fuels was related to small, but significant reductions in FEV₁ (–81 ml), FVC (–122 ml), and FEV₁% predicted (–4.7) (7). The effects were smaller compared with previous hospital-based case-control studies (9–11), but still comparable to the estimates from environmental tobacco smoke exposure in adults (16). Taking into account the extent of biomass fuel exposure in rural communities, the effect is likely to have significant impact on the health of women in developing countries. Questionnaire-reported current exposure and cumulative hour-years were significantly related to occurrence of cough and phlegm production, with a small increase also detected for wheezing in the past. The study also adjusted for potential confounding factors, such as age, passive smoking and socioeconomic status. Income served as a proxy for the latter and is a relevant confounder, since people's possibilities for using cleaner fuels are strongly related to their income.

A limitation of the study is its small reference group of women with exposure to gas only, which reduces the power of this otherwise rather large study. Even more important is the fact that practically all women had previous exposure to biomass fuels, most of them in their childhood. This means that current biomass fuel exposure could not be contrasted to “never exposure,” but instead to “no recent exposure,” so the effect estimates are underestimated. The strongest effects on lung function were detected in relation to high current concentrations of PM₁₀, so how much of the effect is acute and perhaps reversible and how much of it is long-term reduction that cannot be recovered remains somewhat unclear. Distinguishing these two types of effects is important from a preventive point of view and, in the future, development of better assessment methods for long-term exposures and longitudinal assessment of the change in lung