Is the Hygiene Hypothesis an Example of Hormesis?

John A. Bukowski^{1*} and R. Jeffrey Lewis¹

¹*ExxonMobil Biomedical Science, Inc., 1545 Route 22 East, P.O. Box 971, Annandale, NJ 08801-0971*

ABSTRACT

The “hygiene hypothesis” has been suggested to explain the rising incidence of allergic disorders in developed countries. The postulated mechanism is that infectious and/or microbial agents stimulate the immune system toward Th1 (allergy fighting) rather than Th2 (allergy promoting) response. This paper reviews the evidence related to early life infectious/microbial exposures and subsequent atopic disorders and evaluates whether these data suggest a hormetic effect. Our review indicates an insufficient and contradictory association for bacterial/viral infections, with protective effects being either absent or specific to certain infections and/or populations. Chronic, heavy parasitic burdens appear to confer protection against atopic disorders, but are associated with considerable pathology. Moreover, light parasitic burden may increase allergic responses (i.e., no “low dose” beneficial effect). In contrast, there is consistent evidence that general microbial exposures, particularly gut commensals, may be protective against allergy development, which is consistent with a hormetic effect (i.e., potentially beneficial effects at low doses and detrimental effects at high levels). **Conclusion:** General microbial exposures in relation to the “hygiene hypothesis” may represent a hormetic effect, although further research with more rigorous study methods (i.e., prospective designs and measurement of exposure timing, dose, route, etc.) are needed.

Key Words: hormesis, hygiene hypothesis, atopy, allergies, infection, parasites.

INTRODUCTION

In recent years, children from developed countries like the United States and the United Kingdom have experienced large increases in the incidence of asthma and atopic (i.e. inhaled allergic) diseases. It has been suggested that these increases may be attributable to improved “hygienic” conditions typically found in developed

* Corresponding author: ExxonMobil Biomedical Science, Inc., 1545 Route 22 East, P.O. Box 971, Annandale, NJ 08801-0971. (908)730-1105 (telephone). (908)730-1192 (fax). john.a.bukowski@exxonmobil.com

societies, a hypothesis referred to as the “hygiene hypothesis” (Strachan 1989; Lordan and Holgate 1999; Martinez and Holt 1999; Strachan 1999; Abramson and Walters 2000).

The mechanism postulated for the “hygiene hypothesis” is that infectious and/or microbial agents may serve to “educate” the immune system to avoid inappropriate responses to harmless external antigens. The immune system of nonatopic individuals is characterized by a predisposition toward a type-1 helper-T-cell (Th1) response, with production of INF- γ and inhibition of type-2 helper-T-cell (Th2) cells. Conversely, atopic individuals have a predisposition toward a Th2 response and production of cytokines such as IL-4 and IL-5, which promote IgE production and eosinophilia. The newborn immune system also exhibits a Th2 bias. The hygiene hypothesis suggests that microorganism exposures early in life may serve to stimulate the immune system toward a more mature Th1 response, rather than an allergy-promoting Th2 response (Martinez 1994; Romagnani 1996; Martinez and Holt 1999; Openshaw and Walzl 1999; Abramson and Walters 2000).

The concept of hormesis suggests that exposure to low doses of an agent that is detrimental at higher doses may produce a beneficial effect (Sagan 1987; Neafsey 1990; Bukowski and Lewis 2000). Based on this definition, a natural question is whether the “hygiene hypothesis” may represent a hormetic effect. Accordingly, this paper reviews the evidence related to the “hygiene hypothesis” and compares the findings with the concept of hormesis. This paper is not intended as a comprehensive or exhaustive review of the literature. Rather, it is meant as a high-level overview of the evidence on this topic, with the hope of stimulating discussion and awareness. Our review focuses on three types of exposures — bacterial/viral infectious illness, general microbial exposures, and parasitism — and evaluates whether the evidence related to each of these exposures suggests a possible hormetic effect.

BACTERIAL/VIRAL INFECTIONS

Findings from studies of early life exposure to bacterial/viral infections and subsequent risk of atopic disorders have produced variable results. For example, a study of Italian military cadets found that a history of exposure to food-borne and oral-fecal infections (*H. pylori*, *Toxoplasma gondii*, and hepatitis A), but not airborne viruses, was associated with reduced atopy and allergies (Matricardi *et al.* 1997, 2000). Measles infection, an inducer of a Th1 immune response, has been associated with a reduced risk of atopic disorders in some studies (Shaheen *et al.* 1996; Bodner *et al.* 1998; Farooqi and Hopkin 1998) but not others (Paunio *et al.* 2000). A report from Japan suggests that children demonstrating a positive immune response to the bacille Calmette-Gue‘rin (BCG) vaccine (another inducer of type 1 immune response) have less allergenic sensitization and fewer atopic diseases compared with children having a negative response (Shirakawa *et al.* 1997).

Studies having less specific information on type of viral infection have reported similar results. For example, Martinez *et al.* (1995) found that nonwheezing lower

respiratory tract infections (LRI) during the first 3 years of life were associated with lower IgE levels and lower skin test reactivity compared with children with wheezing and/or absent LRI. A longitudinal study of 1314 German children by Illi *et al.* (2001) found that repeated viral infections other than lower respiratory tract infections early in life may reduce risk of subsequent asthma development. Additionally, there is a relatively consistent association between smaller sibship size, (Strachan 1989; von Mutius *et al.* 1994; Bodner *et al.* 1998) higher birth order (Strachan 1989; von Mutius *et al.* 1994; Strachan *et al.* 1997; Bodner *et al.* 1998) and increased risk of atopic disorders. Interestingly, two studies reported a reduced level of IgE in cord blood of pregnancies having higher birth order, suggesting that the sibling effect may originate in utero (Bergmann *et al.* 1999; Karmaus *et al.* 2001). In contrast to the above studies, there is evidence that some respiratory viruses, particularly respiratory syncytical virus (RSV), may cause asthma (Weiss 1998a; Hogg 1999; Gern 2000). Additionally, several studies have found either similar or increased risk of atopy and/or asthma associated with childhood infection with airborne viruses (Bodner *et al.* 1998; Wold 1998; von Mutius *et al.* 1999; Matricardi *et al.* 2000).

Another source of information about the role of childhood infections in the development of atopic diseases is studies of childhood immunizations, antibiotic use and daycare attendance. With the exception of vaccines like the BCG, most vaccinations given to children induce a Th2 response (Rook and Stanford 1998). Several studies suggest that infant and childhood immunizations, particularly pertussis vaccine, may be associated with an increased risk of asthma (Kemp *et al.* 1997; Farooqi and Hopkin 1998). With regard to antibiotics use, several investigators have found that antibiotic use in the first few years of life is associated with an increased risk of developing atopic diseases, with risk increasing as number of antibiotic courses taken increases (Farooqi and Hopkin 1998; von Mutius *et al.* 1999; Wickens *et al.* 1999a). However, several studies have reported either weak or no evidence that infant and childhood immunizations increase risk of atopic diseases (Henderson *et al.* 1999; Nilsson *et al.* 1999; Destefano *et al.* 2002). Moreover, one study found antibiotic use to be a risk factor for asthma and allergic disorders only among children with parents having hay fever (Droste *et al.* 2000). Finally, results for daycare attendance and subsequent asthma are mixed, with some studies reporting a protective effect (Oddy *et al.* 1999) and others reporting either no association (Nystad *et al.* 1999; Kilpelainen *et al.* 2000) or an increased risk (Wickens *et al.* 1999b).

Conclusion. The evidence to support a reduced risk of atopic disorders and early life exposure to bacterial/viral infections is insufficient and contradictory. In general, the effect is either absent or specific to certain infections and/or populations. More importantly, there is no evidence that the response is a low-dose phenomenon — a key characteristic of hormetic effects. As such, if a protective effect of bacterial/viral infections is real, this effect is more appropriately viewed as a beneficial response to a pathological process rather than a hormetic effect.

PARASITISM

As infection and microbial exposure has decreased in Western society over time, so has parasitism. Prior to the 20th century, parasitism was quite common throughout the world, and it is still endemic to most regions of the developing world. Also, as with allergy, parasitism is associated with an alteration in the balance between the Th1 and Th2 responses. These factors have caused some to wonder if the rise in atopy has been fueled by the fall in parasitism (Yazdanbakhsh *et al.* 2001, 2002).

There is a fair body of evidence suggesting that parasitism may protect against allergic illness. Schistosomiasis and a variety of intestinal parasite infections have been associated with lower levels of allergic reactivity (Lynch *et al.* 1998; Araujo *et al.* 2000; van den Biggelaar *et al.* 2000; Nyan *et al.* 2001). Removal of parasites through anthelmintic treatment has also been shown to induce a heightened allergic response (Lynch *et al.* 1993). However, there is also evidence against a protective effect from parasitism, with some studies finding no protective effect on allergy, or even a worsening of allergic disease among parasitized individuals (Catapani *et al.* 1997; Lynch *et al.* 1997; Mao *et al.* 2000).

One apparent inconsistency behind the hypothesis that parasitism protects against allergy is the fact that most parasites elicit a strong Th2 immune response in the host (Weiss 1998b). Several mechanisms have been suggested to explain this seemingly paradoxical finding. One possible mechanism is that the polyclonal IgE response induced by parasitism competitively binds to mast cell receptors, inhibiting allergic degranulation. A second proposed mechanism is that this polyclonal IgE expansion exhausts the immune response. However, the most promising mechanistic explanation is that the inflammatory response produced by chronic parasitism also induces a down-regulation of the immune system (e.g., through interleukin 10 production), as a means of protecting the host from runaway inflammation. This would explain why heavy parasitism seems to be protective for allergy/asthma, while lighter parasite burdens are not. This also explains why anthelmintic treatment removes this protective benefit. In essence, this mechanism suggests that parasitism does not prevent allergy development, only the inflammatory manifestations of allergic disease (Yazdanbakhsh *et al.* 2001, 2002).

Conclusion: The best available evidence supporting a protective effect on allergy from parasite exposure suggests a dampening of the allergic response in those with heavy parasite burdens, which is due to an antiinflammatory mechanism. This is not a low-dose phenomenon, given that heavy parasite burdens are needed. Furthermore, the pathology associated with heavy parasite loads probably outweighs any anti-allergy/asthma benefits. Therefore, such a process, if true, does not represent hormesis.

MICROBIAL EXPOSURES

Humans have evolved in an environment characterized by the presence of non-pathogenic microbial exposures (Rook and Stanford 1998). Only in recent years

have these microbial agents been reduced or eliminated through the introduction of improved food handling and safety procedures, the use of antimicrobial soaps and agents, etc. There is increasing evidence and mechanistic rationale to suggest that early life exposure to generally harmless environmental bacteria may confer protection against development of atopy and asthma.

Endotoxin

Endotoxins are lipopolysaccharides that form a part of the outer structure of the cell wall of Gram-negative bacteria. Interestingly, endotoxin that is typically found in households is capable of inducing INF- γ and IL-12 production (Le *et al.* 1986; D'Andrea *et al.* 1992). Martinez (2001) notes that the body's immune system has a sensitive mechanism to detect the presence of lipopolysaccharides (LPS), which results in the production of a set of cytokines and immune mediators (e.g., INF- γ) by antigen presenting cells. This process may prevent early allergic sensitization, thereby ultimately preventing the development of asthma or allergic conditions.

There is consistent epidemiologic evidence that children raised on farms have a reduced risk of developing atopic disorders compared with children raised in non-farm environments (Braun-Fahrlander *et al.* 1999; Ernst and Cormier 2000; Riedler *et al.* 2000) (Kilpelainen *et al.* 2000; Von Ehrenstein *et al.* 2000; von Mutius *et al.* 2000; Leynaert *et al.* 2001; Riedler *et al.* 2001). The apparent protective effect of being raised on a farm is stronger for hay fever and atopic sensitization than for asthma (Lewis 2000). Currently, it is unclear whether this association may be due to increased microbial exposures resulting from contact with livestock (Riedler *et al.* 2000; Von Ehrenstein *et al.* 2000), consumption of farm milk (Riedler *et al.* 2001), or some other aspect of the traditional farming lifestyle.

Studies have reported that dust samples collected from farming homes have significantly higher levels of endotoxin compared with non-farming homes (von Mutius *et al.* 2000; Gehring *et al.* 2002). Gereda *et al.* (2000) reported significantly lower levels of endotoxin in dust samples collected in the homes of allergen-sensitized infants compared with nonsensitized infants. In this same study, blood T-cells collected from infants living in homes with higher endotoxin levels showed significantly increased INF- γ responses. Braun-Fahrlander *et al.* (2002) reported that endotoxin levels in the dust taken from a child's mattress were inversely related to the occurrence of hay fever, atopic asthma, and atopic sensitization, but not with nonatopic wheeze. An inverse association was also reported between endotoxin levels and the capacity of circulating leukocytes to produce inflammatory cytokines after stimulation with LPS. These findings provide perhaps the strongest evidence for an association between endotoxin exposure and decreased atopic illness (Braun-Fahrlander *et al.* 2002).

While the above data argue that endotoxin may protect against atopic disease, some have pointed out conflicting observations that argue against this hypothesis. For example, exposure to endotoxin in occupational or laboratory settings can produce a number of adverse effects (e.g., fever, respiratory disease, etc.) (Douwes

et al. 2002). However, when examined more closely, these observations are not incompatible with the hygiene hypothesis. Exposed workers (e.g., cotton and grain workers) and subjects in controlled studies experience relatively high-level inhalation exposure to endotoxin, rather than the lower-level exposure to inhaled and ingested endotoxin produced by contact with animals and soil in farm settings. Also, adult exposures occur well after the stimulatory time window suggested by the hygiene hypothesis. Elevated endotoxin levels in metropolitan homes have been associated with exacerbation of existing asthma, but not asthma development (Liu 2002).

Intestinal Flora

Until the 1950s and 1960s, acute infectious diseases of the gastrointestinal tract were a major cause of illness requiring medical treatment in US children, a pattern that still persists today in developing countries. The causes of these illnesses have largely been eliminated in developed countries through vaccinations, better food and water supplies, and increased use of antibiotics (Martinez and Holt 1999). Matricardi and Bonini (2000a; 2000b) and others (Holt *et al.* 1997; Wold 1998) propose that high turnover of microbes (commensal and pathogenic bacteria) at mucosal surfaces, particularly in the gut, may “educate” the immune system to interact safely with nonmicrobial antigens. There is some evidence to support this hypothesis, with several studies indicating compositional differences in gut flora between allergic and nonallergic subjects (Sepp *et al.* 1997; Bjorksten *et al.* 1999; Bottcher *et al.* 2000; Kalliomaki *et al.* 2001a). One of the few prospective studies (Bjorksten *et al.* 2001) found differences in intestinal flora between allergic and nonallergic children in Estonia and Sweden, with allergic children having less lactobacilli and bifidobacteria and more aerobic microorganisms, compared with nonallergic children. Most impressive is a recent double-blind, randomized placebo-controlled trial that showed prenatal administration of lactobacillus GG to mothers of high-risk infants was effective in preventing early atopic disease (Kalliomaki *et al.* 2001b).

One piece of conflicting evidence is that high asthma prevalence in the US is often among urban children who live in environments that are typically thought of as being “dirty”. However, this may reflect a difference in the definition of “dirty” between farm and urban environments (Matricardi and Bonini 2000a). Urban children often live in poorly ventilated homes exposed to cockroaches, vermin, house dust, and volatile chemicals. Farm children are exposed to “dirt” that is rich in organic material.

Conclusion: Considerable evidence supports a protective effect on atopy/asthma from early exposure to microbials, especially those found on farms (e.g., Gram-Negative bacteria and lactobacilli). These low-level beneficial exposures, which are harmful at high levels, suggest a hormetic process.

SUMMARY AND CONCLUSIONS

The evidence to support an association between early bacterial/viral infectious illness and decreased atopy/asthma is insufficient and contradictory. The evidence for such a protective effect from parasitism is stronger, but neither line of evidence suggests a low-level (i.e., hormetic) process. However, considerable evidence does support a protective effect from low-level exposure to general ambient microbials, suggesting that this aspect of the hygiene hypothesis may be an example of hormesis. It should be noted that while evidence supports this aspect of the hygiene hypothesis, the hypothesis still represents a theory that has not been proven. It must also be recognized that asthma/allergy development is complex, with multiple other factors (e.g., fetal development and genetic predisposition), as well as interactions of these factors with the environment, playing an important role (Berg 1996; Tantisira and Weiss 2001; Bukowski *et al.* 2002).

RESEARCH CONSIDERATIONS

This hypothesis, if substantiated, suggests obvious personal or public health interventions that might be useful to decrease the risk of childhood allergy/asthma (e.g., microbial supplementation or decreased emphasis on extensive household disinfection). However, there are currently many unanswered questions regarding the role that early microbial exposures play in the development of allergy and asthma. For example, it is unclear what type, duration, or magnitude of microbial exposure is most associated with a protective effect, although those organisms that induce a strong Th1 response are likely to be good candidates for study (Von Hertzen 1998). Importantly, it is unclear what developmental periods constitute critical “exposure” windows, whether exposures after a certain age (e.g., 3 to 5 years) have an impact, and the role that these exposures play in initiating or maintaining a Th1-type immune response. There is some evidence to suggest that exposures into adulthood may still be important, with several studies suggesting that immigrants may take on the atopy profile of the local population over time (Kalyoncu and Stalenheim 1992). Moreover, atopic disorders have been reported to go into remission after infectious diseases occur among atopic individuals (Serafini 1997).

One controlled trial reported that microbial supplementation during pregnancy may protect against allergy development in children (Kalliomaki *et al.* 2001b). National trials need to be performed to confirm these findings and demonstrate the safety and efficacy of this approach as a possible prophylactic measure.

Finally, most of the studies investigating the “hygiene hypothesis” have been cross-sectional and/or retrospective designs. In order to answer the questions above, larger prospective studies will be needed. National efforts, such as the recently proposed longitudinal National Children’s Study (U.S. Department of Health and Human Services and U.S. Environmental Protection Agency 2001) represent ideal situations within which to evaluate and refine this hypothesis and to identify specific prophylactic exposures.

REFERENCES

- Abramson MJ and Walters EH. 2000. The epidemic of asthma: too much allergen or not enough infection. *Med J Aust* 172:119-121
- Araujo MI, Lopes AA, Medeiros M, Cruz AA, Sousa-Atta L, Sole D, and Carvalho EM. 2000. Inverse association between skin response to aeroallergens and *schistosoma mansoni* infection. *Int Arch Allergy Immunol* 123:145-148.
- Berg RD. 1996. The indigenous gastrointestinal microflora. *Trends Microbiol* 4:430-435
- Bergmann RL, Schulz J, Gunther S, Dudenhausen JW, Bergmann KE, Bauer CP, Dorsch W, Schmidt E, Luck W, and Lau S. 1999. Determinants of cord-blood IgE concentrations in 6401 German neonates. *Allergy* 50:65-71.
- Bjorksten B, Naaber P, Sepp E, and Mikelsaar M. 1999. The intestinal microflora in allergic Estonian and Swedish 2-year old children. *Clin Exper Allergy* 29:342-346.
- Bjorksten B, Sepp E, Julge K, Voor T, and Mikelsaar M. 2001. Allergy development and the intestinal microflora during the first year of life. *J Allergy Clin Immunol* 108:516-520.
- Bodner C, Godden D, and Seaton A. 1998. Family size, childhood infection and atopic diseases. *Thorax* 53:28-32.
- Botthcher M, Sandin A, Norin E, Midtvedt T, and Bjorksten B. 2000. Microflora associated characteristics in faeces from allergic and non-allergic children. *Clin Exp Allergy* 30:590-596.
- Braun-Fahrlander C, Gassner M, Grize L, Neu U, Sennhauser FH, Varonier HS, Vuille JC, and Wuthrich B. 1999. Prevalence of hay fever and allergic sensitization in farmer's children and their peers living in the same rural community. SCARPOL team. Swiss study on childhood allergy and respiratory symptoms with respect to air pollution. *Clin Exp Allergy* 29:28-34..
- Braun-Fahrlander C, Riedler J, Herz U, Eder W, Waser M, Grize L, Maisch S, Carr D, Gerlach F, Bufer A, Lauener RP, Schierl R, Renz H, Nowak D, and von Mutius E. 2002. Environmental exposure to endotoxin and its relation to asthma in school-age children. *N Engl J Med* 347:869-877.
- Bukowski JA and Lewis RJ. 2000. Hormesis and health: a little of what you fancy may be good for you. *South Med J* 93:371-374.
- Bukowski JA, Lewis RJ, Gamble JF, Wojcik NC, and Laumbach RJ. 2002. Range-finding study of risk factors for childhood asthma development and national asthma prevalence. *Hum Ecol Risk Assess* 8:735-765.
- Catapani WR, Pinto PLS, Amato-Neto V, and Mendes E. 1997. Prevalence of allergic diseases in patients with *Schistosomiasis mansoni*. *Allergy Clin Immunol* 100:1764-1765.
- D'Andrea A, Rengaraju M, Valiante NM, Chehimi J, Kubin M, Aste M, Chan SH, Kobayashi M, Young D, and Nickbarg E. 1992. Production of natural killer cell stimulatory factor (interleukin 12) by peripheral blood mononuclear cells. *J Exp Med* 176:1387-1398.
- Destefano F, Gu D, Kramarz P, Truman BI, Iademarco MF, Mullooly JP, Jackson LA, Davis RL, Black SB, Shinefield HR, Marcy SM, Ward JI, and Chen RT. 2002. Childhood vaccinations and risk of asthma. *Pediatr Infect Dis J* 21:498-504.
- Douwes J, Pearce N, and Heederik D. 2002. Does environmental endotoxin exposure prevent asthma? *Thorax* 57:86-90.
- Droste JHJ, Wieringa MH, Weyler JJ, Nelen VJ, Vermeire PA, and Van Bever HP. 2000. Does the use of antibiotics in early childhood increase the risk of asthma and allergic disease? *Clin Exp Allergy* 30:1547-1553.

- Ernst P and Cormier Y. 2000. Relative scarcity of asthma and atopy among rural adolescents raised on a farm. *Am J Respir Crit Care Med* 161:1563–1566.
- Farooqi IS and Hopkin JM. 1998. Early childhood infection and atopic disorder. *Thorax* 53:927–932.
- Gehring U, Bischof W, Fahlbusch B, Wichmann HE, and Heinrich J. 2002. House dust endotoxin and allergic sensitization in children. *Am J Respir Crit Care Med* 166:939–944.
- Gereda JE, Leung DYM, Thatayatikom A, Streib JE, Price MR, Klennert MD, and Liu AH. 2000. Relation between house-dust endotoxin exposure, type 1 T-cell development, and allergen sensitization in infants at high risk of asthma. *Lancet* 355:1680–1683.
- Gern JE. 2000. Viral and bacterial infections in the development and progress of asthma. *J Allergy Clin Immunol* 105:S497–S502.
- Henderson J, North K, Griffiths M, Harvey I, and Golding J. 1999. Pertussis vaccination and wheezing illnesses in young children: prospective cohort study. *Br Med J* 318:1173–1176.
- Hogg JC. 1999. Childhood viral infection and the pathogenesis of asthma and chronic obstructive lung disease. *Am J Respir Crit Care Med* 160:S26–S28.
- Holt PG, Sly PD, and Bjorksten B. 1997. Atopic versus infectious diseases in childhood: a question of balance? *Pediatr Allergy Immunol* 8:53–58.
- Illi S, von Mutius E, Lau S, Bergmann R, Niggemann B, Sommerfeld C, and Wahn U. 2001. Early childhood infectious diseases and the development of asthma up to school age: a birth cohort study. *B Med J* 322:390–395.
- Kalliomaki M, Kirjavainen P, Eerola E, Kero P, Salminen S, and Isolauri E. 2001a. Distinct patterns of neonatal gut microflora in infants developing or not developing atopy. *J Allergy Clin Immunol* 107:129–134.
- Kalliomaki M, Salminen S, Arvilommi H, Kero P, Koskinen P, and Isolauri E. 2001b. Probiotics in primary prevention of atopic disease: a randomised placebo-controlled trial. *Lancet* 357:1076–1079.
- Kalyoncu AF and Stalenheim G. 1992. Serum IgE levels and allergic spectra in immigrants to Sweden. *Allergy* 47:277–280.
- Karmaus W, Arshad H, and Mattes J. 2001. Does the sibling effect have its origin in utero? Investigating birth order, cord blood immunoglobulin E concentration, and allergic sensitization at age 4 years. *Am J Epidemiol* 154:909–915.
- Kemp T, Pearce N, Fitzharris P, Crane J, Fergusson D, St George I, Wickens K, and Beasley R. 1997. Is infant immunization a risk factor for childhood asthma or allergy? *Epidemiology* 8:678–680.
- Kilpelainen M, Terho EO, Helenius H, and Koskenvuo M. 2000. Farm environment in childhood prevents the development of allergies. *Clin Exp Allergy* 30:201–208.
- Le J, Lin JX, Henriksen-Destefano D, and Vilcek J. 1986. Bacterial lipopolysaccharide-induced interferon-gamma production: roles of interleukin 1 and interleukin 2. *J Immunol* 136:4525–4530.
- Lewis SA. 2000. Animals and allergy. *Clin Exp Allergy* 30:153–157.
- Leynaert B, Neukirch C, Jarvis D, Chinn S, Burney P, and Neukirch F. 2001. Does living on a farm during childhood protect against asthma, allergic rhinitis, and atopy in adulthood? *Am J Respir Crit Care Med* 164:1829–1834.
- Liu AH. 2002. Endotoxin exposure in allergy and asthma: reconciling a paradox. *J Allergy Clin Immunol* 109:379–392.
- Lordan JL and Holgate ST. 1999. Recent developments in the pathogenesis and epidemiological trends of asthma. *J R Coll Physicians Lond* 33:418–425.

- Lynch NR, Hagel I, Perez M, Di Prisco MC, Lopez R, and Alvarez N. 1993. Effect of anthelmintic treatment on the allergic reactivity of children in a tropical slum. *J Allergy Clin Immunol* 92:404–411.
- Lynch NR, Hagel IA, Palenque ME, Di Prisco MC, Escudero JE, Corao LA, Sandia JA, Ferreira LJ, Botto C, Perez M, and Le Souef PN. 1998. Relationship between helminthic infection and IgE response in atopic and nonatopic children in a tropical environment. *J Allergy Clin Immunol* 101:217–221.
- Lynch NR, Palenque M, Hagel I, and Di Prisco MC. 1997. Clinical improvement of asthma after anthelmintic treatment in a tropical situation. *Am J Respir Crit Care Med* 156:50–54.
- Mao XQ, Sun DJ, Miyoshi A, Feng Z, Handzel ZT, Hopkin JM, and Shirakawa T. 2000. The link between helminthic infection and atopy. *Parasitol Today* 16:186–187.
- Martinez FD. 1994. Role of viral infections in the inception of asthma and allergies during childhood: could they be protective? *Thorax* 49:1189–1191.
- Martinez FD. 2001. The coming-of-age of the hygiene hypothesis. *Respir Rev* 2:129–132.
- Martinez FD and Holt PG. 1999. Role of microbial burden in aetiology of allergy and asthma. *Lancet* 354:12–15.
- Martinez FD, Stern DA, Wright AL, Taussig LM, and Halonen M. 1995. Association of non-wheezing lower respiratory tract illnesses in early life with persistently diminished serum IgE levels. *Thorax* 50:1067–1072.
- Matricardi PM and Bonini S. 2000a. High microbial turnover rate preventing atopy: a solution to inconsistencies impinging on the hygiene hypothesis? *Clin Exp Allergy* 30:1506–1510.
- Matricardi and Bonini S. 2000b. Mimicking microbial “education” of the immune system: a strategy to revert the epidemic trend of atopy and allergic asthma? *Respir Res* 1:129–132.
- Matricardi PM, Rosmini F, Ferrigno L, Nisini R, Rapicetta M, Chionne P, Stroffolini T, Pasquini P, and D’Amelio R. 1997. Cross sectional retrospective study of prevalence of atopy among Italian military students with antibodies against hepatitis a virus. *Br Med J* 314:999–1007.
- Matricardi PM, Rosmini F, Riondino S, Fortini M, Ferrigno L, Rapicetta M, and Bonini S. 2000. Exposure to foodborne and orofecal microbes versus airborne viruses in relation to atopy and allergic asthma: epidemiological study. *Br Med J* 320:412–417.
- Neafsey PJ. 1990. Longevity hormesis, a review. *Mech Ageing Dev* 51:1–31.
- Nilsson L, Kjellman NIM, and Bjorksten B. 1999. A randomized controlled trial of the effect of pertussis vaccines on atopic disease. *Arch Pediatr Adolesc Med* 152:734–738.
- Nyan OA, Walraven GEL, Banya WAS, Milligan P, van der Sande M, Ceesay SM, Del Prete G, and McAdam KPWJ. 2001. Atopy, intestinal helminth infection and total serum IgE in rural and urban adult Gambian communities. *Clin Exp Allergy* 31:1672–1678.
- Nystad W, Skrandal A, and Magnus P. 1999. Day care attendance, recurrent respiratory tract infections and asthma. *Int J Epidemiol* 28:882–887.
- Oddy WH, Holt PG, Sly PD, Read AW, Landau LI, Stanley FJ, Kendall GE, and Burton PR. 1999. Association between breast feeding and asthma in 6-year-old children: findings of a prospective birth control study. *B Med J* 319:815–819.
- Openshaw PJM and Walzl G. 1999. Infections prevent the development of asthma — true, false or both? *J Royal Soc Med* 92:495–499.
- Paunio M, Heinonen OP, Virtanen M, Leinikki P, Patja A, and Peltola H. 2000. Measles history and atopic diseases: a population-based cross-sectional study. *J Am Med Assoc* 283:343–346.

Hygiene Hypothesis and Hormesis

- Riedler J, Braun-Fahrlander C, Eder W, Schreuer M, Waser M, Maisch S, Carr D, Schjeri R, Nowak D, and von Mutius E. 2001. Exposure to farming in early life and development of asthma and allergy: a cross-sectional survey. *Lancet* 358:1129–1133.
- Riedler J, Eder W, Oberfeld G, and Schreuer M. 2000. Austrian children living on a farm have less hay fever, asthma and allergic sensitization. *Clin Exp Allergy* 30:194–200.
- Romagnani S. 1996. Induction of Th1 and Th2 responses: a key role for the “natural” immune response? *Immunol Today* 13:379–381.
- Rook GAW and Stanford JL. 1998. Give us this day our daily germs. *Immunol Today* 19:113–116.
- Sagan LA. 1987. What is hormesis and why haven’t we heard about it before? *Health Physics* 52:521–525.
- Sepp E, Julge K, Vasar M, Naaber P, Bjorksten B, and Mikelsaar M. 1997. Intestinal microflora of Estonia and Swedish infants. *Acta Paediatr* 86:956–961.
- Serafini U. 1997. Do infections protect against asthma and atopy? *Allergy* 52:955–997.
- Shaheen SO, Aaby P, Hall AJ, Barker DJ, Heyes CB, Shiell AW, and Goudiaby A. 1996. Measles and atopy in Guinea-Bissau. *Lancet* 347:1792–1796.
- Shirakawa T, Enomoto T, Shimazu S, and Hopkin JM. 1997. The inverse association between tuberculin responses and atopic disorder. *Science* 275:77–79.
- Strachan DP. 1989. Hay fever, hygiene, and household size. *Br Med J* 299:1259–1260.
- Strachan DP. 1999. Family size, infection and atopy: the first decade of the “hygiene hypothesis”. *Thorax* 55:554–558.
- Strachan DP, Harkins LS, Johnston ID, and Anderson HR. 1997. Childhood antecedents of allergic sensitization in young British adults. *J Allergy Clin Immunol* 99:6–12.
- Tantisira KG and Weiss ST. 2001. Childhood infections and asthma: at the crossroads of the hygiene and Barker hypothesis. *Respir Res* 2:324–327.
- U.S. Department of Health and Human Services, and U.S. Environmental Protection Agency. 2001. *The National Children’s Study. November 2001. Web Page. Available at: <http://www.nichd.nih.gov/despr/cohort/>.*
- van den Biggelaar AHJ, van Ree R, Rodrigues LC, Lell B, Deelder AM, Kremsner PG, and Yazdanbakhsh M. 2000. Decreased atopy in children infected with *schistosoma haematobium*: a role for parasite-induced interleukin-10. *Lancet* 356:1723–1727.
- Von Ehrenstein OS, von Mutius E, Illi S, Baumann L, Bohm O, and von Kries R. 2000. Reduced risk of hay fever and asthma among children of farmers. *Clin Exp Allergy* 30:187–193.
- Von Hertzen LC. 1998. The hygiene hypothesis in the development of atopy and asthma—still a matter of controversy? *Q J Med* 91:767–771.
- von Mutius E, Braun-Fahrlander C, Schierl R, Riedler J, Ehlermann S, Maisch S, Waser M, and Nowak D. 2000. Exposure to endotoxin or other bacterial components might protect against the development of atopy. *Clin Exp Allergy* 30:1230–1234.
- von Mutius E, Illi S, Hirsch T, Leupold W, Keil U, and Weiland SK. 1999. Frequency of infections and risk of asthma, atopy and airway hyperresponsiveness in children. *Eur Respir J* 14:4–11.
- von Mutius E, Martinez FD, Fritsch C, Nicolai T, Reitmeir P, and Thiemann HH. 1994. Skin test reactivity and number of siblings. *Br Med J* 308:692–695.
- Weiss ST. 1998a. Environmental risk factors in childhood asthma. *Clin Exp Allergy* 28:29–34.
- Weiss ST. 1998b. The relationship of parasites to asthma. *Clin Exp Allergy* 28:38–39.

- Wickens K, Pearce N, Crane J, and Beasley R. 1999a. Antibiotic use in early childhood and the development of asthma. *Clin Exp Allergy* 29:766–771.
- Wickens KL, Crane J, Kemp TJ, Lewis SJ, D’Souza WJ, Sawyer GM, Stone ML, Tohill SJ, Kennedy JC, Slater TM, and Pearce NE. 1999b. Family size, infections, and asthma prevalence in New Zealand children. *Epidemiology* 10:699–705.
- Wold AE. 1998. The hygiene hypothesis revised: is the rising frequency of allergy due to changes in the intestinal flora? *Allergy* 53(Suppl 46):20–25.
- Yazdanbakhsh M, Kremsner PG, and van Ree R. 2002. Allergy, parasites, and the hygiene hypothesis. *Science* 296:490–494.
- Yazdanbakhsh M, van den Biggelaar A, and Maizels RM. 2001. Th2 responses without atopy: immunoregulation in chronic helminth infections and reduced allergic disease. *Trends Immunol* 22:372–377.