Published in final edited form as:

Immunol Cell Biol. 2009 January ; 87(1): 46–49. doi:10.1038/icb.2008.88.

Re-entry of mature T cells to the thymus: an epiphenomenon?

Jonathan Sprent¹ and Charles D Surh²

¹The Garvan Institute of Medical Research, Sydney, Australia

²The Scripps Research Institute, San Diego, USA

Abstract

In certain situations mature post-thymic T cells are able to leave their residence in the secondary lymphoid tissues and re-enter the thymus. The physiological significance of this phenomenon is discussed.

The pool of mature T cells

Typical mature T cells expressing $\alpha\beta$ T cell receptor (TCR) molecules arise in the thymus and are released into the periphery as a mixture of mature CD4 and CD8 cells¹. These cells are immunologically naïve and, in mice, are marked by low (lo) expression of CD44 and high (hi) levels of CD62L (L-selectin) and chemokine receptor CCR7^{2–7}. Naïve T cells live for prolonged periods in interphase and recirculate continuously between blood and lymph through defined regions of the secondary lymphoid tissues, ie spleen, lymph nodes (LN) and Peyer's patches. T cell entry to these organs is highly specific and, for LN, involves sequential contact between CD62L and CCR7 on T cells interacting with vascular addressins and CCL19, 21, respectively, expressed on the surface of high endothelial venules. Homing of naïve T cells to nonlymphoid tissues is generally thought to be minimal, although this view has recently been contested⁸.

Contact with foreign antigen in the spleen and LN drives naïve T cells to proliferate extensively and differentiate into activated effector cells^{4, 9, 10}. These cells tend to downregulate CD62L and CCR7 expression but synthesize a new range of homing molecules which enables the cells to migrate into nonlymphoid tissues. The receptor/ligand interactions involved in migration to nonlymphoid tissues are highly complex^{6, 7}. Expression of CLA and CCR4 is important for activated T cell entry to skin, while several other homing molecules, including $\alpha 4\beta 7$, $\alpha 4\beta 1$ (VLA-4) and CCR9, control migration to the gut. CD44 is involved in T cell entry to the liver while CCR5 is important for migration to a number of nonlymphoid sites, including sites of inflammation involved in autoimmune disease.

At the end of the immune response, elimination of the pathogen concerned causes most effector cells to die rapidly, leaving only a few cells to survive and become long-lived memory cells^{4, 9, 10}. Most of these cells revert to resting cells and reacquire expression of CD62L and CCR7, thus allowing the cells to re-join the recirculating lymphocyte pool and cease to migrate through nonlymphoid sites. Nevertheless, memory cells differ from naïve cells in retaining the CD44^{hi} phenotype of effector cells. T cells with the same phenotype as memory cells are found naturally in normal unimmunised mice. These memory-phenotype (MP) cells comprise a small proportion (10–20%) of T cells in young mice but become a dominant population in old age. MP cells are presumed to be the descendants of cells responding to a variety of foreign and self antigens.

T cell migration to the thymus

As mentioned above, normal resting naïve and memory cells reside within the confines of the recirculating lymphocyte pool, migration to nonlymphoid sites being minimal. In the case of the thymus, early studies showed that injection of mice with mature lymphocytes caused almost none ($<0.1\%$) of the donor cells to enter the thymus^{11–14}, thus showing that this organ is not part of the recirculating lymphocyte pool¹⁵. These and other findings have consolidated the view that the thymus is a “primary” lymphoid organ: once released from the thymus as newly-formed cells, mature T cells rarely if ever return to the thymus. Subsequent studies have confirmed that “back-migration” of normal T cells to the thymus is extremely limited. Nevertheless, parabiosis experiments have shown that some normal T cells do reenter the thymus. Thus, after establishing parabiosis, around 0.3% of total cells in the thymus were partner derived at day 14 and were almost entirely situated in the medulla¹⁶; partner-derived cells were not seen in the cortex until day 21, indicating that the cells in the medulla at day 14 were mature T cells derived from the circulation rather than from cortical stem cells.

Significant migration of mature T cells to the thymus is also detectable when very large numbers of T cells are injected into normal mice. Thus, studies in which allotype-marked T cells were detected in tissue sections after intravenous (IV) injection of 1×10^8 total LN cells showed that about 0.2% of cells in the thymic medulla were of donor origin¹⁷; in the cortex, by contrast, donor cells were barely detectable, ie 0.01% or less. Significantly, the cells in the medulla remained at constant levels for at least 24 weeks. Since the proportion of donor cells in LN declined 4-fold during this time, the data implied that the cells reaching the host thymus stayed there and were not part of the recirculating pool.

In the latter study, it is of interest that most of the donor cells in the medulla had a CD62L^{lo} phenotype, suggesting that migration to the thymus was restricted to a small subset of cells with an activated phenotype. In fact, prior experiments with rat T cell lines had shown that, after IV injection, a surprisingly high proportion, about 1%, of the injected cells reached the host thymus within 3 days and remained there for at least 2 months¹⁸; migration to the thymus depended on antigen activation of the T cell lines just prior to injection. These findings were supported by studies in mice on T cell responses to transplantation antigens¹⁹. Here, it was shown that some of the responding T cells were activated in spleen or LN and then migrated into the thymus.

Quantitative evidence on this issue came from a comparison of thymic migration by normal T cells versus activated T cells after IV injection²⁰. To enhance detection of allotype-marked donor cells in the host thymus, thymic suspensions were depleted of immature HSA^{hi} cells prior to FACS analysis, thus enriching for the small subset (5%) of mature medullary thymocytes. When normal small resting T cells were injected, the % of donor cells in the mature component of host thymocytes was undetectable ($<0.01\%$) even with injection of large numbers of cells, ie 4×10^7 . However, significant migration to the thymus occurred following injection of activated T cells (which were obtained as T blast cells from thoracic duct lymph of irradiated H-2-different recipients of naïve T cells). With injection of large numbers of syngeneic activated T cells, about 0.5% of HSA⁺ thymocytes were of donor origin; when the donor blasts were labelled with ¹²⁵I-5-iodo-2' deoxyuridine, a DNA precursor, the % of donor cells in thymus suspensions reached 5%, suggesting that thymic homing was skewed to cycling cells. In sections, the cells reaching the thymus were largely restricted to the medulla and remained there for at least 1 month.

Additional evidence that thymic homing is largely restricted to activated T cells came from studies in rats. Thus, homing of mature T cells to the rat thymus was very low with CD45RC⁺ cells, ie cells with the phenotype of resting naïve and memory cells, but clearly

detectable with CD45RC⁺ cells, which include activated effector cells as well as resting cells²¹. In a follow up study, however, appreciable thymic homing was observed with CD45RC⁺ cells, leading the authors to conclude that thymic homing can also apply to small resting T cells²². In this study it is notable that the medulla of the rat strain used contained a high proportion of non-T cells, including 12% B cells, implying that ... "the thymic medulla shares many features of peripheral lymphoid organs such as lymph nodes" ²². This is not the case in the mouse where the proportion of B cells in the thymus is normally very low. However, certain mouse strains prone to autoimmune disease, eg (NZB × SJL)F₁ mice, do have features of secondary lymphoid tissues, and mature T and B cells readily enter the thymus of these mice²³. This situation also applies to the preneoplastic thymus of AKR mice – see below.

The immunodeficient thymus

All of the above data refer to thymic homing in normal adult animals. When immunodeficient hosts are used, the results are somewhat different. If the cellularity of the thymus is reduced by >95% following heavy irradiation or treatment with cortisone, migration of naïve LN T cells into the host thymus after adoptive transfer is very low^{20, 24}, indicating that acute depletion of cells does not make the thymus more permeable to circulating T cells. When the thymus is congenitally acellular, by contrast, peripheral T cells readily enter the thymus. Thus, after IV injection of 4×10^7 LN T cells into SCID mice, 1–2% of cells in thymocyte suspensions were of donor origin²⁵. Although the SCID thymus contains very few lymphoid cells, total yields of donor cells in the thymus after IV injection were about 30-fold higher for SCID hosts than for normal hosts. Nearly all of the donor cells recovered from the SCID thymus had the phenotype of typical naïve T cells and comprised a mixture of CD4 and CD8 cells. These findings indicate therefore that naïve T cells are excluded from entering the normal thymus but do enter the atrophic thymus of congenitally T-deficient SCID mice. Similar findings were subsequently reported for other congenitally-deficient mice, namely RAG-2^{-/-} mice, following parabiosis to normal mice²⁶.

Whereas the thymus of SCID and RAG-deficient mice show a drastic (>98%) reduction in total numbers of lymphoid cells, the thymus of mice deficient in pre-T α -which guides early thymocyte differentiation into CD4+CD8⁺ cells – contains significant numbers of T cells (10% of normal), mostly of the $\gamma\delta$ lineage²⁷. When pre-T α ^{-/-} mice were parabiosed to normal mice for 1 week, the remarkable finding was that around 50% of mature CD4 and CD8 T cells in the pre-T α ^{-/-} thymus were derived from the normal partner strain²⁸; most of the immigrant cells were naïve CD44^{lo} cells. In a follow up study, IV injection of 2×10^7 LN T cells led to the appearance of 2% donor cells in thymocyte suspensions from pre-T α ^{-/-} mice but only 0.001% donor cells in thymocytes from normal mice²⁶. Taking into consideration the differing size of the thymuses, it was calculated that the total number of immigrant mature T cells was about 1×10^5 cells/thymus and was no higher in the lymphopenic pre-T α ^{-/-} thymus than in the normal thymus. The implication therefore is that, in contrast to the SCID thymus (see above), the lymphopenic thymus of pre-T α ^{-/-} mice is not more permeable to circulating mature T cells than the normal thymus. In this study, it should be mentioned that, based on the results of injecting a dye, CFSE, into the spleen and then searching for labelled cells in the thymus, about 50% of the immigrant T cells in the normal thymus were found to be naïve CD44^{lo} cells. This finding is surprising and seemingly at odds with the conclusion from the earlier studies mentioned above that the normal thymus is impermeable to naïve resting naïve T cells.

The neonatal and aged thymus

The evidence that circulating naïve T cells are generally excluded from re-entering the thymus applies to normal adult animals. The situation in neonates is quite different. Thus, studies in which newborn mice were injected IV with LN T cells enriched for resting cells showed that

donor cells comprised a high proportion, 1–5%, of the component of HSA⁺ cells in the host thymus, compared with only 0.02% for donor cells reaching the thymus of adult control mice²⁹. In neonatal hosts, the donor cells comprised a mixture of naïve CD44^{lo} CD4 and CD8 cells and localized largely in the medulla where they remained for at least 2 months.

As in the neonate, T cells readily enter the thymus of aged mice³⁰. After IV injection of naïve T cells (TCR transgenic CD8 cells), localization of the donor cells in the host thymus was about 10-fold higher for old (1.5 years) mice as hosts than for young mice. These findings applied to aged C57BL/6 mice. For AKR mice, ie mice prone to developing thymic lymphomas, the preneoplastic hyperplastic thymus becomes permeable to circulating T and B cells, and these cells lodge in the medulla in substantial numbers after IV injection^{31, 32}. Migration of mature lymphocytes into the AKR thymus increases considerably with age until the mice eventually develop lymphoma.

Discussion

As a whole, the above data indicate that mature T cell traffic in the normal adult thymus is unidirectional: after export from the thymus, naïve T cells circulate in the peripheral lymphoid tissues for prolonged periods but rarely if ever return to the thymus. This scenario is in keeping with the view that the thymus is a primary lymphoid organ. As discussed above, however, the “gate” to the thymus is partly ajar in the neonatal period and old age, and also in certain pathological states in adults, especially in mice with congenital lymphopenia. Why naïve T cells return to the thymus in these situations is unclear. In the case of the normal adult thymus, the failure of naïve T cells to re-enter the thymus presumably reflects that the small blood vessels of this organ do not express sufficient levels of the “homing” ligands such as vascular addressins (PNAd and MadCAM-1) and/or certain chemokines that naïve T cells use for their recirculation through secondary lymphoid tissues. It would seem likely therefore that permeability of the thymus to circulating mature T cells reflects increased expression of these homing ligands.

Although direct information on this issue is sparse, it is of interest that the increased permeability of the preneoplastic AKR thymus for mature T cells with age correlates with a progressive increase in the expression of PNAd and MadCAM-1 on small blood vessels in the thymic medulla³². Similar increased expression of these vascular addressins is seen in the peri-islet lymphocyte infiltrations of pre-diabetic NOD mice³³; though enriched for MP cells, the islets of NOD mice contain substantial numbers of typical naïve T cells, implying that the influx of these cells reflects de novo synthesis of vascular addressins. Whether this scenario is also applicable to T cell homing to the thymus of normal neonates and mice with congenital lymphopenia has yet to be resolved. For neonates, it is of interest that, in mice, the expression of vascular addressins is different at birth, at least in LN³⁴; moreover, both for sheep³⁵ and humans³⁶, the pattern of blood-to-lymph recirculation of naïve T cells during the perinatal period differs from adults in showing prominent migration through the skin and other nonlymphoid tissues. For mice, the permeability of the neonatal thymus to circulating T cells correlates with the medulla being relatively immature at birth and occupied largely by $\gamma\delta$ cells rather than $\alpha\beta$ cells^{34, 37}. The medulla is also poorly formed in congenitally lymphopenic SCID mice, but is rapidly reconstituted after an influx of mature T cells following IV injection²⁵. It is conceivable therefore that a poorly formed medulla somehow makes the thymus leaky and allows entry of naïve T cells. Direct evidence on this issue is lacking.

For the normal thymus, re-entry of mature T cells from the circulation is largely restricted to activated cells and certain subsets of memory cells. As discussed earlier, these cells express high levels of a variety of selectins, selectin ligands, integrins and chemokine receptors, but which of these newly-expressed homing molecules allow the cells to localize in the thymus is

uncertain. However, there could be parallels with the recruitment of thymic stem cells. In this respect, the normal thymus is known to express low but significant levels of several vascular addressins and selectins^{38, 39}, and interaction with these ligands is considered to guide stem cell entry into the thymus. Such entry involves a number of counter-receptors on stem cells, including CD44^{40, 41}, PSGL-1³⁹, $\alpha 4$ integrins⁴¹, as well as chemokine receptors such as CXCR4 and CCR5⁴². These and other homing/adhesion molecules are highly expressed on subsets of activated T cells and are probably largely responsible for controlling the entry of these cells into the normal thymus. Direct evidence on this issue is sparse, however, although T cell homing to the rat thymus is reported to involve $\alpha 4$ integrin-VCAM-1 interaction²¹.

The physiological significance of T cell re-entry to the thymus is a matter for speculation. One possibility is that the antigens on immigrant T cells contribute to tolerance induction. We became interested in this possibility when injection of newborn mice with allogeneic naïve CD8 T cells was found to cause profound intrathymic tolerance to the foreign Mtv antigens on the T cells⁴³. The implication was that the newborn thymus is directly permeable to naïve T cells, which we then proved in later experiments – see above. But the tolerance model used in these experiments was highly contrived and of dubious relevance to tolerance induction to self antigens. Self tolerance is thought to be controlled through intrathymic contact with epithelial cells and resident DC, plus immigrant DC from the periphery, and it is difficult to envisage how tolerance could be further improved by exposure to antigens expressed on immigrant mature T cells. Nevertheless, as for Mtv antigens, it is possible that migration of antigen-bearing T cells into the thymus could contribute to acquired tolerance to foreign antigens, especially alloantigens^{44, 45}.

Another suggestion for T cells returning to the thymus is that these cells may contribute to positive selection. In fact, direct evidence in favour of this idea has come from studies in which the ligand controlling positive selection in the thymus was expressed selectively on immigrant mature T cells²⁶. Under normal conditions, however, positive selection is known to be controlled via immature T cell contact with the rich network of epithelial cells in the cortex, whereas most immigrant T cells settle in the medulla with only very small numbers reaching the cortex. Hence, it seems doubtful whether positive selection mediated by T cells in experimental models is physiologically relevant.

A third possibility is that migration of mature T cells into the thymus plays a role in the development of CD4⁺CD25⁺Foxp3⁺ T regulatory cells (Tregs). These cells are heavily dependent on IL-2 for their survival in the periphery and also seem to require contact with IL-2 for their formation in the thymus^{46–49}. Thus, numbers of Tregs in the thymus are significantly reduced in the absence of IL-2. What then is the source of IL-2 in the thymus? In considering this question, it is notable that Tregs are undetectable in the mouse thymus until a few days after birth^{50, 51}. Since T cells released from the neonatal thymus undergo substantial proliferation and activation in the periphery, presumably to foreign antigens⁵², one can envisage a scenario where some of the activated T cells rapidly re-enter the thymus; here, the immigrant cells release IL-2 locally, thus promoting the development of Tregs. In addition to IL-2, these migrant cells might be needed to release other γc cytokines in the thymus. Thus, Treg development is only partly reduced (by 2-fold) in the absence of IL-2 but is undetectable for $\gamma c^{-/-}$ thymocytes⁴⁶. There is also evidence that Treg formation requires intrathymic contact with thymic stromal derived lymphopoietin (TSLP) made by medullary epithelial cells⁵⁰. Since the medulla is poorly formed at birth (see above), re-entry of activated T cells to the thymus during this period might also be needed to promote formation of the medulla and induce synthesis of TSLP. This scenario might not be applicable to humans because Treg development is evident in the fetus at the end of the first trimester and presumably occurs in a relatively antigen-free environment^{53, 54}.

The least interesting explanation for backmigration of T cells to the thymus is that such migration is merely an epiphenomenon: after appropriate activation by specific antigen and expression of new homing molecules, activated T cells can migrate to almost any site in the body, including the thymus, to mediate their effector function. Precisely where the cells localize, the thymus or elsewhere, is largely an accident. Hopefully, future experiments will come up with an interesting new idea to dispel this gloomy scenario.

References

1. Starr TK, Jameson SC, Hogquist KA. Positive and negative selection of T cells. *Annual review of immunology* 2003;21:139–176.
2. Sprent J. Lifespans of naive, memory and effector lymphocytes. *Current opinion in immunology* 1993;5:433–438. [PubMed: 8347304]
3. Butcher EC, Picker LJ. Lymphocyte homing and homeostasis. *Science* (New York, N.Y. 1996;272:60–66.
4. Sprent J, Surh CD. T cell memory. *Annual review of immunology* 2002;20:551–579.
5. Nourshargh S, Marelli-Berg FM. Transmigration through venular walls: a key regulator of leukocyte phenotype and function. *Trends in immunology* 2005;26:157–165. [PubMed: 15745858]
6. Mora JR, von Andrian UH. T-cell homing specificity and plasticity: new concepts and future challenges. *Trends in immunology* 2006;27:235–243. [PubMed: 16580261]
7. Marelli-Berg FM, Cannella L, Dazzi F, Mirenda V. The highway code of T cell trafficking. *The Journal of pathology* 2008;214:179–189. [PubMed: 18161751]
8. Cose S, Brammer C, Khanna KM, Masopust D, Lefrancois L. Evidence that a significant number of naive T cells enter non-lymphoid organs as part of a normal migratory pathway. *European journal of immunology* 2006;36:1423–1433. [PubMed: 16708400]
9. Kaeck SM, Wherry EJ, Ahmed R. Effector and memory T-cell differentiation: implications for vaccine development. *Nature reviews* 2002;2:251–262.
10. Harty JT, Badovinac VP. Shaping and reshaping CD8+ T-cell memory. *Nature reviews* 2008;8:107–119.
11. Gesner BM, Gowans JL. The fate of lethally irradiated mice given isologous and heterologous thoracic duct lymphocytes. *British journal of experimental pathology* 1962;43:431–440. [PubMed: 13898114]
12. Ford CE, Micklem HS. The thymus and lymph-nodes in radiation chimaeras. *Lancet* 1963;1:359–362. [PubMed: 13958695]
13. Zatz MM, Lance EM. The distribution of chromium 51-labelled lymphoid cells in the mouse. A survey of anatomical compartments. *Cellular immunology* 1970;1:3–17. [PubMed: 5523574]
14. Sprent J. Circulating T and B lymphocytes of the mouse. I. Migratory properties. *Cellular immunology* 1973;7:10–39. [PubMed: 4540428]
15. Gowans JL, Knight EJ. The Route of Re-Circulation of Lymphocytes in the Rat. *Proceedings of the Royal Society of London. Series B, Containing papers of a Biological character* 1964;159:257–282.
16. Hirokawa K, Utsuyama M, Sado T. Immunohistological analysis of immigration of thymocyte-precursors into the thymus: evidence for immigration of peripheral T cells into the thymic medulla. *Cellular immunology* 1989;119:160–170. [PubMed: 2646025]
17. Michie SA, Kirkpatrick EA, Rouse RV. Rare peripheral T cells migrate to and persist in normal mouse thymus. *The Journal of experimental medicine* 1988;168:1929–1934. [PubMed: 2903215]
18. Naparstek Y, Holoshitz J, Eisenstein S, et al. Effector T lymphocyte line cells migrate to the thymus and persist there. *Nature* 1982;300:262–264. [PubMed: 6183591]
19. Fink PJ, Bevan MJ, Weissman IL. Thymic cytotoxic T lymphocytes are primed in vivo to minor histocompatibility antigens. *The Journal of experimental medicine* 1984;159:436–451. [PubMed: 6607314]
20. Agus DB, Surh CD, Sprent J. Reentry of T cells to the adult thymus is restricted to activated T cells. *The Journal of experimental medicine* 1991;173:1039–1046. [PubMed: 2022918]

21. Bell EB, Sparshott SM, Ager A. Migration pathways of CD4 T cell subsets in vivo: the CD45RC-subset enters the thymus via alpha 4 integrin-VCAM-1 interaction. *International immunology* 1995;7:1861–1871. [PubMed: 8580084]
22. Westermann J, Smith T, Peters U, et al. Both activated and nonactivated leukocytes from the periphery continuously enter the thymic medulla of adult rats: phenotypes, sources and magnitude of traffic. *European journal of immunology* 1996;26:1866–1874. [PubMed: 8765033]
23. Dumont FJ, Barrois R, Jacobson EB. Migration of peripheral T and B cells into the thymus of aging (NZB X SJL)F1 female mice. *Cellular immunology* 1984;83:292–301. [PubMed: 6607128]
24. Kadish JL, Basch RS. Hematopoietic thymocyte precursors. III. A population of thymocytes with the capacity to return ("home") to the thymus. *Cellular immunology* 1977;30:12–24. [PubMed: 301062]
25. Surh CD, Ernst B, Sprent J. Growth of epithelial cells in the thymic medulla is under the control of mature T cells. *The Journal of experimental medicine* 1992;176:611–616. [PubMed: 1500862]
26. Kirberg J, Bosco N, Deloulme JC, Ceredig R, Agenes F. Peripheral T lymphocytes recirculating back into the thymus can mediate thymocyte positive selection. *J Immunol* 2008;181:1207–1214. [PubMed: 18606674]
27. Fehling HJ, Krotkova A, Saint-Ruf C, von Boehmer H. Crucial role of the pre-T-cell receptor alpha gene in development of alpha beta but not gamma delta T cells. *Nature* 1995;375:795–798. [PubMed: 7596413]
28. Bosco N, Agenes F, Rolink AG, Ceredig R. Peripheral T cell lymphopenia and concomitant enrichment in naturally arising regulatory T cells: the case of the pre-Talpha gene-deleted mouse. *J Immunol* 2006;177:5014–5023. [PubMed: 17015684]
29. Surh CD, Sprent J, Webb SR. Exclusion of circulating T cells from the thymus does not apply in the neonatal period. *The Journal of experimental medicine* 1993;177:379–385. [PubMed: 8426109]
30. Hale JS, Boursalian TE, Turk GL, Fink PJ. Thymic output in aged mice. *Proceedings of the National Academy of Sciences of the United States of America* 2006;103:8447–8452. [PubMed: 16717190]
31. Michie SA, Rouse RV. Traffic of peripheral B and T lymphocytes to hyperplastic, preneoplastic thymuses of AKR mice. *The American journal of pathology* 1991;138:1015–1025. [PubMed: 2012169]
32. Michie SA, Streeter PR, Butcher EC, Rouse RV. L-selectin and alpha 4 beta 7 integrin homing receptor pathways mediate peripheral lymphocyte traffic to AKR mouse hyperplastic thymus. *The American journal of pathology* 1995;147:412–421. [PubMed: 7543735]
33. Faveeuw C, Gagnerault MC, Lepault F. Expression of homing and adhesion molecules in infiltrated islets of Langerhans and salivary glands of nonobese diabetic mice. *J Immunol* 1994;152:5969–5978. [PubMed: 8207221]
34. Mebius RE, Streeter PR, Michie S, Butcher EC, Weissman IL. A developmental switch in lymphocyte homing receptor and endothelial vascular addressin expression regulates lymphocyte homing and permits CD4+ CD3- cells to colonize lymph nodes. *Proceedings of the National Academy of Sciences of the United States of America* 1996;93:11019–11024. [PubMed: 8855301]
35. Cahill RN, Kimpton WG, Washington EA, Holder JE, Cunningham CP. The ontogeny of T cell recirculation during foetal life. *Seminars in immunology* 1999;11:105–114. [PubMed: 10329497]
36. Zippelius A, Bioley G, Le Gal FA, et al. Human thymus exports naive CD8 T cells that can home to nonlymphoid tissues. *J Immunol* 2004;172:2773–2777. [PubMed: 14978076]
37. Farr A, Hosier S, Nelson A, Itohara S, Tonegawa S. Distribution of thymocytes expressing gamma delta receptors in the murine thymus during development. *J Immunol* 1990;144:492–498. [PubMed: 2136893]
38. Lepique AP, Palencia S, Irjala H, Petrie HT. Characterization of vascular adhesion molecules that may facilitate progenitor homing in the post-natal mouse thymus. *Clinical & developmental immunology* 2003;10:27–33. [PubMed: 14575155]
39. Rossi FM, Corbel SY, Merzaban JS, et al. Recruitment of adult thymic progenitors is regulated by P-selectin and its ligand PSGL-1. *Nature immunology* 2005;6:626–634. [PubMed: 15880112]
40. O'Neill HC, De Cello G, O'Neill TJ. Characterization of lymphoid cell lines from murine spleen which can localize in thymus. *International immunology* 1992;4:207–214. [PubMed: 1622895]

41. Kawakami N, Nishizawa F, Sakane N, et al. Roles of integrins and CD44 on the adhesion and migration of fetal liver cells to the fetal thymus. *J Immunol* 1999;163:3211–3216. [PubMed: 10477589]
42. Robertson P, Means TK, Luster AD, Scadden DT. CXCR4 and CCR5 mediate homing of primitive bone marrow-derived hematopoietic cells to the postnatal thymus. *Experimental hematology* 2006;34:308–319. [PubMed: 16543065]
43. Webb SR, Sprent J. Induction of neonatal tolerance to Mls antigens by CD8+ T cells. *Science (New York, N.Y)* 1990;248:1643–1646.
44. Gopinathan R, De Paz HA, Oluwole OO, et al. Role of reentry of in vivo alloMHC peptide-activated T cells into the adult thymus in acquired systemic tolerance. *Transplantation* 2001;72:1533–1541. [PubMed: 11707742]
45. Tian C, Bagley J, Forman D, Iacomini J. Induction of central tolerance by mature T cells. *J Immunol* 2004;173:7217–7222. [PubMed: 15585843]
46. Fontenot JD, Rasmussen JP, Gavin MA, Rudensky AY. A function for interleukin 2 in Foxp3-expressing regulatory T cells. *Nature immunology* 2005;6:1142–1151. [PubMed: 16227984]
47. Kim JM, Rudensky A. The role of the transcription factor Foxp3 in the development of regulatory T cells. *Immunological reviews* 2006;212:86–98. [PubMed: 16903908]
48. Sakaguchi S, Ono M, Setoguchi R, et al. Foxp3+ CD25+ CD4+ natural regulatory T cells in dominant self-tolerance and autoimmune disease. *Immunological reviews* 2006;212:8–27. [PubMed: 16903903]
49. Bayer AL, Yu A, Malek TR. Function of the IL-2R for thymic and peripheral CD4+CD25+ Foxp3+ T regulatory cells. *J Immunol* 2007;178:4062–4071. [PubMed: 17371960]
50. Jiang Q, Su H, Knudsen G, Helms W, Su L. Delayed functional maturation of natural regulatory T cells in the medulla of postnatal thymus: role of TSLP. *BMC immunology* 2006;7:6. [PubMed: 16579866]
51. Fontenot JD, Dooley JL, Farr AG, Rudensky AY. Developmental regulation of Foxp3 expression during ontogeny. *The Journal of experimental medicine* 2005;202:901–906. [PubMed: 16203863]
52. Min B, McHugh R, Sempowski GD, Mackall C, Foucras G, Paul WE. Neonates support lymphopenia-induced proliferation. *Immunity* 2003;18:131–140. [PubMed: 12530982]
53. Cupedo T, Nagasawa M, Weijer K, Blom B, Spits H. Development and activation of regulatory T cells in the human fetus. *European journal of immunology* 2005;35:383–390. [PubMed: 15682453]
54. Darrasse-Jeze G, Marodon G, Salomon BL, Catala M, Klatzmann D. Ontogeny of CD4+CD25+ regulatory/suppressor T cells in human fetuses. *Blood* 2005;105:4715–4721. [PubMed: 15731180]