Published in final edited form as:

J Am Chem Soc. 2009 January 14; 131(1): 14–15. doi:10.1021/ja805541u.

AcOLeDMP and BnOLeDMP: Conformationally Restricted Nucleophilic Catalysts for Enantioselective Rearrangement of Indolyl Acetates and Carbonates

Trisha A. Duffey, Scott A. Shaw, and Edwin Vedejs*

Department of Chemistry, University of Michigan, Ann Arbor, Michigan 48109

Abstract

The rate of indolyl *O*- to *C*-acetyl or carboxyl rearrangement is accelerated by the electron withdrawing *N*-diphenylacetyl group (DPA) using the conformationally restricted chiral catalysts AcOLeDMP (**12b**) and BnOLeDMP (**13b**). Highly enantioselective conversion to quaternary *C*-acetylated and *C*-carboxylated oxindoles is observed, even for substrates containing branched substituents. The rearrangement of the carboxylate substrates **19** occurs with complementary enantiofacial selectivity using catalyst **13b** compared to the acetyl migrations of **16** catalyzed by **12b**. Access to *N*-unsubstituted oxindoles is demonstrated by DPA cleavage with Et₂NH.

The catalytic asymmetric synthesis of all-carbon oxindole quaternary centers has been a difficult challenge. Significant advances have been described using phase-transfer,¹ transition metal,^{2–5} and chiral nucleophilic catalysts.^{6–9} Examples of the latter process include the highly enantioselective rearrangement of indolyl carbonates **1** to the oxindoles **2** catalyzed by planar-chiral pyridines (2 days, 35 °C, 10 mol% catalyst).⁶ Catalyst **3** (TADMAP) also promoted the rearrangement, but a marginal enantioselectivity/reactivity profile at 10% catalyst loading discouraged detailed optimization.⁷ Herein, we report a dramatic substitution effect that solves the reactivity problem. We also describe the development of practical new catalysts featuring a chiral sidechain that prefers a uniquely advantageous conformation. These advances enable enantioselective acyl as well as carboxyl migration in the oxindole series.

Preparation of versatile substrates related to **1** from *N*-protected oxindoles was challenging due to competing reaction at *O* and *C*, but kinetic *O*-acetylation of oxindole **4** with acetyl chloride/2,6-lutidine afforded enol acetate **5** in acceptable 81% yield.¹⁰ Deprotonation of **5** followed by trapping with reactive electrophiles then gave differentially protected indolyl acetates **6** for initial evaluation. Upon catalysis with DMAP (3%), the *N*-benzyl (**6a**) or *N*-alkoxycarbonyl (**6b**) derivatives rearranged slowly to the oxindoles *rac*-**7** (89% and 78% conversion, 5h, rt). On the other hand, the *N*-nosyl (**6c**) or *N*-acyl (**6d–f**; **8**) analogues rearranged completely within 20 min (>95%) while **6g** rearranged to the extent of 84%. Therefore, reactivity increases when more electron-withdrawing substituents are placed at indole nitrogen.

When **3** was used to catalyze the rearrangement of indolyl acetate **6d** to **7d**, much improved reactivity was indeed observed (98% isolated after 3 h, rt; 10% catalyst), but the 20% ee prompted a re-evaluation of the catalyst. One concern was that synthesis of enantiopure **3** requires classical resolution. Another issue is that the trityl group offers few options for catalyst modification short of repeating the entire 5-step synthesis/ resolution sequence. Accordingly, a new family of catalysts was designed that incorporates both electronic and steric factors expected to favor a specific sidechain conformer. Thus, (*S*)-*N*-benzoylvalinol¹¹ was oxidized to aldehyde **10a** and 3-Li- DMAP (from the bromide)⁷ was added to afford the alcohol **11a**. Subsequent acylation yielded **12a** (AcOVaDMAP, 6:1 dr, 72% from **10a**, Scheme 2). A similar route from (*S*)-*tert*-leucinol afforded **12b** (AcOLeDMAP, >98:2 dr; 60% from **10b**). Because diastereomer separation was not necessary with **12b**, this catalyst was used for most of the optimization studies. A third catalyst **13b** (BnOLeDMAP) was prepared by *O*-benzylation of isolated **11**. The stereochemistry and the expected geometry of **12** and **13** (*anti* DMAP and *t*Bu groups; *gauche* OAc and NHBz substituents) were confirmed by X-ray crystallography as shown in **12b**^{*}, and by $J_{1,2} < 1\text{ Hz}$ for the OCHCHNBz protons of **12b** and **13b**. This evidence points to a strong preference for a well-defined catalyst geometry having the benzamido substituent near the catalytic site at the nucleophilic pyridine nitrogen.

Catalyst **12b** effected rearrangement of the *N*-acetyl substrate **6d** to **7d** with promising 61% ee (THF, rt). The *N*-nosyl indole **6c** gave similar results (58% ee), while the *N*-*i*-butyryl indole **6e** rearranged with increased selectivity (77% ee). Reactivity dropped significantly with the *N*-pivalyl analogue **6g**, but the *N*-diphenylacetyl (*N*-DPA) indole **6f** gave good enantioselectivity in THF (89% ee) without impeding the reaction. Optimal 92% ee was obtained in ethyl acetate at 0 °C, although other common solvents also gave good results.

Preparation of **6f** according to Scheme 1 afforded modest yields, so a new route to *N*-DPA indolyl esters was developed. Heating **4** in neat Ph₂CHCOCl (1.5 eq) yielded **14** (85%), and reaction with AcCl/Et₃N afforded the easily purified, crystalline indolyl acetate **15a** as well as ca. 5% of the *C*-acetylated isomer (Scheme 3). Various indolyl acetates prepared in this way were then subjected to catalysis by **12b** (Table 1). Unbranched alkyl groups at the 3-position promoted rearrangement to **16** with good selectivity and reactivity (entries 1–6). A branched alkyl group (*i*-Pr) decreased the rate, but gave oxindole **16f** with 94% ee, while the more reactive 3-phenyl derivative **15g** rearranged with lower selectivity (entry 8).¹² The 5-bromoindolyl acetate **17** reacted completely within 20 min, and the purified major oxindole **18** was found to have the (*S*) configuration by X-ray crystallography (anomalous dispersion). Since the oxindoles **16** and **18** have the same sign of optical rotation and similar chromophores, and also have similar relative mobility for major vs. minor enantiomers on chiral hplc supports, **16a–g** and **18** were assigned the same configuration by analogy.

Using 1 mol% of catalyst **12b**, **15a** rearranged on gram scale within 24 h (entry 2) and recrystallization upgraded the resulting oxindole **16a** to 99% ee. The valine derived catalyst **12a** (AcOVaDMAP) was also effective (**16a**: 88% yield, 90% ee). If desired, the *N*-DPA products **16** can be deprotected to the *N*-H oxindoles **19**. Strong base¹³ or primary amines removed DPA as well as acetyl groups in **16**, but diethyl amine selectively cleaved DPA to give the parent oxindole (**19a**, 75%; **19c** 65%). Retention of configuration was confirmed in the latter case (94% ee). To further illustrate synthetic potential, **16a** (88% ee) was converted into **20a** (73% yield; 87% ee) under Baeyer-Villiger conditions (MCPBA/NaHCO₃, CH₂Cl₂/ reflux).

Catalyst **12b** was also evaluated with the indolyl carbonate substrates. Rearrangement from **8** to **9** occurred readily (5 h, rt), but enantioselectivity was low (4 % ee). The reason became clear when NMR/MS assay of recovered catalyst revealed clean conversion of **12b** to an oxazoline resulting from loss of the *O*-acetyl group.¹⁴ The stable catalyst **13b** (BnOLeDMAP)

catalyzed the conversion from **8** to **9** with modest 33% ee, but good enantioselectivity was achieved by optimizing substituents. Thus, the *N*-DPA indolyl carbonates **21** or **23** (available from 1-naphthylmethyl chloroformate) were converted into oxindoles **22** or **24** with 90–94% ee using **13b** in several representative examples (entries 10–14; CHCl₃, –20 °C). Remarkably, these reactions afforded oxindoles having the *opposite* configuration compared to **16** or **18** obtained using catalyst **12b** according to X-ray crystallography data for **25**, obtained in 94% yield by treatment of **24a** with diethyl amine.¹⁵

In summary, carboxyl and acetyl migration of indolyl esters is strongly accelerated by *N*-acyl groups. Easily accessible AcOLeDMAP (**12b**) and BnOLeDMAP (**13b**) are the most practical and versatile nucleophilic catalysts reported to date for enantioselective rearrangement of indolyl acetates and carbonates to oxindoles containing chiral quaternary carbon. The interplay between migrating group substituents and catalyst modifications at the benzylic oxygen has striking consequences, as illustrated by the complementary enantiofacial selectivity for **13b** with **21/23** vs. **12b** with **15/17**.¹⁶ Furthermore, the catalyst design highlights a conformationally restricted sidechain that may have other uses in situations where convergent functionality in a chirotopic environment is required.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgement

This work was supported by the National Institutes of Health (CA17918). The authors thank Jeff W. Kampf for X-ray crystallographic analysis.

References

1. Poulsen TB, Bernardi L, Alemán J, Overgaard J, Jørgensen KA. J. Am. Chem. Soc 2007;129:441. [PubMed: 17212424]
2. (a) Ashimori A, Bachand B, Overman LE, Poon DJ. J. Am. Chem. Soc 1998;120:6477. (b) Dounay AB, Hatanaka K, Kodanko JJ, Oestreich M, Overman LE, Pfeifer LA, Weiss MM. J. Am. Chem. Soc 2003;125:6261. [PubMed: 12785859]
3. Lee S, Hartwig JF. J. Org. Chem 2001;66:3402. [PubMed: 11348123]
4. (a) Trost BM, Frederiksen MU. Angew. Chem. Int. Ed 2005;44:308. (b) Trost BM, Zhang Y. J. Am. Chem. Soc 2006;128:4590. [PubMed: 16594693] (c) Trost BM, Zhang Y. J. Am. Chem. Soc 2007;129:14548. [PubMed: 17985899] (d) Kündig EP, Seidel TM, Jia Y-X, Bernardinelli G. Angew. Chem. Int. Ed 2007;46:8484.
5. Corkey BK, Toste FD. J. Am. Chem. Soc 2007;129:2764. [PubMed: 17305344]
6. Hills ID, Fu GC. Angew. Chem. Int. Ed 2003;42:3921.
7. (a) Shaw SA, Aleman P, Vedejs E. J. Am. Chem. Soc 2003;125:13368. [PubMed: 14583027] (b) Shaw SA, Aleman P, Christy J, Kampf JW, Va P, Vedejs E. J. Am. Chem. Soc 2006;128:925. [PubMed: 16417383]
8. For reviews of chiral nucleophilic amines see: (a) Spivey AC, Maddaford A, Redgrave AJ. Org. Prep. Proced. Int 2000;32:331. (b) France S, Guerin DJ, Miller SJ, Lectka T. Chem. Rev 2003;103:2985. [PubMed: 12914489] (c) Wurz RP. Chem. Rev 2007;107:5570. [PubMed: 18072804]
9. For other examples of chiral nucleophile catalyzed carboxyl rearrangements see: (a) Nguyen HV, Butler DCD, Richards CJ. Org. Lett 2006;8:769. [PubMed: 16468763] (b) Busto E, Gotor-Fernández V, Gotor V. Adv. Synth. Catal 2006;348:2626.
10. The use of less hindered or more basic amines led to non-selective oxindole acetylation as did less reactive acetyl donors.
11. Menges F, Neuburger M, Pfaltz A. Org. Lett 2002;4:4713. [PubMed: 12489968]

12. Enantiopurity remained the same upon resubjecting oxindole **15b** to the reaction conditions, ruling out a reversible reaction.
13. Kawasaki T, Takamiya W, Okamoto N, Nagaoka M, Hirayama T. Tetrahedron Lett 2006;47:5379.
14. The oxazoline is formed with retention, suggesting that ester cleavage to the alcohol may be the initiating event. Decomposition of catalyst **12b** was not detected in any of the analogous indolyl acetate rearrangements.
15. Oxindole **22a** has the same configuration as **24a** according to correlation by debromination of **24a** with Bu₃SnH. Oxindoles **22** and **24a** have the opposite sense of optical rotation compared to **16** and **18**. The chemical correlation of **22a** with **24a** also supports the analogies used to assign the absolute configurations of **16**.
16. Using **13b** in place of **12b** for Table 2, entry 1, affords the same major enantiomer of **16a** (77% ee; E. McGreevy, unpublished results).

Scheme 1.

Scheme 2.

Scheme 3.

Scheme 4.

Table 1

Table 1A. Rearrangement of indolyl acetates catalyzed by 12b

Entry	indole	R ^d	time	product	ee
1	15a	Me	2 h ^a	16a 94%	92%
2	15a	"	24 h ^b	16a 99%	92%
3	15b	Et	2.5 h ^a	16b 98%	91%
4	15c	Bn	3 h ^a	16c 96%	94% ^c
5	15d	(CH ₂) ₂ OTBDPS	3 h ^a	16d 94%	91%
6	15e	Allyl	2.5 h ^a	16e 98%	86%
7	15f	<i>i</i> -Pr	42 h ^a	16f 82%	94%
8	15g	Ph	2 h ^a	16g 98%	66%
9	17	Me	.33 h ^a	18 95%	85%

Table 1B. Rearrangement of indolyl carbonates catalyzed by 13b

Entry	indole	R ^d	time	product	ee
10	21a	Me	23 h ^d	22a 91%	-90%
11	21b	Bn	23 h ^d	22b 98%	-92%
12	21c	(CH ₂) ₂ OTBDPS	23 h ^d	22c 99%	-94%
13	21d	Ph	2 h ^{d,e}	22d 98%	-91%
14	23a	Me	5 h ^{d,e}	24a 90%	-90%

^a 10 mol % 12b, 0.2M, EtOAc, 0 °C

^b 1 mol % 12b

^c ee of deprotected NH oxindole

^d 10 mol% 13b, 0.8M, CHCl₃, -20 °C

^e 0.2M