PostScript

LETTERS

Dictated versus hand-written accident and emergency discharge documents

New attendances at emergency departments in the UK continue to increase. There were 11.3 million new attendances in 1981, rising to 16 million in 1999.¹ Combined with the government's drive to reduce waiting times, this has resulted in considerable strain being placed on doctors to see, treat and discharge or admit patients quickly, while maintaining accurate records.

One method suggested to reduce doctors' time spent on administration is for discharge documents to be dictated rather than hand-written for all patients sent home directly from the emergency department.

We performed a pilot study assessing whether there was any variance in the quality and content of hand-written versus dictated accident and emergency discharge documentation between two sites in National Health Service Fife. The senior health officers of the accident and emergency department work between the two sites, and facilities for dictation are only available at one of these. This allowed for comparison while eliminating inter-doctor variance. A total of 372 discharge documents were assessed for legibility, the presence or absence of formal diagnoses, investigation results, analgesia given and outcome (discharged, referred or admitted). Dictated documents were superior in every category. Legibility was a particular problem, with 6.6% of hand-written documents being largely illegible. Diagnoses were more often present in dictated records (88% v 69.7%; $p < 0.01$), and, when present, were more often highlighted or obviously located (84% v 44.3%; $p < 0.01$). Radiology results (90% v 81%), analgesia given (100% v 91.7%) and outcome (92% v 86.1%) were also disparate, although these did not achieve statistical significance.

With other studies showing that dictation is faster, improves doctor's satisfaction and departmental productivity,² and that their quality is less adversely affected by increased workload,³ we think this is an area that deserves further attention and hope to conduct a larger-scale study in the near future.

C A Walker

Correspondence to: Dr Craig A Walker, Hairmyres Hospital, East Kilbride, Scotland; dr.walker@fsmail.net

doi: 10.1136/emj.2006.038034

Accepted 30 August 2006

Competing interests: None.

References

- 1 Department of Health, National Assembly for Wales, National Health Service in Scotland. Home of official UK statistics. <http://www.statistics.gov.uk> (accessed 9 Oct 2006).
- 2 Witt DJ. Transcription services in the Emergency Department. *Am J Emerg Med* 1995;13:34–6.

- 3 Dawdy MR, Munter DW, Gilmore RA. Correlation of patient entry rates and physician documentation errors in dictated and handwritten emergency treatment records. *Am J Emerg Med* 1997;15:115–17.

The trauma room: a "one-stop shop" for diagnosis and treatment?

Time taken to complete diagnostic tests and initiate definitive treatment, the "golden hour" of Cowley,¹ is frequently mentioned in the care of seriously injured patients. Surgical care during the first hour is of utmost importance to increase the patient's chances of survival.¹ The availability of the multislice computed tomography (MSCT) enables a complete diagnostic assessment of injured patients in 90–240 s, but is presently carried out only at the end of basic diagnostics. We investigated the effects of a clinical algorithm (fig 1) using MSCT scanning before other measures in the clinical care of seriously injured patients.

In 2004, 139 patients were treated in the trauma room according to the above-mentioned procedure. The mean age of the patients was 42.2 years and the mean Injury Severity Score was 26.93. In all, 12 (8.6%) patients died. The length of stay in the trauma room was markedly reduced from 87 (our data from 2000–1) to 38 min (standard deviation (SD) 19.1 min) on average. The mean duration of complete computed tomography diagnostics (native head computed tomography and contrast medium-supported whole-body computed tomography) was 4 (SD 5.71) min. Diagnostics were completed an average of 13 (SD 8.4) min after arrival of the patient.

Care of seriously injured patients must be carried out as rapidly as possible, and reducing the time needed for clinical care is always

desirable. Rapid computed tomography scanning is sensible and practical. Rapid and complete initial diagnosis by computed tomography markedly reduces the duration in the trauma room, and should considerably improve clinical outcomes. Life-threatening injuries are identified in a matter of minutes and appropriate treatment is initiated. A prerequisite for the use of the concept introduced is the integration of an MSCT into the trauma room, transforming it into a "one-stop shop" for diagnosis and treatment.

P Hilbert, K zur Nieden, R Koch, R Stuttmann
BG-Kliniken Bergmannstrost, Halle/Saale, Germany

Correspondence to: P Hilbert, BG-Kliniken Bergmannstrost, Merseburgerstr 165, Halle/Saale 06112, Germany; dr.peterhilbert@web.de

doi: 10.1136/emj.2006.038554

Accepted 5 September 2006

Competing interests: None declared.

Reference

- 1 Cowley RA. The resuscitation and stabilization of major multiple trauma patients in a trauma center environment. *Clin Med* 1976;83:16–22.

Intravascular haemolysis due to glucose-6-phosphate dehydrogenase deficiency in a patient with aluminium phosphide poisoning

Aluminium phosphide (ALP) poisoning and glucose-6-phosphate dehydrogenase (G6PD) deficiency are two commonly seen clinical presentations in northern India.^{1,2} However,

Figure 1 Effects of a clinical algorithm. CT, computed tomography.