PostScript

LETTERS

Cervical manipulation leading to dissection was not performed by a chiropractor

I read with interest the article by Chen *et al* (2006) "Vertebral artery dissection and cerebellar infarction following chiropractic manipulation".¹ On communication with Dr Chen, it was clarified that this manipulation was performed by an individual who graduated from a college associated with 'Chinese herbs'. This individual did not graduate from an accredited chiropractic college, nor is he licensed to practise chiropractic by any other means. As the manipulation was not done by a qualified chiropractor, it was misleading to identify this as a chiropractic adjustment. Similar to the fact that only qualified surgeons can perform surgeries, only qualified chiropractors can perform chiropractic manipulation.

In addition, the authors report that chiropractors perform adjustments without acknowledging relevant risks. This statement is also false. Chiropractors are duly trained to identify risk factors for spinal manipulation and, in many jurisdictions, are required to obtain informed consent. These measures are taken because it has been well established that there are risks associated with cervical manipulation. These risks have been proved with scientific research and are recognised in the profession.

I understand that the intent of this article was to convey to other emergency ward doctors the diagnostic pitfalls associated with the presentation of cerebellar infarction. However, it is important that this issue be clarified because the use of the term "chiropractic" was misleading to the public and potentially harmful to the profession.

D Liu

Correspondence to: Dr D Liu, Department of Biochemistry/Physiology, Canadian Memorial Chiropractic College, 6100 Leslie Street, Toronto, Ontario, Canada M2H 3J1; dliu@cmcc.ca

Accepted 3 October 2006

Competing interests: None declared.

Reference

- 1 Chen WL, Chern CH, Wu YL, *et al*. Vertebral artery dissection and cerebellar infarction following chiropractic manipulation. *Emerg Med J* 2006;23:84.

What's in a name?

Over the decades, the development of our speciality has been reflected in the changes to the name of our working environment.¹ The debate has been reignited recently by the change in name of the College and by the fact that the Department of Health has changed the name of the speciality on Certificates of Completion of Training.² A survey conducted by the college last year received replies from 52

departments, of which 23 had changed their name to emergency department.

We have noticed that in our Trust a variety of names are used, both for the departments in which we work and for the people working there; and that this provokes confusion for staff and patients alike.

We therefore conducted a survey in the two departments in our Trust, asking everyone who worked in the department (whatever their role) to vote for their preferred name. The options given were "Accident Department", "Accident & Emergency Department", "Casualty" or "Emergency Department".

Only 52% of the 163 people who were sent a ballot paper responded. Of the votes which were received, 40 (47%) people voted for Emergency Department and 38 (45%) for Accident & Emergency Department. Six (7%) people preferred Casualty and 1 (1%) opted for Accident Department.

This leaves us in a quandary. More people voted for Emergency Department than for any other single option, but most still preferred different names. We all agree that our speciality is changing rapidly for the better, and that the College of Emergency Medicine presides over something very different from that overseen by the Casualty Surgeons Association.

However, we cannot agree on the name. Is this important? Will people care or even notice the difference? Some of us feel passionately that with our speciality developing so quickly, we need to mark the difference between the future and the past with a name change. Others believe that whatever we say or do, we will not change people's perception of our work.³ Perhaps it would be of interest to survey our local population especially in this era of increasing public and patient involvement.⁴

For the time being, we have agreed among ourselves to answer the telephone, introduce ourselves and sign our letters each in our own preferred way. It seems as though pronounced change in our Trust is unlikely in the immediate future, unless centrally driven nationwide change occurs.

Nick Jenkins, Rosie Furse, David Potts, Simon Smith

Buckinghamshire Hospitals National Health Service Trust, UK

Correspondence to: Dr N Jenkins, Buckinghamshire Hospitals National Health Service Trust, High Wycombe, HP11 2TT, UK; n.jenkins@doctors.org.uk

doi: 10.1136/emj.2006.041699

Accepted 4 September 2006

Competing interests: None.

References

- 1 Rocke L. Accident and emergency medicine or emergency medicine? *J Accid Emerg Med* 1999;16:74.
- 2 Lockey A. Emergency medicine. *BMJ Career Focus* 2003;327:155.
- 3 Smith S. Is a career in emergency medicine associated with stigma? *Eur J Emerg Med* 2003;10:13-15.
- 4 Gray C. What is clinical governance? *BMJ Career Focus* 2005;330:254.

Substance use among patients attending an accident and emergency department

Alcohol and drug usage is associated with attendance to accident and emergency misuse departments (AEDs).¹ However, despite a clear relationship between substance misuse and trauma, we are not aware of any UK studies that have investigated substance misuse and presentation to the AED. We have recently undertaken a cross-sectional survey of substance use among 350 patients presenting to an inner-London AED, using the Paddington Alcohol Test and the Severity of Dependence Scale (SDS).

Levels of hazardous drinking were at a level similar to that reported in the Alcohol Harm Reduction Strategy.² Levels of substance use were lower than expected; however, the relationship between a positive score on the SDS (for alcohol or other substances) and previous attendance, and the proportion of such patients willing to accept advice, was of interest.

In line with previous studies, we found that 38% of patients were hazardous consumers of alcohol and 19% used other illicit substances, with about one third of such patients reporting signs of dependence. Although hazardous alcohol consumption did not predict previous attendances to the AED, possible dependence on illicit drugs did, with about two thirds of such users willing to accept help or advice. Patients with a positive SDS score were almost three times as likely to have attended the AED on three or more occasions in the previous 12 months (odds ratio 2.84, 95% confidence interval 1.5 to 5.38).

We therefore recommend that a brief screening of all patients for substance use be introduced as part of routine practice. Those patients identified as hazardous consumers could then be directed towards appropriate help and advice, such as an alcohol health worker, leading to a potential reduction in the number of attendances.³

Robert Patton, William Smythe, Helen Kelsall, Francis B Selemon

South London & Maudsley National Health Services Trust, London, UK

Correspondence to: Mr R Patton, National Addiction Centre, 4 Windsor Walk, London SE5 8BB, UK

Informed consent was obtained for publication of the patient's details in this article.

doi: 10.1136/emj.2006.042200

Accepted 2 November 2006

Competing interests: None declared.

References

- 1 Soderstrom CA, Smith GS, Dischinger PC. Psychoactive substance abuse disorders among seriously injured trauma centre patients. *J Am Med Assoc* 1997;277:1769-74.
- 2 Strategy Unit. *Alcohol harm reduction strategy for England*. London: The Cabinet Office, 2004.
- 3 Crawford MJ, Patton R, Touquet R, *et al*. Screening and referral for brief intervention of alcohol misusing patients in an accident and emergency department: a pragmatic randomised controlled trial. *Lancet* 2004;364:1334-9.