Published in final edited form as:

Org Lett. 2008 October 2; 10(19): 4231–4234. doi:10.1021/ol801605w.

Cu(I)-Catalyzed Asymmetric Diamination of Conjugated Dienes

Haifeng Du, Baoguo Zhao, Weicheng Yuan, and Yian Shi*

Department of Chemistry, Colorado State University, Fort Collins, Colorado, 80523

Abstract

A Cu(I)-catalyzed asymmetric diamination for a variety of conjugated dienes and a triene with encouraging ee's has been effectively achieved using (R)-DTBM-SEGPHOS as chiral ligand and di-*tert*-butyldiaziridinone as nitrogen source.

Direct diamination of olefins provides an efficient approach to synthesis of vicinal diamines, which are biologically and chemically important functional moieties.¹ While various metal-mediated and catalyzed diamination has been achieved,^{2–8} catalytic asymmetric diamination of olefins has been less well developed and still remains a challenge in organic synthesis. Earlier, Muñiz and coworkers reported chiral auxiliary based^{9a} and chiral Lewis acid-catalyzed^{9b} asymmetric diamination using bisimidoosmium as reagent. Recently, we reported Pd(0)-catalyzed asymmetric diamination of conjugated dienes^{10a,b} and asymmetric allylic and homoallylic C-H diamination of terminal olefins.^{10c} Previously, we have shown that Cu(I)-catalyzed diamination of conjugated dienes and triene using di-*tert*-butyldiaziridinone (**2**)^{11, 12} as nitrogen source occurs regioselectively at the terminal double bond under mild reaction conditions (Scheme 1)^{8a}, which provides complementary regioselectivity to the Pd(0)-catalyzed diamination.⁷ It is highly desirable develop an asymmetric version of this diamination process to enhance its synthetic utility. However, the Cu(I)-catalyzed diamination likely proceeds via a radical mechanism (Figure 1).^{8,13–15} The involvement of radical intermediates presents a challenge for asymmetric control.

Asymmetric diamination was initially investigated using 10 mol % of CuCl with various commercially available or easily prepared chiral monophosphorus and nitrogen-containing ligands¹⁶ and *trans*-1-phenylbutadiene (**7a**) as test substrate at room temperature for 6 h (Scheme 2). As shown in Figure 2, all the reactions went smoothly to give diamination product **3a** with high conversions, but with 0 to 12% ee. To search for more promising ligands, a series of commercially available chiral bisphosphine ligands¹⁷ were subsequently studied for the diamination of *trans*-1-phenylbutadiene (**7a**) with a 2/1 ratio of CuCl and ligand. It was found that steric bulkiness on the phosphine atoms had a large impact on the enantioselectivities (Figure 3). Encouragingly, the diamination of **7a** with (R)-DTBM-SEGPHOS (**L12**)^{17f} gave 90% conversion and 56% ee (Figure 3).

yian@lamar.colostate.edu.

To improve the enantioselectivity, the reaction conditions, including solvent, temperature, and the ratio of ligand and CuCl were further investigated. It was found that solvent has a significant effect on both reactivity and enantioselectivity (Table 1, entries 1–7), benzene-*d*₆ proved to be the best solvent. The ratio of ligand and CuCl was also very important, only 24% conversion was obtained with slightly higher ee when 1:1 CuCl and **L12** was used (Table 1, entry 8). Lowering temperature improved the enantioselectivity (Table 1, entries 9–13), increasing to 65% ee at 0 °C (small amount of toluene was added to avoid solidification of benzene-*d*₆). Overall, the best reaction conditions involve 10 mol % CuCl and 5.5 mol % **L12** in benzene-*d*₆ with a small amount of toluene at 0 °C (Table 1, entry 10).

Under the optimized conditions, a variety of conjugated dienes can be regio- and enantioselectively diaminated at the terminal double bond in 59–93% yield with 62–74% ee's (Table 2, entries 1–10). The ee could be further improved after recrystallization (Table 2, entry 2). For *cis*-1-phenylbutadiene, isomerization of the *cis* double bond occurred during the reaction and mainly gave *E*-isomer product in 70% ee which was a little higher than direct diamination of *trans*-1-phenylbutadiene (Table 2, entry 1 vs 5). When a mixture of *trans*- and *cis*-dienes was subjected to the reaction conditions, the diamination product of *E* isomer was formed predominately with only a trace amount of *Z* isomer (Table 2, entries 6 and 7). Asymmetric diamination of 1,1-disubstituted butadienes gave slightly higher ee's (Table 2, entries 8–10). When a triene was diaminated at room temperature in benzene-*d*₆, the diamination product was obtained in 58% ee (Table 2, entry 11). Asymmetric diamination of *trans*-1-phenyl-3-methylbutadiene led to 90% yield with only 23% ee (Table 2, entry 12), suggesting that the steric effect and radical stability are important factors for the enantioselectivity.

In summary, catalytic asymmetric diamination for a variety of conjugated dienes and a triene with encouraging ee's has been achieved using CuCl/**L12** as the catalyst and di-*tert*-butyldiaziridinone as the nitrogen source. These results show that Cu(I)-catalyzed asymmetric diamination is feasible despite the fact that the diamination likely involves radical intermediates. Ligand (*R*)-DTBM-SEGPHOS (**L12**) provides a very promising lead for further improvement. Searches for more effective chiral ligands, studies of different nitrogen sources, and expansion of the substrate scope will be carried out.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgement

We are grateful to the generous financial support from the General Medical Sciences of the National Institutes of Health (GM083944-01).

References

1. For leading reviews, see: (a) Lucet D, Gall TL, Mioskowski C. *Angew. Chem., Int. Ed* 1998;37:2580. (b) Mortensen MS, O'Doherty GA. *Chemtracts: Organic Chemistry* 2005;18:555. (c) Kotti SRSS, Timmons C, Li G. *Chem. Biol. Drug. Des* 2006;67:101. [PubMed: 16492158]
2. For examples of metal-mediated diaminations, see: Co: (a) Becker PN, White MA, Bergman RG. *J. Am. Chem. Soc* 1980;102:5676. Hg: (b) Barluenga J, Alonso-Cires L, Asensio G. *Synthesis* 1979;962:962. Mn: (c) Fristad WE, Brandvold TA, Peterson JR, Thompson SR. *J. Org. Chem* 1985;50:3647. Os: (d) Chong AO, Oshima K, Sharpless KB. *J. Am. Chem. Soc* 1977;99:3420. (e) Muñiz K. *Eur. J. Org. Chem* 2004:2243. Pd: (f) Bäckvall J-E. *Tetrahedron Lett* 1978:163. Ti: (g) Aranda VG, Barluenga J, Aznar F. *Synthesis* 1974:504.

3. For recent Cu(II)-mediated intramolecular diamination, see: (a) Zabawa TP, Kasi D, Chemler SR. J. Am. Chem. Soc 2005;127:11250. [PubMed: 16089447] (b) Zabawa TP, Chemler SR. Org. Lett 2007;9:2035. [PubMed: 17447781]
4. For Rh(II)-, Fe(III)-catalyzed diamination with TsNCl₂, see: (a) Li G, Wei H-X, Kim SH, Carducci MD. Angew. Chem., Int. Ed 2001;40:4277. (b) Wei H-X, Kim SH, Li G. J. Org. Chem 2002;67:4777. [PubMed: 12098288]
5. For a recent Pd(II)-catalyzed intermolecular diamination of conjugated dienes, see: Bar GLJ, Lloyd-Jones GC, Booker-Milburn KI. J. Am. Chem. Soc 2005;127:7308. [PubMed: 15898768]
6. For recent Pd(II)- and Ni-catalyzed intramolecular diamination of olefins, see: (a) Streuff J, Hövelmann CH, Nieger M, Muñiz K. J. Am. Chem. Soc 2005;127:14586. [PubMed: 16231907] (b) Muñiz K, Streuff J, Hövelmann CH, Núñez A. Angew. Chem., Int. Ed 2007;46:7125. (c) Muñiz K. J. Am. Chem. Soc 2007;129:14542. [PubMed: 17985900] (d) Muñiz K, Hövelmann CH, Streuff J. J. Am. Chem. Soc 2008;130:763. [PubMed: 18081279]
7. For Pd(0)-catalyzed intermolecular diamination, see: (a) Du H, Zhao B, Shi Y. J. Am. Chem. Soc 2007;129:762. [PubMed: 17243803] (b) Xu L, Du H, Shi Y. J. Org. Chem 2007;72:7038. [PubMed: 17676910] (c) Du H, Yuan W, Zhao B, Shi Y. J. Am. Chem. Soc 2007;129:7496. [PubMed: 17521192]
8. For Cu(I)-catalyzed intermolecular diamination, see: (a) Yuan W, Du H, Zhao B, Shi Y. Org. Lett 2007;9:2589. [PubMed: 17536815] (b) Zhao B, Yuan W, Du H, Shi Y. Org. Lett 2007;9:4943. [PubMed: 17973481] (c) Zhao B, Du H, Shi Y. Org. Lett 2008;10:1087. [PubMed: 18302393]
9. For leading references on asymmetric diamination using bisimidoosmium as reagent, see: (a) Muñiz K, Nieger M. Synlett 2003:211. (b) Muñiz K, Nieger M. Chem. Commun 2005:2729.
10. (a) Du H, Yuan W, Zhao B, Shi Y. J. Am. Chem. Soc 2007;129:11688. [PubMed: 17803307] (b) Xu L, Shi Y. J. Org. Chem 2008;73:749. [PubMed: 18095707] (c) Du H, Zhao B, Shi Y. J. Am. Chem. Soc 2008;130:8590. [PubMed: 18549207]
11. Greene FD, Stowell JC, Bergmark WR. J. Org. Chem 1969;34:2254.
12. For a leading review on diaziridinone, see: HeineHWHassnerA The Chemistry of Heterocyclic Compounds 1983 Hoboken, NJ John Wiley & Sons, Inc 547
13. For a leading review on metal-promoted radical reactions, see: Iqbal J, Bhatia B, Nayyar NK. Chem. Rev 1994;94:519.
14. For leading reviews on CuX-catalyzed atom transfer reactions see: (a) Patten TE, Matyjaszewski K. Acc. Chem. Res 1999;32:895. (b) Clark AJ. Chem. Soc. Rev 2002;31:1. [PubMed: 12108978]
15. For a leading review on nitrogen-centered radicals, see: StellaLRenaudPSibiMPRadicals in Organic Synthesis 2001 Vol. 2 Weinheim, Germany Wiley-VCH 407
16. For leading references on **L1–L6**, see: (a) Hattori T, Shijo M, Kumagai S, Miyano S. Chem. Express 1991;6:335. (b) ref. ^{10a}. (c) Sewald N, Wendisch V. Tetrahedron: Asymmetry 1998;9:1341. (d) Hayashi T, Yamamoto K, Kumada M. Tetrahedron Lett 1974;15:4405. (e) von Matt P, Pfaltz A. Angew. Chem., Int. Ed. Engl 1993;32:566. (f) Evans DA, Woerpel KA, Hinman MM, Faul MM. J. Am. Chem. Soc 1991;113:726. (g) Lowenthal RE, Masamune S. Tetrahedron Lett 1991;32:7373.
17. For leading references on **L7–L12**, see: Kagan HB, Dang T-P. J. Am. Chem. Soc 1972;94:6429. (b) Miyashita A, Yasuda A, Takaya H, Toriumi K, Ito T, Souchi T, Noyori R. J. Am. Chem. Soc 1980;102:7932. (c) Mashima K, Matsumura Y-i, Kusano K-h, Kumobayashi H, Sayo N, Hori Y, Ishizaki T, Akutagawa S, Takaya H. J. Chem. Soc., Chem. Commun 1991:609. (d) Lipshutz BH, Noson K, Chrisman W. J. Am. Chem. Soc 2001;123:12917. [PubMed: 11749557] (e) Hatano M, Terada M, Mikami K. Angew. Chem., Int. Ed 2001;40:249. (f) Lipshutz BH, Lower A, Noson K. Org. Lett 2002;4:4045. [PubMed: 12423082]
18. The reactions were carried out with *trans*-1-phenylbutadiene (0.20 mmol), di-*tert*-butyldiaziridinone (**2**) (0.30 mmol), CuCl (0.02 mmol), ligand **L1–L5** (0.02 mmol), except for **L6** (0.01 mmol) in benzene-*d*₆ (0.60 mL) at room temperature for 6 h. The conversions were determined by crude ¹H NMR, and the ee's were determined by chiral HPLC.
19. The same conditions as ref. ¹⁸ except for **L7–L12** (0.01 mmol).

Figure 1.
A Proposed Catalytic Cycle for Cu(I)-catalyzed Diamination

Figure 2. Asymmetric Diamination of Diene **7a** with Selected Monophosphorus and Nitrogen-containing Ligands¹⁸

Figure 3.
Asymmetric Diamination of Diene **7a** with Selected Bisphosphine Ligands ¹⁹

Scheme 1.

Scheme 2.

Table 1
Studies on Reaction Conditions for Asymmetric Diamination of **7a**^a

entry	CuCl (mol %)	L12 (mol %)	solvent	conv. (%) ^b	ee (%) ^c
1	10	5	DME	63	44
2	10	5	Et ₂ O	43	42
3	10	5	CDCl ₃	78	19
4	10	5	CD ₂ Cl ₂	100	35
5	10	5	THF	21	33
6	10	5	Toluene- <i>d</i> ₈	77	53
7	10	5	C₆D₆	90	56
8	10	10	C ₆ D ₆	24	58
9	10	5	C ₆ D ₆ /PhCH ₃ (11/1, v/v)	73	63
10	10	5.5	C₆D₆/PhCH₃ (11/1, v/v)	72	65
11	12	5	C ₆ D ₆ /PhCH ₃ (11/1, v/v)	74	61
12	15	5	C ₆ D ₆ /PhCH ₃ (11/1, v/v)	89	56
13	10	5.5	PhCH ₃	30	65

^a All the reactions were carried out with *trans*-1-phenylbutadiene (**7a**) (0.20 mmol), di-*tert*-butyldiaziridine (**2**) (0.30 mmol) and solvent (0.60 mL). For entries 1–8, the reactions were carried out at rt for 6 h. For entries 9–13, the reactions were carried out at 0 °C for 20 h.

^b The conversion was determined by crude ¹H NMR.

^c The ee was determined by chiral HPLC (Chiralpak AD-H column).

Table 2
Catalytic Asymmetric Diamination of Dienes and Triene ^a

entry	substrate	product	yield (%) ^b	ee (%) ^f
1	X = H		69	65
2	X = 4-OMe		76	67 (>99) ^g
3	X = 2-OMe		70	67
4			80	62
5			60 ^c	70
				nd ^h
6			65 ^d	66
7			78 ^d	66
8			93	72
9	Ar = <i>p</i> -tolyl		59	74
10	Ar = 2-thienyl		79	66
11			55 ^e	58
12			90	23

^a All the reactions were carried out with olefin (0.20 mmol), di-*tert*-butyldiaziridinone (**2**) (0.30 mmol), CuCl (0.02 mmol), **L12** (0.011 mmol), benzene-*d*₆ (0.55 mL) and toluene (0.050 mL) at 0 °C under argon for 20 h unless otherwise stated.

^b Isolated yield based on diene or triene.

^c The ratio was determined by ¹H NMR.

^d Only trace amount of isomer was observed by ¹H NMR.

^e The reaction was carried out in benzene-*d*₆ at rt for 20 h.

^f The ee was determined by chiral HPLC (Chiralpak AD-H column).

^g The ee after recrystallization from hexanes.

^h Not determined.