Published in final edited form as:

Semin Immunol. 2009 February ; 21(1): 28–35. doi:10.1016/j.smim.2008.08.002.

Role of Retinoic Acid in the Imprinting of Gut-Homing IgA-Secreting Cells

J. Rodrigo Mora^a and Ulrich H. von Andrian^b

^a *Gastrointestinal Unit, Massachusetts General Hospital, Harvard Medical School, 55 Fruit St., GRJ-815, Boston, MA 02114, USA*

^b *Immune Disease Institute & Department of Pathology, Harvard Medical School, 77 Ave. Louis Pasteur, Room 836, Boston, MA 02115, USA*

Abstract

Antibody-secreting cells (ASCs) lodging in the mucosa of the small intestine are derived from activated B cells that are thought to arise in gut-associated lymphoid tissues (GALT). Upon leaving the GALT, B cells return to the blood where they must express the gut-homing receptors $\alpha 4\beta 7$ and CCR9 in order to emigrate into the small bowel. Recent evidence indicates that gut-associated dendritic cells (DCs) in GALT induce gut-homing receptors on B cells via a mechanism that depends on the vitamin A metabolite retinoic acid (RA). In addition, although ASC associated with other mucosal tissues secrete IgA in an RA-independent fashion, the presence of high levels of RA in intestine and GALT can promote B cell class switching to IgA and, thus, boost the production of IgA in the intestinal mucosa. Here we discuss the role of RA in the imprinting of gut-homing ASC and the evidence linking RA with the generation of intestinal IgA-ASCs.

Keywords

B cells; retinoic acid; IgA; antibody-secreting cells; homing

Introduction

IgA is the most abundant immunoglobulin isotype produced in the body (around 3 g/day) and it is estimated that around 80% of all IgA-ASCs reside in the gut mucosa^{1,2}. Mice either lacking IgA or impaired in its secretion are more susceptible to intestinal toxins and pathogens^{3,4}. In addition, migration of B cells and ASCs to the gut is critical for conferring protection against intestinal pathogens⁵⁻⁹. Therefore, both IgA secretion and homing of ASCs to the gut are important in conferring protection at this anatomical site.

Naïve B lymphocytes migrate to secondary lymphoid organs (SLOs), such as lymph nodes, Peyer's patches and the spleen, where they are activated by their cognate antigen¹⁰. Conventional (B2) B cells can be activated by T cell-dependent (TD) antigens, i.e., antigens that elicit concomitant "helper" CD4 T cell responses (usually proteins), and become either ASCs or memory B cells (B_{Mem})¹¹. On the other hand, B cells can also be activated by T cell-

Address correspondence to: J. Rodrigo Mora (j_rodrigo_mora@hms.harvard.edu) or Ulrich H. von Andrian (uva@hms.harvard.edu).

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

independent (TI) antigens, either type-I (polyclonal activators, such as LPS, CpG, poly-IC) or type-II (polysaccharides, such as capsular bacterial polysaccharides) and become mostly short-lived IgM-ASCs¹¹.

Peyer's patches (PPs), and to a lesser extent mesenteric lymph nodes (MLNs), are the main SLOs where B cells differentiate into IgA-ASCs^{2,12}. As we will discuss below, the preferential induction of IgA-ASCs in these sites also explain why they acquire preferential migration to the gut¹³⁻¹⁵. Homing of ASCs to the intestinal mucosa requires the expression of the integrin $\alpha 4\beta 7$ ^{16,17}, which binds to its receptor MAdCAM-1, which is displayed on intestinal postcapillary endothelial cells¹⁸. Moreover, in the case of the small bowel, ASCs also need to express the chemokine receptor CCR9 in order to migrate efficiently to this compartment¹⁹⁻²¹. Another chemokine receptor, CCR10, has been proposed as a "general" mucosal homing receptor²²⁻²⁵. In fact, most IgA-ASCs express CCR10^{24,26} and one of its ligands, the chemokine CCL28/MEC, is expressed by most mucosal epithelia^{23,27}. However, although CCR10 is apparently necessary for the homing of ASCs to the colon²¹ and the lactating mammary gland²², its role in ASC migration to the small bowel remains controversial^{21,28}.

Peritoneal B1 B cells can also give rise to intestinal IgA-ASCs, although the extent of their contribution remains controversial, ranging from 1-50% of all intestinal lamina propria IgA-ASCs, depending on the experimental system and readout²⁹⁻³². In addition, it has been described that conventional B1 B cells are not be found in most mammals (including several mouse strains). Instead, another B cell subset (Bw B cells) has been described in most mouse strains³³. These cells are mostly found in the peritoneal cavity and the spleen and they may play an important role in the production of natural autoantibodies. However, the migratory properties of this particular B cell subset has not been characterized. Moreover, in humans, peritoneal B1 B cells do not seem to be a significant source of intestinal IgA-ASCs³⁴. Thus, the relative contribution of B1 B cells to the pool of intestinal IgA-ASCs and their relevance to gut immunity remain to be determined. It is also unknown what traffic molecules B1 cells need to home to the gut mucosa. Interestingly, a recent study showed that the peritoneal cavity environment can also imprint gut-homing capacity and induce IgA class-switching/secretion on plasmablasts³⁵. In this setting, TLR ligands³⁶ and sphingosine-1-phosphate (S1P)³⁷ may also play important roles allowing the mobilization of peritoneal B1 B cells in order to become intestinal IgA-ASCs. Regarding the latter, it is important to highlight that S1P and S1P receptor type 1 (S1P₁) are also essential for lymphocyte exit from lymphoid compartments, such as lymph nodes and PP^{38,39}. Consistent with this notion, S1P is also important for the egress of IgA plasmablasts from PPs⁴⁰.

Gut-associated dendritic cells and retinoic acid in the imprinting of gut-tropic ASCs

It has been shown that oral vaccination induces higher levels of the gut-homing integrin $\alpha 4\beta 7$ on B cells and ASCs than parenteral administration of the same antigen^{5,6,41-44}. Thus, the site of antigen entry into the body determines the microenvironment where B cells are activated, which in turn strongly influences the homing commitment of the resulting ASCs. In the lymphoid microenvironment, DCs are not only essential for T cell activation^{45,46}, but they can also influence B cell responses by enhancing their differentiation to ASCs and survival^{47,48}. Moreover, DCs can present unprocessed antigens to B cells *in vivo*⁴⁹⁻⁵¹. Since several reports have shown that DCs from PPs and MLNs (GALT-DCs) are sufficient to induce $\alpha 4\beta 7$ and CCR9 and gut-homing capacity on activated T cells⁵²⁻⁵⁷, it was plausible that they could also modulate B cells in a tissue-specific manner. In fact, previous data showed that DCs from PPs, but not from the spleen, promoted IgA class-switching in activated B cells^{58,59}. These findings were recently reproduced and extended to other systems⁶⁰⁻⁶³. Moreover,

similar to their effect on T cells, it was recently shown that PP-DCs and DCs from the lamina propria of the small intestine can also imprint $\alpha 4\beta 7$, CCR9 and gut-homing capacity on ASCs^{61,62}.

Insights into the mechanism by which gut-associated DCs imprint gut-homing T cells was provided in a seminal study by Iwata *et al.*⁶⁴ in which it was shown that the vitamin A metabolite *all-trans* retinoic acid (RA) is sufficient to induce $\alpha 4\beta 7$ and CCR9 on activated T cells, and that blocking RA-receptors of the RAR family decreased the induction of gut-homing receptors by PP-DCs and MLN-DCs. Consistent with a pivotal role of RA in gut-homing imprinting, it was shown more recently that RA is also necessary for the induction of gut-homing receptors on B cells and IgA-ASCs^{61,62} (Fig. 1). These findings provided a molecular explanation for older observations that vitamin A-deficient rats exhibit impaired migration of recently activated MLN lymphocytes to the intestinal mucosa⁶⁵, and that these animals had also a marked decrease in the number of IgA-ASCs and CD4 T cells in their ileum⁶⁶.

Of note, subcapsular sinus macrophages can also present lymph-borne antigens and activate naïve B cells in skin-draining lymph nodes⁶⁷⁻⁶⁹. Since macrophages can secrete BAFF (B-cell activating factor/Blys)⁴⁷ and intestinal lamina propria macrophages secrete RA⁷⁰, it will be interesting to determine whether subcapsular sinus macrophages in the GALT can also imprint tissue-specific homing and/or promote specific IgA class-switching.

The reason why GALT-DCs and lamina propria DCs can secrete RA is explained, at least in part, by their selective expression of retinal dehydrogenases (RALDH), which are critical enzymes for RA synthesis^{62-64,71}. However, other cells in the gut, e.g., intestinal epithelial cells (IEC), also express RALDH and can synthesize RA^{64,72}. Also, extraintestinal sources of RA have been identified in lungs⁷³ and liver⁷⁴. However, the role of RA in those extraintestinal anatomic sites remains to be defined. Of interest, it has been reported that recently activated B cells (plasmablasts) are also imprinted with gut-tropism in the peritoneal cavity³⁵. It will be interesting to assess whether this “peritoneal imprinting” also relies on RA.

Even though CCR10 is expressed on most IgA-ASCs, it is unknown how this receptor is induced on ASCs *in vivo*. 1,25(OH)2VD3, the active form of vitamin D, has been reported to induce CCR10 on *ex vivo* activated human T cells⁷⁵ and ASCs⁷⁶. However, the physiological relevance of 1,25(OH)2VD3 for CCR10 induction is presently unclear. Interestingly, a recent report showed that CCR10 is induced on murine ASCs upon intra-rectal, but not oral, immunization⁷⁷. In the latter study it was proposed that CCR10 upregulation happens in the cecal patches and iliac lymph node, although the molecular mechanism for the induction of this receptor remains to be determined⁷⁷.

Interestingly, B cells may also exhibit homing plasticity⁴⁴. In fact, they can be reeducated and acquire or lose gut-homing potential when they are restimulated with or without RA, respectively⁶¹ (Fig. 1). Similar homing malleability has been documented for T cells^{56,57}. Given that plasma cells are terminally differentiated cells and do not divide, it is likely that homing plasticity operates at the level of B_{Mem} when they are reactivated and proliferate to become ASC. In addition, during a restimulation, and depending on the activation conditions, B cells may also switch to another immunoglobulin isotype. For example, B_{Mem} expressing either IgM, IgG or IgE may, in theory, switch to IgA when reactivated in MALT. In fact, sequential immunoglobulin switching from IgG2b to IgA or, in humans, from IgA1 to IgA2, has been described⁷⁸.

Role of gut-associated DCs and RA in the generation of intestinal IgA-ASCs

The different and complex mechanisms implicated in inducing mucosal IgA-ASCs have been reviewed in detail elsewhere^{10,79-81}. Here we will focus on summarizing and discussing the evidence linking RA with the generation of IgA-ASCs, which also establishes a mechanistic link between the imprinting signals for gut-homing and the modulation of B cell effector function.

It had been known for some time that GALT-DCs can induce IgA-ASCs when cocultured with activated B cells *in vitro*⁵⁸⁻⁶³, even in the absence of T cells⁶¹⁻⁶³. As mentioned above, RA is synthesized/secreted by GALT-DCs and it is essential for the imprinting of gut-homing receptors on T and B cells^{61,64}. Since RA also induces IgA secretion in LPS-activated splenocytes⁸²⁻⁸⁷, we tested the possibility that GALT-DCs may rely on RA for inducing IgA class-switching (Fig. 2). Indeed, it was recently demonstrated that the IgA-promoting effect of GALT-DCs or lamina propria DCs is at least partially dependent on RA⁶¹⁻⁶³. The effect of RA on IgA secretion may be mediated, at least in part, by increased IgA class-switching in RA-exposed B cells⁸⁷. However, the extent to which RA directly influences class-switching⁸⁷ or enhances the proliferation/differentiation of already switched IgA plasmablasts^{88,89} remains to be determined. Nonetheless, consistent with the effect of RA on IgA secretion *in vitro*, oral administration of a RA receptor (RAR)-agonist significantly increases serum IgA levels in rats⁹⁰.

RA may also have effects on other immunoglobulin isotypes. Although vitamin A depletion (hence RA depletion) increases total serum IgG levels^{91,92}, antigen-specific IgG1 responses are decreased⁹³. The latter effect probably reflects the impaired Th2 differentiation observed in the setting of vitamin A deficiency^{94,95} and not a direct effect on B cells. In fact, supplementation of RA inhibits IgG1 production *in vitro* and *in vivo*^{82,90,96,97}. Finally, it has been reported that RA also blocks the production of IgE *in vitro*⁹⁸. However, this IgE-blocking effect was not observed *in vivo*⁹⁹.

The interplay of RA and other DC-derived signals in the induction of IgA-ASCs

It has been reported previously that either IL-5 or IL-6 can influence IgA secretion¹⁰⁰⁻¹¹⁰. In fact, RA-induced IgA secretion requires either exogenous IL-5 or the presence of T cells producing this cytokine^{84,96}. Also, RA induces autocrine production of IL-6 by B cells, which may further contribute to IgA secretion¹¹¹. Moreover, both RA and IL-6 are required for optimal IgA induction by GALT-DCs *in vitro*^{60,61}. Furthermore, RA plus either IL-5, IL-6 or LPS synergize and are sufficient to induce IgA secretion by activated B cells in the presence of non-intestinal DCs^{61,62}. However, IL-5, IL-6 or LPS are probably not directly involved in specific IgA class-switching but are rather permissive for immunoglobulin class-switching by inducing activation-induced cytidine deaminase (AID, an essential enzyme for immunoglobulin class-switching and somatic hypermutation)^{79,112} or by promoting proliferation/differentiation of already switched IgA plasmablasts^{10,113}.

Since it is well established that TGF β is critical for IgA responses *in vivo*¹¹⁴⁻¹¹⁷, it is likely that GALT-DCs rely, at least partially, on TGF β for their IgA-inducing capacity. In fact, GALT-DCs can produce active TGF β ^{70,71,118,119} and blocking TGF β decreases the capacity of PP-DCs to induce IgA-ASCs⁶³. Moreover, mucosal DCs express the integrins $\alpha\beta 6$ and $\alpha\beta 8$, which play an essential role in activating latent TGF β (TGF β non-covalently associated to the latency-associated peptide) *in vivo*¹¹⁹⁻¹²³ (Fig. 2). Nonetheless, it is possible that GALT-DCs also promote IgA in a TGF β -independent manner. In fact, blocking TGF β does not completely abrogate the capacity of PP-DCs to induce IgA-ASCs⁶³. Moreover, although RA also induces TGF β activity in LPS-activated splenocytes and other cells⁸²,

^{124,125}, the IgA-inducing effect of RA is only partially dependent on this cytokine ^{63,82}. These observations are in line with the notion that while the essential *in vivo* role of TGF β in IgA class-switching is well demonstrated, its *in vitro* effects on IgA class-switching/secretion seem to vary significantly depending on the experimental system (this issue is discussed in detail elsewhere ¹⁰).

Interestingly, the inducible form of nitric oxide synthase (iNOS) and nitric oxide (NO) play essential roles for both thymus-dependent and thymus-independent IgA responses ¹²⁶. iNOS/NO seem to be important for the normal expression of TGF- β RII and Smad proteins (involved in TGF β signal transduction) in B cells as well as for the production of APRIL (a proliferation-inducing ligand) and BAFF by DCs ¹²⁶ (Fig. 2). Of note, iNOS is expressed by DCs from small intestinal lamina propria and GALT-DC, but not by spleen DC, and its expression depends on signals driven by commensal flora and TLR signals ¹²⁶, which may contribute to explaining the lower induction of IgA by GALT-DCs isolated from germ-free mice ⁶³. Regarding a potential relationship of iNOS/NO and RA, the iNOS gene promoter has a RA-response element (RARE) that is directly activated by RA bound to its nuclear RAR α /RXR heterodimeric receptor ^{127,128}. In fact, intraperitoneal administration of RAR-agonists, including RA, potentiates LPS-induced iNOS expression in several organs, and also increases plasma levels of nitrate/nitrite in rats ^{128,129}. Thus, RA may also indirectly contribute to IgA secretion by inducing iNOS/NO expression.

APRIL and BAFF (which signal through TACI and BCMA on B cells) can also induce IgA-ASCs, and these factors are important during thymus-independent B cell responses ^{10,79}. It has been reported that the intestinal flora and TLR signals induce BAFF and APRIL in DCs ^{78,130}. Therefore, the decreased IgA induction by GALT-DCs isolated from germ-free mice ⁶³ may be explained, at least in part, by a reduced production of APRIL and BAFF. In addition, since iNOS/NO is necessary to induce BAFF and APRIL secretion by GALT-DC ¹²⁶, RA may also play an indirect role in APRIL/BAFF-mediated IgA responses by upregulating iNOS ^{128,129}. However, it should be mentioned that PP-DCs can still induce IgA responses on TACI- and BCMA-deficient B cells when they are activated with either CD40L or LPS ⁶³. This is analogous to the IgA induction by TGF- β 1 plus LPS, which is also TACI- and BCMA-independent ¹³¹. Thus, the relative role of APRIL and BAFF in IgA induction will ultimately depend on the B cell activation context.

Vasoactive intestinal peptide (VIP) can also induce IgA secretion by human activated B cells ¹³²⁻¹³⁴. Interestingly, it has been reported that RA can induce both VIP and VIP receptors in a neuroblastoma cell line ^{135,136}. However, whether RA plays a role influencing VIP responses on B cells and/or DCs remains to be determined.

To summarize, RA has a direct IgA-promoting effect on activated B cells and it also appears to synergize with several other mechanisms that are thought to promote IgA production in the gut. Consistent with an important *in vivo* role of RA in gut IgA production, rats depleted of vitamin A have decreased levels of total IgA in intestinal lavages and decreased mucosal antigen-specific IgA responses ¹³⁷⁻¹⁴¹. Similarly, vitamin A-depleted mice show impaired IgA secretion and protection at mucosal sites ^{92,142}, as well as impaired IgA responses to bacterial toxins either after oral ¹⁴³ or transcutaneous ¹⁴⁴ immunization. However, it should be kept in mind that vitamin A deficiency may have other effects on the immune system. In fact, the greater susceptibility to intestinal infections and toxins observed in vitamin A-deficient animals may also be explained, at least in part, by a decreased epithelial expression of the polymeric immunoglobulin receptor (pIgR) and therefore a decreased IgA secretion to the intestinal lumen ^{137,141,142,145}. Moreover, although vitamin A-deficient mice have a greatly reduced number of IgA-ASCs in the small bowel ^{61,66}, they have normal serum IgA levels ⁶¹. This indicates that retinoids are not absolutely required for IgA production in tissues other

than the small intestine. Nonetheless, the critical role of RA in T and B cell gut-homing imprinting^{61,64}, as well as its IgA-ASC promoting potential in the gut^{61,82}, may contribute to explain the classical epidemiological observation that vitamin A deficiency is associated with impaired intestinal immune responses^{91,14392,137-142} and markedly increased mortality in children in the developing world¹⁴⁶. It also provides a plausible mechanism to explain the empirical observation that vitamin A supplementation decreases diarrhea and mortality in HIV-infected or malnourished children¹⁴⁷⁻¹⁵¹.

Concluding remarks

It is already well established that gut-associated DCs, including DCs from GALT and lamina propria, can strongly influence T and B cell responses in a tissue-specific manner. Gut-associated DCs, owing to their selective ability to produce and secrete RA, imprint gut-homing capacity on both T and B cells. Moreover, gut-associated DCs can induce B cells to become IgA-ASCs by a mechanism that is, at least in part, dependent on RA. Thus, gut-associated DCs and RA modulate intestinal immune responses by affecting both lymphocyte migration and effector activity.

As discussed above, it is also apparent that RA can potentially interact with other mechanisms inducing IgA-ASCs, such as TGF β , iNOS/NO and probably others. However, the overall relevance of RA for TD and/or TI IgA responses *in vivo* remains to be defined. Also, although RA influences the steady-state lymphocyte composition/numbers in the gut, it has been recently reported that during some viral vaccinations gut-homing imprinting and induction of IgA-ASC may also happen outside the GALT¹⁵². Thus, it will be important to determine how essential RA is for lymphocyte migration during acute immune responses or in various settings of inflammation.

Finally, it will be important to address how gut-associated DCs are “educated” to acquire the capacity to synthesize RA and, thus, to imprint gut-homing lymphocytes and IgA-ASCs. Recent work suggests that commensal flora is necessary to confer GALT-DCs with the capacity to induce IgA-ASCs. Whether this is also true for imprinting gut-homing lymphocytes remains to be determined. If so, it will be interesting to address whether TLR signals and/or commensal bacteria are sufficient to confer non-gut DCs or their precursors with gut-imprinting and/or IgA inducing capacity.

Acknowledgements

We thank Susan Davis for editorial assistance. JRM is indebted to Ingrid Ramos for constant support. JRM is supported by grants from Crohn's & Colitis Foundation of America (CCFA), Cancer Research Institute (CRI), Center for the Study of IBD (CSIBD, DK 43351), Massachusetts Life Sciences Center (MLSC) and the Howard M. Goodman Fellowship (MGH). UHvA is supported by NIH grants AI061663, AI069259, AI072252, HL56949 and AR42689.

References

1. Suzuki K, Ha SA, Tsuji M, Fagarasan S. Intestinal IgA synthesis: a primitive form of adaptive immunity that regulates microbial communities in the gut. *Semin Immunol* 2007;19(2):127–135. [PubMed: 17161619]
2. Macpherson AJ, Slack E. The functional interactions of commensal bacteria with intestinal secretory IgA. *Curr Opin Gastroenterol* 2007;23(6):673–678. [PubMed: 17906446]
3. Lycke N, Erlandsson L, Ekman L, Schon K, Leanderson T. Lack of J chain inhibits the transport of gut IgA and abrogates the development of intestinal antitoxic protection. *J Immunol* 1999;163(2):913–919. [PubMed: 10395687]
4. Wijburg OL, Uren TK, Simpfendorfer K, et al. Innate secretory antibodies protect against natural *Salmonella typhimurium* infection. *J Exp Med* 2006;203(1):21–26. [PubMed: 16390940]

5. Youngman KR, Franco MA, Kuklin NA, et al. Correlation of tissue distribution, developmental phenotype, and intestinal homing receptor expression of antigen-specific B cells during the murine anti-rotavirus immune response. *J Immunol* 2002;168(5):2173–2181. [PubMed: 11859103]
6. Gonzalez AM, Jaimes MC, Cajiao I, et al. Rotavirus-specific B cells induced by recent infection in adults and children predominantly express the intestinal homing receptor alpha4beta7. *Virology* 2003;305(1):93–105. [PubMed: 12504544]
7. Williams MB, Rose JR, Rott LS, et al. The memory B cell subset responsible for the secretory IgA response and protective humoral immunity to rotavirus expresses the intestinal homing receptor, alpha4beta7. *J Immunol* 1998;161(8):4227–4235. [PubMed: 9780197]
8. Kuklin NA, Rott L, Feng N, et al. Protective intestinal anti-rotavirus B cell immunity is dependent on alpha 4 beta 7 integrin expression but does not require IgA antibody production. *J Immunol* 2001;166(3):1894–1902. [PubMed: 11160237]
9. Weitkamp JH, Kallewaard NL, Bowen AL, et al. VH1-46 Is the dominant immunoglobulin heavy chain gene segment in rotavirus-specific memory B cells expressing the intestinal homing receptor {alpha}4{beta}7. *J Immunol* 2005;174(6):3454–3460. [PubMed: 15749880]
10. Mora JR, von Andrian UH. Differentiation and Homing of IgA-Secreting Cells. *Mucosal Immunology* 2008;1(2):96–109. [PubMed: 19079167]
11. McHeyzer-Williams LJ, McHeyzer-Williams MG. Antigen-specific memory B cell development. *Annu Rev Immunol* 2005;23:487–513. [PubMed: 15771579]
12. Butcher EC, Rouse RV, Coffman RL, et al. Surface phenotype of Peyer's patch germinal center cells: Implications for the role of germinal centers in B cell differentiation. *J Immunol* 1982;129:2698–2707. [PubMed: 6982940]
13. Rudzik O, Perey DY, Bienenstock J. Differential IgA repopulation after transfer of autologous and allogeneic rabbit Peyer's patch cells. *J Immunol* 1975;114(1 Pt 1):40–44. [PubMed: 1089713]
14. McDermott MR, Bienenstock J. Evidence for a common mucosal immunologic system. I. Migration of B immunoblasts into intestinal, respiratory, and genital tissues. *J Immunol* 1979;122(5):1892–1898. [PubMed: 448111]
15. Tseng J. Transfer of lymphocytes of Peyer's patches between immunoglobulin allotype congenic mice: repopulation of the IgA plasma cells in the gut lamina propria. *J Immunol* 1981;127(5):2039–2043. [PubMed: 7028869]
16. Farstad IN, Halstensen TS, Lazarovits AI, et al. Human intestinal B-cell blasts and plasma cells express the mucosal homing receptor integrin alpha 4 beta 7. *Scand J Immunol* 1995;42(6):662–672. [PubMed: 8552990]
17. Wagner N, Lohler J, Kunkel EJ, et al. Critical role for beta7 integrins in formation of the gut-associated lymphoid tissue. *Nature* 1996;382:366–370. [PubMed: 8684468]
18. Nakache M, Berg EL, Streeter PR, Butcher EC. The mucosal vascular addressin is a tissue-specific endothelial cell adhesion molecule for circulating lymphocytes. *Nature* 1989;337:179–181. [PubMed: 2911352]
19. Bowman EP, Kuklin NA, Youngman KR, et al. The intestinal chemokine thymus-expressed chemokine (CCL25) attracts IgA antibody-secreting cells. *J Exp Med* 2002;195(2):269–275. [PubMed: 11805153]
20. Pabst O, Ohl L, Wendland M, et al. Chemokine receptor CCR9 contributes to the localization of plasma cells to the small intestine. *J Exp Med* 2004;199(3):411–416. [PubMed: 14744993]
21. Hieshima K, Kawasaki Y, Hanamoto H, et al. CC chemokine ligands 25 and 28 play essential roles in intestinal extravasation of IgA antibody-secreting cells. *J Immunol* 2004;173(6):3668–3675. [PubMed: 15356112]
22. Wilson E, Butcher EC. CCL28 controls immunoglobulin (Ig)A plasma cell accumulation in the lactating mammary gland and IgA antibody transfer to the neonate. *J Exp Med* 2004;200(6):805–809. [PubMed: 15381732]
23. Pan J, Kunkel EJ, Gossler U, et al. A novel chemokine ligand for CCR10 and CCR3 expressed by epithelial cells in mucosal tissues. *J Immunol* 2000;165(6):2943–2949. [PubMed: 10975800]
24. Kunkel EJ, Kim CH, Lazarus NH, et al. CCR10 expression is a common feature of circulating and mucosal epithelial tissue IgA Ab-secreting cells. *J Clin Invest* 2003;111(7):1001–1010. [PubMed: 12671049]

25. Kunkel EJ, Butcher EC. Plasma-cell homing. *Nat Rev Immunol* 2003;3(10):822–829. [PubMed: 14523388]
26. Lazarus NH, Kunkel EJ, Johnston B, et al. A common mucosal chemokine (mucosae-associated epithelial chemokine/CCL28) selectively attracts IgA plasmablasts. *J Immunol* 2003;170(7):3799–3805. [PubMed: 12646646]
27. Wang W, Soto H, Oldham ER, et al. Identification of a novel chemokine (CCL28), which binds CCR10 (GPR2). *J Biol Chem* 2000;275(29):22313–22323. [PubMed: 10781587]
28. Feng N, Jaimes MC, Lazarus NH, et al. Redundant role of chemokines CCL25/TECK and CCL28/MEC in IgA+ plasmablast recruitment to the intestinal lamina propria after rotavirus infection. *J Immunol* 2006;176(10):5749–5759. [PubMed: 16670280]
29. Kroese FG, Butcher EC, Stall AM, Herzenberg LA. A major peritoneal reservoir of precursors for intestinal IgA plasma cells. *Immunol Invest* 1989;18(14):47–58. [PubMed: 2786500]
30. Kroese FG, Butcher EC, Stall AM, et al. Many of the IgA producing plasma cells in murine gut are derived from self-replenishing precursors in the peritoneal cavity. *Int Immunol* 1989;1(1):75–84. [PubMed: 2487677]
31. Macpherson AJ, Gatto D, Sainsbury E, et al. A primitive T cell-independent mechanism of intestinal mucosal IgA responses to commensal bacteria. *Science* 2000;288(5474):2222–2226. [PubMed: 10864873]
32. Thurnheer MC, Zuercher AW, Cebra JJ, Bos NA. B1 cells contribute to serum IgM, but not to intestinal IgA, production in gnotobiotic Ig allotype chimeric mice. *J Immunol* 2003;170(9):4564–4571. [PubMed: 12707334]
33. Thiriot A, Drapier AM, Vieira P, et al. The Bw cells, a novel B cell population conserved in the whole genus *Mus*. *J Immunol* 2007;179(10):6568–6578. [PubMed: 17982046]
34. Boursier L, Farstad IN, Mellembakken JR, Brandtzaeg P, Spencer J. IgVH gene analysis suggests that peritoneal B cells do not contribute to the gut immune system in man. *Eur J Immunol* 2002;32(9):2427–2436. [PubMed: 12207327]
35. Berberich S, Forster R, Pabst O. The peritoneal micromilieu commits B cells to home to body cavities and the small intestine. *Blood* 2007;109(11):4627–4634. [PubMed: 17289810]
36. Ha SA, Tsuji M, Suzuki K, et al. Regulation of B1 cell migration by signals through Toll-like receptors. *J Exp Med* 2006;203(11):2541–2550. [PubMed: 17060475]
37. Kunisawa J, Kurashima Y, Gohda M, et al. Sphingosine 1-phosphate regulates peritoneal B-cell trafficking for subsequent intestinal IgA production. *Blood* 2007;109(9):3749–3756. [PubMed: 17234743]
38. Schwab SR, Cyster JG. Finding a way out: lymphocyte egress from lymphoid organs. *Nat Immunol* 2007;8(12):1295–1301. [PubMed: 18026082]
39. Massberg S, von Andrian UH. Fingolimod and sphingosine-1-phosphate--modifiers of lymphocyte migration. *N Engl J Med* 2006;355(11):1088–1091. [PubMed: 16971715]
40. Gohda M, Kunisawa J, Miura F, et al. Sphingosine 1-phosphate regulates the egress of IgA plasmablasts from Peyer's patches for intestinal IgA responses. *J Immunol* 2008;180(8):5335–5343. [PubMed: 18390715]
41. Kantele A, Westerholm M, Kantele JM, Makela PH, Savilahti E. Homing potentials of circulating antibody-secreting cells after administration of oral or parenteral protein or polysaccharide vaccine in humans. *Vaccine* 1999;17(3):229–236. [PubMed: 9987158]
42. Kantele A, Kantele JM, Savilahti E, et al. Homing potentials of circulating lymphocytes in humans depend on the site of activation: oral, but not parenteral, typhoid vaccination induces circulating antibody-secreting cells that all bear homing receptors directing them to the gut. *J Immunol* 1997;158(2):574–579. [PubMed: 8992970]
43. Quiding-Jarbrink M, Nordstrom I, Granstrom G, et al. Differential expression of tissue-specific adhesion molecules on human circulating antibody-forming cells after systemic, enteric, and nasal immunizations. A molecular basis for the compartmentalization of effector B cell responses. *J Clin Invest* 1997;99(6):1281–1286. [PubMed: 9077537]
44. Kantele A, Arvilommi H, Iikkanen K, et al. Unique characteristics of the intestinal immune system as an inductive site after antigen reencounter. *J Infect Dis* 2005;191(2):312–317. [PubMed: 15609243]

45. Jung S, Unutmaz D, Wong P, et al. In vivo depletion of CD11c(+) dendritic cells abrogates priming of CD8(+) T cells by exogenous cell-associated antigens. *Immunity* 2002;17(2):211–220. [PubMed: 12196292]
46. Probst HC, van den Broek M. Priming of CTLs by Lymphocytic Choriomeningitis Virus Depends on Dendritic Cells. *J Immunol* 2005;174(7):3920–3924. [PubMed: 15778347]
47. Craxton A, Magaletti D, Ryan EJ, Clark EA. Macrophage- and dendritic cell-dependent regulation of human B-cell proliferation requires the TNF family ligand BAFF. *Blood* 2003;101(11):4464–4471. [PubMed: 12531790]
48. Poeck H, Wagner M, Battiany J, et al. Plasmacytoid dendritic cells, antigen, and CpG-C license human B cells for plasma cell differentiation and immunoglobulin production in the absence of T-cell help. *Blood* 2004;103(8):3058–3064. [PubMed: 15070685]
49. Wykes M, Pombo A, Jenkins C, MacPherson GG. Dendritic cells interact directly with naive B lymphocytes to transfer antigen and initiate class switching in a primary T-dependent response. *J Immunol* 1998;161(3):1313–1319. [PubMed: 9686593]
50. Bergtold A, Desai DD, Gavhane A, Clynes R. Cell surface recycling of internalized antigen permits dendritic cell priming of B cells. *Immunity* 2005;23(5):503–514. [PubMed: 16286018]
51. Qi H, Egen JG, Huang AY, Germain RN. Extrafollicular activation of lymph node B cells by antigen-bearing dendritic cells. *Science* 2006;312(5780):1672–1676. [PubMed: 16778060]
52. Stagg AJ, Kamm MA, Knight SC. Intestinal dendritic cells increase T cell expression of alpha4beta7 integrin. *Eur J Immunol* 2002;32(5):1445–1454. [PubMed: 11981833]
53. Mora JR, Bono MR, Manjunath N, et al. Selective imprinting of gut-homing T cells by Peyer's patch dendritic cells. *Nature* 2003;424(6944):88–93. [PubMed: 12840763]
54. Johansson-Lindbom B, Svensson M, Wurbel MA, et al. Selective generation of gut tropic T cells in gut-associated lymphoid tissue (GALT): requirement for GALT dendritic cells and adjuvant. *J Exp Med* 2003;198(6):963–969. [PubMed: 12963696]
55. Dudda JC, Simon JC, Martin S. Dendritic cell immunization route determines CD8+ T cell trafficking to inflamed skin: role for tissue microenvironment and dendritic cells in establishment of T cell-homing subsets. *J Immunol* 2004;172(2):857–863. [PubMed: 14707056]
56. Mora JR, Cheng G, Picarella D, et al. Reciprocal and dynamic control of CD8 T cell homing by dendritic cells from skin- and gut-associated lymphoid tissues. *J Exp Med* 2005;201(2):303–316. [PubMed: 15642741]
57. Dudda JC, Lembo A, Bachtanian E, et al. Dendritic cells govern induction and reprogramming of polarized tissue-selective homing receptor patterns of T cells: important roles for soluble factors and tissue microenvironments. *Eur J Immunol* 2005;35(4):1056–1065. [PubMed: 15739162]
58. Spalding DM, Williamson SI, Koopman WJ, McGhee JR. Preferential induction of polyclonal IgA secretion by murine Peyer's patch dendritic cell-T cell mixtures. *J Exp Med* 1984;160(3):941–946. [PubMed: 6332171]
59. Spalding DM, Griffin JA. Different pathways of differentiation of pre-B cell lines are induced by dendritic cells and T cells from different lymphoid tissues. *Cell* 1986;44(3):507–515. [PubMed: 2417730]
60. Sato A, Hashiguchi M, Toda E, et al. CD11b+ Peyer's patch dendritic cells secrete IL-6 and induce IgA secretion from naive B cells. *J Immunol* 2003;171(7):3684–3690. [PubMed: 14500666]
61. Mora JR, Iwata M, Eksteen B, et al. Generation of gut-homing IgA-secreting B cells by intestinal dendritic cells. *Science* 2006;314(5802):1157–1160. [PubMed: 17110582]
62. Uematsu S, Fujimoto K, Jang MH, et al. Regulation of humoral and cellular gut immunity by lamina propria dendritic cells expressing Toll-like receptor 5. *Nat Immunol* 2008;9(7):769–776. [PubMed: 18516037]
63. Massacand JC, Kaiser P, Ernst B, et al. Intestinal bacteria condition dendritic cells to promote IgA production. *PLoS ONE* 2008;3(7):e2588. [PubMed: 18596964]
64. Iwata M, Hirakiyama A, Eshima Y, et al. Retinoic acid imprints gut-homing specificity on T cells. *Immunity* 2004;21(4):527–538. [PubMed: 15485630]
65. McDermott MR, Mark DA, Befus AD, et al. Impaired intestinal localization of mesenteric lymphoblasts associated with vitamin A deficiency and protein-calorie malnutrition. *Immunology* 1982;45(1):1–5. [PubMed: 6799390]

66. Bjersing JL, Telemo E, Dahlgren U, Hanson LA. Loss of ileal IgA+ plasma cells and of CD4+ lymphocytes in ileal Peyer's patches of vitamin A deficient rats. *Clin Exp Immunol* 2002;130(3): 404–408. [PubMed: 12452829]
67. Phan TG, Grigorova I, Okada T, Cyster JG. Subcapsular encounter and complement-dependent transport of immune complexes by lymph node B cells. *Nat Immunol* 2007;8(9):992–1000. [PubMed: 17660822]
68. Carrasco YR, Batista FD. B cells acquire particulate antigen in a macrophage-rich area at the boundary between the follicle and the subcapsular sinus of the lymph node. *Immunity* 2007;27(1):160–171. [PubMed: 17658276]
69. Junt T, Moseman EA, Iannacone M, et al. Subcapsular sinus macrophages in lymph nodes clear lymph-borne viruses and present them to antiviral B cells. *Nature*. 2007
70. Denning TL, Wang YC, Patel SR, Williams IR, Pulendran B. Lamina propria macrophages and dendritic cells differentially induce regulatory and interleukin 17-producing T cell responses. *Nat Immunol* 2007;8(10):1086–1094. [PubMed: 17873879]
71. Coombes JL, Siddiqui KR, Arancibia-Carcamo CV, et al. A functionally specialized population of mucosal CD103+ DCs induces Foxp3+ regulatory T cells via a TGF- β - and retinoic acid-dependent mechanism. *J Exp Med* 2007;204(8):1757–1764. [PubMed: 17620361]
72. Lampen A, Meyer S, Arnhold T, Nau H. Metabolism of vitamin A and its active metabolite all-trans-retinoic acid in small intestinal enterocytes. *J Pharmacol Exp Ther* 2000;295(3):979–985. [PubMed: 11082432]
73. Dirami G, Massaro GD, Clerch LB, et al. Lung retinol storing cells synthesize and secrete retinoic acid, an inducer of alveolus formation. *Am J Physiol Lung Cell Mol Physiol* 2004;286(2):L249–256. [PubMed: 12832282]
74. Martucci RB, Ziulkoski AL, Fortuna VA, et al. Beta-carotene storage, conversion to retinoic acid, and induction of the lipocyte phenotype in hepatic stellate cells. *J Cell Biochem* 2004;92(2):414–423. [PubMed: 15108365]
75. Sigmundsdottir H, Pan J, Debes GF, et al. DCs metabolize sunlight-induced vitamin D3 to 'program' T cell attraction to the epidermal chemokine CCL27. *Nat Immunol* 2007;8(3):285–293. [PubMed: 17259988]
76. Shirakawa AK, Nagakubo D, Hieshima K, et al. 1,25-Dihydroxyvitamin D3 Induces CCR10 Expression in Terminally Differentiating Human B Cells. *J Immunol* 2008;180(5):2786–2795. [PubMed: 18292499]
77. Chang SY, Cha HR, Uematsu S, et al. Colonic patches direct the cross-talk between systemic compartments and large intestine independently of innate immunity. *J Immunol* 2008;180(3):1609–1618. [PubMed: 18209057]
78. He B, Xu W, Santini PA, et al. Intestinal bacteria trigger T cell-independent immunoglobulin A(2) class switching by inducing epithelial-cell secretion of the cytokine APRIL. *Immunity* 2007;26(6): 812–826. [PubMed: 17570691]
79. Cerutti A. The regulation of IgA class switching. *Nat Rev Immunol* 2008;8(6):421–434. [PubMed: 18483500]
80. Fagarasan S. Evolution, development, mechanism and function of IgA in the gut. *Curr Opin Immunol* 2008;20(2):170–177. [PubMed: 18456485]
81. Cerutti A, Rescigno M. The biology of intestinal immunoglobulin A responses. *Immunity* 2008;28(6):740–750. [PubMed: 18549797]
82. Tokuyama H, Tokuyama Y. Retinoids enhance IgA production by lipopolysaccharide-stimulated murine spleen cells. *Cell Immunol* 1993;150(2):353–363. [PubMed: 8370078]
83. Tokuyama Y, Tokuyama H. Retinoic acid and steroid hormones regulate IgA production by LPS-stimulated murine spleen cells. *Immunopharmacology* 1994;28(2):145–151. [PubMed: 8002288]
84. Tokuyama H, Tokuyama Y. Endogenous cytokine expression profiles in retinoic acid-induced IgA production by LPS-stimulated murine splenocytes. *Cell Immunol* 1995;166(2):247–253. [PubMed: 7497526]
85. Tokuyama Y, Tokuyama H. Retinoids as Ig Isotype-Switch Modulators. *Cell Immunol* 1996;170(2): 230–234.

86. Tokuyama Y, Tokuyama H. Retinoids as Ig isotype-switch modulators. The role of retinoids in directing isotype switching to IgA and IgG1 (IgE) in association with IL-4 and IL-5. *Cell Immunol* 1996;170(2):230–234.
87. Tokuyama H, Tokuyama Y. Retinoic acid induces the expression of germ-line C alpha transcript mainly by a TGF-beta-independent mechanism. *Cell Immunol* 1997;176(1):14–21. [PubMed: 9070313]
88. Morikawa K, Nonaka M. All-trans-retinoic acid accelerates the differentiation of human B lymphocytes maturing into plasma cells. *Int Immunopharmacol* 2005;5(1314):1830–1838. [PubMed: 16275619]
89. Chen Q, Ross AC. Retinoic acid promotes mouse splenic B cell surface IgG expression and maturation stimulated by CD40 and IL-4. *Cell Immunol* 2007;249(1):37–45. [PubMed: 18082674]
90. Kuwabara K, Shudo K, Hori Y. Novel synthetic retinoic acid inhibits rat collagen arthritis and differentially affects serum immunoglobulin subclass levels. *FEBS Lett* 1996;378(2):153–156. [PubMed: 8549823]
91. Wiedermann U, Hanson LA, Bremell T, Kahu H, Dahlgren UI. Increased translocation of *Escherichia coli* and development of arthritis in vitamin A-deficient rats. *Infect Immun* 1995;63(8):3062–3068. [PubMed: 7622231]
92. Stephensen CB, Moldoveanu Z, Gangopadhyay NN. Vitamin A deficiency diminishes the salivary immunoglobulin A response and enhances the serum immunoglobulin G response to influenza A virus infection in BALB/c mice. *J Nutr* 1996;126(1):94–102. [PubMed: 8558330]
93. Carman JA, Smith SM, Hayes CE. Characterization of a helper T lymphocyte defect in vitamin A-deficient mice. *J Immunol* 1989;142(2):388–393. [PubMed: 2463304]
94. Iwata M, Eshima Y, Kagechika H. Retinoic acids exert direct effects on T cells to suppress Th1 development and enhance Th2 development via retinoic acid receptors. *Int Immunol* 2003;15(8):1017–1025. [PubMed: 12882839]
95. Dawson HD, Collins G, Pyle R, et al. Direct and indirect effects of retinoic acid on human Th2 cytokine and chemokine expression by human T lymphocytes. *BMC Immunol* 2006;7:27. [PubMed: 17118196]
96. Tokuyama H, Tokuyama Y. The regulatory effects of all-trans-retinoic acid on isotype switching: retinoic acid induces IgA switch rearrangement in cooperation with IL-5 and inhibits IgG1 switching. *Cell Immunol* 1999;192(1):41–47. [PubMed: 10066345]
97. Tokuyama H, Tokuyama Y, Nakanishi K. Retinoids inhibit IL-4-dependent IgE and IgG1 production by LPS-stimulated murine splenic B cells. *Cell Immunol* 1995;162(1):153–158. [PubMed: 7704904]
98. Worm M, Krah JM, Manz RA, Henz BM. Retinoic acid inhibits CD40 + interleukin-4-mediated IgE production in vitro. *Blood* 1998;92(5):1713–1720. [PubMed: 9716600]
99. Worm M, Herz U, Krah JM, Renz H, Henz BM. Effects of retinoids on in vitro and in vivo IgE production. *Int Arch Allergy Immunol* 2001;124(13):233–236. [PubMed: 11306979]
100. Nikawa T, Ikemoto M, Kano M, et al. Impaired vitamin A-mediated mucosal IgA response in IL-5 receptor-knockout mice. *Biochem Biophys Res Commun* 2001;285(2):546–549. [PubMed: 11444878]
101. Harriman GR, Kunimoto DY, Elliott JF, Paetkau V, Strober W. The role of IL-5 in IgA B cell differentiation. *J Immunol* 1988;140(9):3033–3039. [PubMed: 3258891]
102. Schoenbeck S, McKenzie DT, Kagnoff MF. Interleukin 5 is a differentiation factor for IgA B cells. *Eur J Immunol* 1989;19(6):965–969. [PubMed: 2787753]
103. Matsumoto R, Matsumoto M, Mita S, et al. Interleukin-5 induces maturation but not class switching of surface IgA-positive B cells into IgA-secreting cells. *Immunology* 1989;66(1):32–38. [PubMed: 15493259]
104. Ramsay AJ, Husband AJ, Ramshaw IA, et al. The role of interleukin-6 in mucosal IgA antibody responses in vivo. *Science* 1994;264(5158):561–563. [PubMed: 8160012]
105. McGhee JR, Fujihashi K, Beagley KW, Kiyono H. Role of interleukin-6 in human and mouse mucosal IgA plasma cell responses. *Immunol Res* 1991;10(34):418–422. [PubMed: 1955769]
106. Deviere J, Content J, Denys C, et al. High interleukin-6 serum levels and increased production by leucocytes in alcoholic liver cirrhosis. Correlation with IgA serum levels and lymphokines production. *Clin Exp Immunol* 1989;77(2):221–225. [PubMed: 2505958]

107. Bromander AK, Ekman L, Kopf M, Nedrud JG, Lycke NY. IL-6-deficient mice exhibit normal mucosal IgA responses to local immunizations and *Helicobacter felis* infection. *J Immunol* 1996;156(11):4290–4297. [PubMed: 8666800]
108. McGhee JR, Fujihashi K, Lue C, et al. Role of IL-6 in human antigen-specific and polyclonal IgA responses. *Adv Exp Med Biol* 1991;310:113–121. [PubMed: 1808988]
109. Brandt SJ, Bodine DM, Dunbar CE, Nienhuis AW. Dysregulated interleukin 6 expression produces a syndrome resembling Castleman's disease in mice. *J Clin Invest* 1990;86(2):592–599. [PubMed: 2384605]
110. Beagley KW, Eldridge JH, Lee F, et al. Interleukins and IgA synthesis. Human and murine interleukin 6 induce high rate IgA secretion in IgA-committed B cells. *J Exp Med* 1989;169(6):2133–2148. [PubMed: 2786548]
111. Ballow M, Xiang S, Wang W, Brodsky L. The effects of retinoic acid on immunoglobulin synthesis: role of interleukin 6. *J Clin Immunol* 1996;16(3):171–179. [PubMed: 8734361]
112. Park SR, Kim HA, Chun SK, Park JB, Kim PH. Mechanisms underlying the effects of LPS and activation-induced cytidine deaminase on IgA isotype expression. *Mol Cells* 2005;19(3):445–451. [PubMed: 15995363]
113. Coffman RL, Lebman DA, Shrader B. Transforming growth factor β specifically enhances IgA production by lipopolysaccharide-stimulated murine B lymphocytes. *J Exp Med* 1989;170:1039–1044. [PubMed: 2788703]
114. van Ginkel FW, Wahl SM, Kearney JF, et al. Partial IgA-deficiency with increased Th2-type cytokines in TGF-beta 1 knockout mice. *J Immunol* 1999;163(4):1951–1957. [PubMed: 10438931]
115. Cazac BB, Roes J. TGF-beta receptor controls B cell responsiveness and induction of IgA in vivo. *Immunity* 2000;13(4):443–451. [PubMed: 11070163]
116. Borsutzky S, Cazac BB, Roes J, Guzman CA. TGF-beta receptor signaling is critical for mucosal IgA responses. *J Immunol* 2004;173(5):3305–3309. [PubMed: 15322193]
117. Klein J, Ju W, Heyer J, et al. B cell-specific deficiency for Smad2 in vivo leads to defects in TGF-beta-directed IgA switching and changes in B cell fate. *J Immunol* 2006;176(4):2389–2396. [PubMed: 16455997]
118. Iwasaki A, Kelsall BL. Freshly isolated Peyer's patch, but not spleen, dendritic cells produce interleukin 10 and induce the differentiation of T helper type 2 cells. *J Exp Med* 1999;190(2):229–239. [PubMed: 10432286]
119. Travis MA, Reizis B, Melton AC, et al. Loss of integrin $\alpha(v)\beta(8)$ on dendritic cells causes autoimmunity and colitis in mice. *Nature* 2007;449(7160):361–365. [PubMed: 17694047]
120. Munger JS, Huang X, Kawakatsu H, et al. The integrin $\alpha v \beta 6$ binds and activates latent TGF $\beta 1$: a mechanism for regulating pulmonary inflammation and fibrosis. *Cell* 1999;96(3):319–328. [PubMed: 10025398]
121. Mu D, Cambier S, Fjellbirkeland L, et al. The integrin $\alpha(v)\beta(8)$ mediates epithelial homeostasis through MT1-MMP-dependent activation of TGF-beta1. *J Cell Biol* 2002;157(3):493–507. [PubMed: 11970960]
122. Yang Z, Mu Z, Dabovic B, et al. Absence of integrin-mediated TGFbeta1 activation in vivo recapitulates the phenotype of TGFbeta1-null mice. *J Cell Biol* 2007;176(6):787–793. [PubMed: 17353357]
123. Lacy-Hulbert A, Smith AM, Tissire H, et al. Ulcerative colitis and autoimmunity induced by loss of myeloid $\{\alpha\}v$ integrins. *Proc Natl Acad Sci U S A* 2007;104(40):15823–15828. [PubMed: 17895374]
124. Glick AB, McCune BK, Abdulkarem N, et al. Complex regulation of TGF beta expression by retinoic acid in the vitamin A-deficient rat. *Development* 1991;111(4):1081–1086. [PubMed: 1879351]
125. Glick AB, Flanders KC, Danielpour D, Yuspa SH, Sporn MB. Retinoic acid induces transforming growth factor-beta 2 in cultured keratinocytes and mouse epidermis. *Cell Regul* 1989;1(1):87–97. [PubMed: 2519621]
126. Tezuka H, Abe Y, Iwata M, et al. Regulation of IgA production by naturally occurring TNF/iNOS-producing dendritic cells. *Nature* 2007;448(7156):929–933. [PubMed: 17713535]

127. Zou F, Liu Y, Liu L, et al. Retinoic acid activates human inducible nitric oxide synthase gene through binding of RARalpha/RXRalpha heterodimer to a novel retinoic acid response element in the promoter. *Biochem Biophys Res Commun* 2007;355(2):494–500. [PubMed: 17306764]
128. Devaux Y, Grosjean S, Seguin C, et al. Retinoic acid and host-pathogen interactions: effects on inducible nitric oxide synthase in vivo. *Am J Physiol Endocrinol Metab* 2000;279(5):E1045–1053. [PubMed: 11052959]
129. Seguin-Devaux C, Devaux Y, Latger-Cannard V, et al. Enhancement of the inducible NO synthase activation by retinoic acid is mimicked by RARalpha agonist in vivo. *Am J Physiol Endocrinol Metab* 2002;283(3):E525–535. [PubMed: 12169446]
130. Litinskiy MB, Nardelli B, Hilbert DM, et al. DCs induce CD40-independent immunoglobulin class switching through BLYS and APRIL. *Nat Immunol* 2002;3(9):822–829. [PubMed: 12154359]
131. Castigli E, Wilson SA, Scott S, et al. TACI and BAFF-R mediate isotype switching in B cells. *J Exp Med* 2005;201(1):35–39. [PubMed: 15630136]
132. Kimata H, Fujimoto M. Vasoactive intestinal peptide specifically induces human IgA1 and IgA2 production. *Eur J Immunol* 1994;24(9):2262–2265. [PubMed: 7522170]
133. Kimata H, Fujimoto M. Induction of IgA1 and IgA2 production in immature human fetal B cells and pre-B cells by vasoactive intestinal peptide. *Blood* 1995;85(8):2098–2104. [PubMed: 7536491]
134. Fujieda S, Waschek JA, Zhang K, Saxon A. Vasoactive intestinal peptide induces S(alpha)/S(mu) switch circular DNA in human B cells. *J Clin Invest* 1996;98(7):1527–1532. [PubMed: 8833899]
135. Waschek JA, Muller JM, Duan DS, Sadee W. Retinoic acid enhances VIP receptor expression and responsiveness in human neuroblastoma cell, SH-SY5Y. *FEBS Lett* 1989;250(2):611–614. [PubMed: 2546814]
136. Agoston DV, Colburn S, Krajniak KG, Waschek JA. Distinct regulation of vasoactive intestinal peptide (VIP) expression at mRNA and peptide levels in human neuroblastoma cells. *Neurosci Lett* 1992;139(2):213–216. [PubMed: 1319016]
137. Sirisinha S, Darip MD, Moongkarndi P, Ongsakul M, Lamb AJ. Impaired local immune response in vitamin A-deficient rats. *Clin Exp Immunol* 1980;40(1):127–135. [PubMed: 7389210]
138. Kozakova H, Hanson LA, Stepankova R, et al. Vitamin A deficiency leads to severe functional disturbance of the intestinal epithelium enzymes associated with diarrhoea and increased bacterial translocation in gnotobiotic rats. *Microbes Infect* 2003;5(5):405–411. [PubMed: 12737996]
139. Wiedermann U, Chen XJ, Enerback L, et al. Vitamin A deficiency increases inflammatory responses. *Scand J Immunol* 1996;44(6):578–584. [PubMed: 8972739]
140. Wiedermann U, Hanson LA, Holmgren J, Dahlgren UI. Impaired mucosal immune response in vitamin A deficient rats immunized with oral cholera vaccine. *Adv Exp Med Biol* 1995;371B:1629–1631. [PubMed: 7502871]
141. Wiedermann U, Hanson LA, Kahu H, Dahlgren UI. Aberrant T-cell function in vitro and impaired T-cell dependent antibody response in vivo in vitamin A-deficient rats. *Immunology* 1993;80(4):581–586. [PubMed: 8307607]
142. Gangopadhyay NN, Moldoveanu Z, Stephensen CB. Vitamin A deficiency has different effects on immunoglobulin A production and transport during influenza A infection in BALB/c mice. *J Nutr* 1996;126(12):2960–2967. [PubMed: 9001362]
143. Wiedermann U, Hanson LA, Holmgren J, Kahu H, Dahlgren UI. Impaired mucosal antibody response to cholera toxin in vitamin A-deficient rats immunized with oral cholera vaccine. *Infect Immun* 1993;61(9):3952–3957. [PubMed: 8359917]
144. Chang SY, Cha HR, Igarashi O, et al. Cutting edge: Langerin+ dendritic cells in the mesenteric lymph node set the stage for skin and gut immune system cross-talk. *J Immunol* 2008;180(7):4361–4365. [PubMed: 18354155]
145. Sarkar J, Gangopadhyay NN, Moldoveanu Z, Mestecky J, Stephensen CB. Vitamin A is required for regulation of polymeric immunoglobulin receptor (pIgR) expression by interleukin-4 and interferon-gamma in a human intestinal epithelial cell line. *J Nutr* 1998;128(7):1063–1069. [PubMed: 9649586]
146. Sommer A, Tarwotjo I, Hussaini G, Susanto D. Increased mortality in children with mild vitamin A deficiency. *Lancet* 1983;2(8350):585–588. [PubMed: 6136744]

147. Sommer A, Tarwotjo I, Djunaedi E, et al. Impact of vitamin A supplementation on childhood mortality. A randomised controlled community trial. *Lancet* 1986;1(8491):1169–1173. [PubMed: 2871418]
148. West KP Jr, Pokhrel RP, Katz J, et al. Efficacy of vitamin A in reducing preschool child mortality in Nepal. *Lancet* 1991;338(8759):67–71. [PubMed: 1676467]
149. Lie C, Ying C, Wang EL, Brun T, Geissler C. Impact of large-dose vitamin A supplementation on childhood diarrhoea, respiratory disease and growth. *Eur J Clin Nutr* 1993;47(2):88–96. [PubMed: 8436094]
150. Semba RD, Ndugwa C, Perry RT, et al. Effect of periodic vitamin A supplementation on mortality and morbidity of human immunodeficiency virus-infected children in Uganda: A controlled clinical trial. *Nutrition* 2005;21(1):25–31. [PubMed: 15661475]
151. Mehta S, Fawzi W. Effects of vitamins, including vitamin A, on HIV/AIDS patients. *Vitam Horm* 2007;75:355–383. [PubMed: 17368322]
152. Thompson JM, Nicholson MG, Whitmore AC, et al. Nonmucosal alphavirus vaccination stimulates a mucosal inductive environment in the peripheral draining lymph node. *J Immunol* 2008;181(1): 574–585. [PubMed: 18566424]

Figure 1. Homing imprinting on B cells and ASC

GALT-DC or *all-trans* retinoic acid (RA) induce the expression of $\alpha 4\beta 7$ and CCR9 on ASC and probably also on memory B cells (B_{Mem}), endowing them with the capacity to home to the small bowel. In addition, IgA-ASC migrating to all mucosal tissues express CCR10 and the CCR10 ligand MEC/CCL28 is expressed in all mucosal compartments. However, it is unknown how CCR10 is induced on ASC. Like T cells, B cells also show plasticity regarding their homing commitment. If non gut-homing B cells are restimulated in the presence of RA, they readily upregulate $\alpha 4\beta 7$ and CCR9. On the other hand, B cells with gut-homing capacity lose $\alpha 4\beta 7$ and CCR9 if they are reactivated without RA. Since ASC are terminally differentiated and do not divide, it is likely that the capacity to be “reprogrammed” in their

homing potential resides at the level of B_{Mem} . Finally, whether homing to the bone marrow or sites of inflammation represents a default pathway in the absence of RA or other mucosal signals remains to be determined. (+): agonist/inductive effect. (-): antagonist/blocking effect. Dashed lines: hypothetical/speculative scenario.

Figure 2. Retinoic acid and induction of IgA-ASC

TGF β is directly involved in IgA class-switching and is virtually essential for IgA responses in all mucosal compartments. In the gut, intestinal epithelial cells (IEC) and GALT-DC are among the potential sources of TGF β . DC also express the integrin $\alpha\text{v}\beta 8$, which is essential for TGF β activation *in vivo*. Gut IgA responses to thymus-independent (TI) antigens require APRIL binding to TACI on B cells. APRIL is produced by IEL upon stimulation by commensal flora or TLR signals. These stimuli also induce TSLP secretion by IEL, which in turn induces more APRIL expression by GALT-DC. Mucosal DC (including GALT-DC) express the inducible form of nitric oxide synthase (iNOS), which is also induced by commensal flora and TLR signals. iNOS generates nitric oxide (NO), which is critical for both TD and TI IgA

responses. NO synthesis is necessary for proper TGF- β signaling on B cells and also for APRIL synthesis by GALT-DC. CD40L, APRIL and TLR ligands contribute to IgA class-switching by upregulating the enzyme activation-induced cytidine deaminase (AID), which is essential for both immunoglobulin class-switching and somatic hypermutation. APRIL and TLR ligands may also directly contribute to IgA class-switching. Vasoactive intestinal peptide (VIP) can also induce IgA class-switching *in vitro*, although its significance *in vivo* remains to be determined.

RA, which can be synthesized by intestinal epithelial cells (IEC) and GALT-DC, is probably interrelated to some of the IgA-inducing mechanism mentioned above. RA plus IL-5, IL-6, or TLR signals can promote the differentiation of IgA-ASC in the presence of DC. Whether the latter effect represents mainly a direct effect of RA on IgA class-switching remains to be clarified. RA may also induce IgA class-switching indirectly by upregulating TGF- β secretion and iNOS/NO. RA has also been shown to induce VIP and its receptors in some cell lines.

However, whether this is also relevant for DC and/or B cells remains to be defined. It will also be interesting to determine whether RA can induce the expression of $\alpha\beta 8$ on DC. (+): synthesis, induction or activation. Dashed lines: hypothetical speculative scenario. Questions marks: unknown mechanism.