Published in final edited form as:

Org Lett. 2008 November 20; 10(22): 5257–5260. doi:10.1021/ol8018538.

Intermolecular Silacarbonyl Ylide Cycloadditions: A Direct Pathway to Oxasilacyclopentenenes

Laura E. Bourque and K. A. Woerpel*

Department of Chemistry, University of California, Irvine, California 92697-2025

Abstract

Silacarbonyl ylides, generated by metal-catalyzed silylene transfer to carbonyls, participate in formal intermolecular 1,3-dipolar cycloaddition reactions with carbonyl compounds and alkynes to form dioxasilacyclopentane acetals and oxasilacyclopentenenes in an efficient, one-step process.

The reactions of silylenes with carbonyl compounds have produced a number of mechanistically and synthetically useful reactions. Silylenes (**1**) react with aldehydes and ketones (**2**) to give either oxasilacyclopropanes **3** or silacarbonyl ylides **4** (Scheme 1),¹ and these isomeric products can be interconverted.² The silacarbonyl ylide can be intercepted by π -bonds to form products formally arising from 1,3-dipolar cycloadditions.³ This reaction, which forms new carbon–carbon and carbon–silicon bonds, has not been found to be preparatively useful because yields are often low, even in intramolecular examples.^{1a} In addition, the generation of silylenes requires forcing conditions such as extended photolysis with excess reagents.⁴

In this Letter, we report that metal-catalyzed silylene transfer to carbonyl compounds in the presence of an electron-deficient alkyne produced formal 1,3-dipolar cycloaddition products efficiently under mild conditions. The products of these reactions possess functional groups that can be functionalized with control of regioselectivity.

We discovered that silylene transfer to carbonyl compounds led to three-component cyclization reactions in our attempts to form oxasilacyclopropanes. Treatment of cyclohexene silacyclopentane **6**⁵ with two equivalents of benzaldehyde **5** and a catalytic amount of silver triflate did not yield the expected three-membered ring. Instead, it gave the product of silylene transfer to the C–O double bond and subsequent insertion of the second equivalent of aldehyde to form dioxasilacyclopentane **7** (Scheme 2).⁶ A screen of metal salts known to catalyze silylene transfer to alkenes⁷ did not lead to formation of oxasilacyclopropanes. When *n*-butyraldehyde (**8**) was subjected to the same catalyst screen, copper (II) bromide-catalyzed silylene transfer afforded the intermolecular 1,3-dipolar cycloaddition product, dioxasilacyclopentane acetal **9**, in 89% yield (Scheme 2). A competition experiment between aldehydes **5** and **8** yielded dioxasilacyclopentane **7** as the sole product, suggesting that the difference in regioselectivity of the corresponding products **7** and **9** is caused by a divergence in the mechanistic pathways.

Silylene transfer to benzaldehyde in the presence of electron-deficient alkyne **10** led to a different three-component coupling reaction. When one equivalent of alkyne **10** and one equivalent of benzaldehyde (**5**) were treated under metal-catalyzed silylene transfer conditions, oxasilacyclopentene **11** was formed in 85% yield (Scheme 3). This reaction is complementary to our earlier synthesis of oxasilacyclopentenones.⁸ That method involved a two-step process whereupon silacycloprenes, prepared from alkynes, underwent insertion reactions with carbonyl compounds. Electron-deficient alkynes, however, did not participate in that reaction because the silylenoid intermediate is electrophilic, not nucleophilic.⁹

A variety of carbonyl compounds participated in the cycloaddition employing alkyne **10** as the dipolarophile (Table 1). The cycloaddition reaction tolerated α,β -unsaturated aldehydes and formate esters (entries 1 and 2). Formamides also reacted to form *N,O*-acetal oxasilacyclopentene **13c**, although the reaction was slower. Ketones underwent cyclization to form oxasilacyclopentenones containing tetrasubstituted carbon atoms (entry 4). α,β -Unsaturated ester **12e** reacted with alkyne **10** to form oxasilacyclopentene **13e**, although silylene transfer to the α,β -unsaturated ester was also observed (Scheme 4).¹⁰ Excess ester and silacyclop propane were employed to optimize the formation of oxasilacyclopentene **13e** (entry 5).

The highly electron-deficient alkyne **10** was the only alkyne that participated in these reactions. Treatment of silacyclop propane **6** and carbonyl compounds with other dipolarophiles led to either silylene transfer to the alkyne to form silacycloprenes or dimerization with the aldehyde to form dioxasilacyclopentane **7** (Scheme 5).

Control experiments indicated that the formation of the oxasilacyclopentene did not involve a silacyclop propane intermediate. When electron-deficient alkyne **10** was treated with silacyclop propane **6**, formation of silacyclop propane **19** was not detected (Scheme 6). The ²⁹Si NMR spectrum of the reaction mixture revealed two peaks at δ 18.0 and 4.8 ppm, with no peaks around δ -70 ppm, which would be expected for a silacyclop propane.¹¹ The silylene transfer product could not be identified, but the chemical shifts were consistent with the presence of alkyl silanes.

It has been previously reported that electrophilic carbenes react with aldehydes to form carbonyl ylides that then undergo 1,3-dipolar cycloadditions preferentially with electron-deficient alkynes.¹² This precedent, suggests that the formation of the oxasilacyclopentenones occurs through an analogous concerted 1,3-dipolar cycloaddition pathway (Scheme 7).

The oxasilacyclopentenones that possessed acetal moieties could be elaborated by nucleophilic substitution reactions. Oxasilacyclopentene **13b** was treated with allyltrimethylsilane and SnBr₄ to afford allylated oxasilacyclopentene **21** (Scheme 8), although the modest stabilization of the oxocarbenium ion intermediate by the electron-deficient ester required long reaction times (40 hours).¹³ Oxasilacyclopentene **13e** could be allylated under the same reaction conditions as **13b**, but an unanticipated product was observed. Upon Lewis acid activation to form the oxocarbenium ion, allyltrimethylsilane added to the unsubstituted terminus of the conjugated system, resulting in oxasilacyclopentene **22**.¹⁴

The carbonyl groups can be functionalized and appear to be sterically differentiated. Oxasilacyclopentene **11** was reduced by *i*-Bu₂AlH to form diol **23**. When oxasilacyclopentene **11** was treated with two equivalents of methylmagnesium bromide, bicyclic α,β -unsaturated lactone **24** was formed (Scheme 9). Addition of one equivalent of a Grignard reagent resulted in a 1:1 mixture of starting material and lactone **24**. These experiments reveal that the two carbonyls can be functionalized independently or simultaneously, and suggest that the dissimilarity is due to the steric hindrance imparted by the bulky *tert*-butyl groups attached to the silicon atom.

Conversion of oxasilacyclopentene **13d** to allylic alcohol **25** demonstrates the synthetic utility of the cycloaddition reaction. Initial attempts at removing the silicon atom through conventional protodesilylation methods were unsuccessful.¹⁵ The silicon atom could be removed to form allylic alcohol **25** utilizing palladium salts and Bu₄NF as the fluoride source.¹⁶ Oxasilacyclopentene **11** also underwent protodesilylation, but allylic alcohol **26** isomerized to provide ketone **27**.

In conclusion, silylene transfer to carbonyl compounds can be used to produce dioxasilacyclopentane acetals and oxasilacyclopentenones in a single step through formal intermolecular 1,3-dipolar cycloadditions. The oxasilacyclopentene products are useful intermediates that can be elaborated by substitution, addition, and protodesilylation reactions.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgements

This research was supported by the National Institute of General Medical Sciences of the National Institutes of Health (GM-54909). L.E.B. thanks the Department of Education (GAANN) for a predoctoral fellowship. K.A.W. thanks Amgen and Lilly for awards to support research. We thank Dr. Phil Dennison (UCI) for assistance with NMR spectroscopy and Dr. John Greaves and Ms. Shirin Sorooshian (UCI) for mass spectrometry.

References

1. (a) Ando W, Ikeno M, Sekiguchi A. *J Am Chem Soc* 1978;100:3613–3615. (b) Ando W, Hagiwara K, Sekiguchi A. *Organometallics* 1987;6:2270–2271. (c) Sakai N, Fukushima I, Minakata S, Ryu I, Komatsu M. *Chem Commun* 1999:1857–1858.
2. (a) Gordon MS, George C. *J Am Chem Soc* 1984;106:609–611. (b) Becerra R, Cannady JP, Walsh R. *J Phys Chem* 1999;103:4457–4464. (c) Becerra R, Cannady JP, Walsh R. *Phys Chem Chem Phys* 2001;3:2343–2351.
3. Padwa A, Weingarten MD. *Chem Rev* 1996;96:223–270. [PubMed: 11848752]
4. (a) Sakai N, Fukushima T, Okada A, Ohashi S, Minakata S, Komatsu M. *J Organomet Chem* 2003;686:368–372. (b) Minakata S, Ohashi S, Amano Y, Oderaotoshi Y, Komatsu M. *Synthesis* 2007:2481–2484.
5. Boudjouk P, Black E, Kumarathan R. *Organometallics* 1991;10:2095–2096.
6. (a) Jutzi P, Eikenberg D, Bunte EA, Möhrke A, Neumann B, Stammler HG. *Organometallics* 1996;15:1930–1934. (b) Belzner J, Ihmels H, Pauletto L, Noltemeyer M. *J Org Chem* 1996;61:3315–3319. (c) Franz AK, Woerpel KA. *J Am Chem Soc* 1999;121:949–957.
7. Ćiraković J, Driver TG, Woerpel KA. *J Org Chem* 2004;69:4007–4012. [PubMed: 15176824]
8. (a) Seyferth D, Vick SC, Shannon ML. *Organometallics* 1984;3:1897–1905. (b) Clark TB, Woerpel KA. *J Am Chem Soc* 2004;126:9522–9523. [PubMed: 15291539]
9. Driver TG, Woerpel KA. *J Am Chem Soc* 2004;126:9993–10002. [PubMed: 15303873]
10. Calad SA, Woerpel KA. *J Am Chem Soc* 2005;127:2046–2047. [PubMed: 15713073]
11. Palmer WS, Woerpel KA. *Organometallics* 1997;16:4824–4827.
12. (a) Hodgson DM, Glen R, Grant GH, Redgrave AJ. *J Org Chem* 2003;68:581–586. [PubMed: 12530886] (b) Russell AE, Brekan J, Gronenberg L, Doyle MP. *J Org Chem* 2004;69:5269–5274. [PubMed: 15287769]
13. Creary X, Geiger CC. *J Am Chem Soc* 1982;104:4151–4162.
14. One alkene isomer was observed. We have assigned the product as the Z-alkene isomer because this configuration should minimize steric interactions. However, spectroscopic experiments showed no NOE between the vinyl hydrogen and the carboethoxy groups.
15. Heitzman CL, Lambert WT, Mertz E, Shotwell JB, Tinsley JM, Va P, Roush WR. *Org Lett* 2005;7:2405–2408. [PubMed: 15932209] and references therein

16. Copper salts have been employed in protodesilylation reactions: Trost BM, Ball ZT, Jöge T. J Am Chem Soc 2002;124:7922–7923. [PubMed: 12095335]

Scheme 1.
Silylene Transfer to Carbonyl Compounds

Scheme 2.
Silylene Transfer to Aldehydes

Scheme 3.
Formation of Oxasilacyclopentene

Scheme 4.
Silylene Transfer to α - β -Unsaturated Ester

Scheme 5.

Dipolarophile Screen

^a As determined by ¹H NMR spectroscopic analysis of the product relative to an internal standard (PhSiMe₃).

Scheme 6.
Silacyclopropene Control Experiment

Scheme 7.
Proposed Cycloaddition

Scheme 8.
Nucleophilic Substitution to Oxasilacyclopentenes

Scheme 9.
Functionalization of Carbonyls

Scheme 10.
Removal of Silicon

Table 1
Silver-Catalyzed Silylcarbonyl Ylide Cycloaddition

entry	substrate (12)	product (13)
1		
2		
3		
4

entry	substrate (12)	product
5		

^a At 50 °C for 12 h.

^b For 3 h.

^c Four equivalents of **6** and four equivalents of **12e**.