Omeprazole and ocular damage

Facts of cases are unclear

EDITOR—In the report by Schönhöfer et al of visual loss related to the proton pump inhibitor omeprazole, several issues need clarification.¹ Six of the nine cases are stated to have had “funduscopically confirmed” irreversible anterior ischaemic optic neuropathy. This clinical entity is not diagnosed solely on the basis of fundal findings but in the light of the nature and pattern of visual loss, the appearances of the optic disc in both eyes—an important clue being a relatively small optic disc in the unaffected eye²—and the results of fluorescein angiography. This is an important consideration because many forms of anterior optic neuropathy, including demyelinative or postviral optic neuritis and neoplastic optic nerve infiltration, can produce the same appearance of the optic nerve head as anterior ischaemic optic neuropathy. Indeed, the authors seem to be unclear whether demyelinative optic neuritis or anterior ischaemic optic neuropathy is associated with multiple sclerosis.

Advice to authors

We receive more letters than we can publish: we can currently accept only about one third. We prefer short letters that relate to articles published within the past four weeks. We also publish some “out of the blue” letters, which usually relate to matters of public policy.

When deciding which letters to publish we favour originality, assertions supported by data or by citation, and a clear prose style. Letters should have fewer than 400 words (please give a word count) and no more than five references (including one to the BMJ article to which they relate); references should be in the Vancouver style. We welcome pictures.

Letters, whether typed or sent by email, should give each author's current appointment and full address. Letters sent by email should give a telephone and fax number when possible. We encourage you to declare any conflict of interest. Please send a stamped addressed envelope if you would like to know whether your letter has been accepted or rejected.

We may post some letters submitted to us on the world wide web before we decide on publication in the paper version. We will assume that correspondents consent to this unless they specifically say no.

Letters will be edited and may be shortened.

It is also stated that two patients had no known risk factors for anterior ischaemic optic neuropathy. In fact, both had gastrointestinal ulcers, which have been shown in a prospective review of over 400 patients to be significantly associated with non-arteritic anterior ischaemic optic neuropathy.³ This in itself generates difficulty when attempting to associate a treatment for gastrointestinal ulcers with an increased risk of anterior ischaemic optic neuropathy.

In their concluding sentence, the authors say that anterior ischaemic optic neuropathy is due to retinal artery ischaemia, whereas it is due to a perfusion deficit in the posterior ciliary circulation. There may well be an association between omeprazole and optic nerve dysfunction, but this paper fails to provide conclusive evidence of such an association or a clear understanding of its possible aetiology.

Paul Riordan-Eva *Consultant ophthalmic surgeon*
West Kent Eye Centre, Farnborough Hospital,
Orpington, Kent BR6 8ND

Michael D Sanders *Consultant ophthalmologist*
Department of Neuro-Ophthalmology, National
Hospital for Neurology and Neurosurgery, Queen
Square, London WC1N 3BG

1 Schönhöfer P, Werner B, Tröger U. Ocular damage associated with proton pump inhibitors. *BMJ* 1997;314:1805. (21 June.)

2 Burde RM. Optic disk risk factors for nonarteritic anterior ischemic optic neuropathy. *Am J Ophthalmol* 1993;116: 759-64.

3 Hayreh SS, Joos KM, Podhajsky PA, Long CR. Systemic diseases associated with nonarteritic anterior ischemic optic neuropathy. *Am J Ophthalmol* 1994; 188:766-80.

Concerns on safety of drug are unwarranted

EDITOR—Schönhöfer et al's article on ischaemic optic neuropathy from omeprazole¹ may have created unwarranted concerns among doctors and patients about the safety of this drug.

Several years ago, after allegations that visual impairment was a side effect of omeprazole, Astra invited me as a neuro-ophthalmologist to review case reports and decide if the evidence supported the allegations. Since then I have intermittently reviewed additional cases. The actual diagnoses in these cases were a hodgepodge of disorders (from uveitis to pre-syncope) but some represent instances of anterior ischemic optic neuropathy consequent to gastrointestinal bleeding, a complication of haemorrhage that was recognised many decades before omeprazole was introduced. I have yet to identify a single patient in

whom drug toxicity caused visual dysfunction. The results of these case analyses were transmitted in detail at a meeting of the Committee for Proprietary Medicinal Products in Brussels in July 1994. The negative result of the “pharmacoepidemiological” study cited in the article also provides reassurance about the safety of omeprazole.

Of the nine cases in the article, six had been discussed at the Brussels meeting. Case 3 was that of a patient terminally ill with metastatic cancer whose vague symptoms were never evaluated in life. Cases 1, 2, 4, and 5 were examples of a bilateral optic neuropathy in which a cause other than drug intoxication was likely. All of these patients received several drugs in addition to omeprazole. In case 6, the correct diagnosis was an inflammatory or demyelinative optic neuropathy.

I suspect that case 7 is an example of anterior ischemic optic neuropathy and apparently omeprazole was the only drug that this patient received; however, as with the other cases it seems inappropriate to implicate omeprazole as the putative cause of the neuropathy. Idiopathic anterior ischemic optic neuropathy is by no means rare even in the absence of risk factors for vascular disease. It also is unlikely that a toxic optic neuropathy would affect only one eye.

The hypothesis proposed by the authors for ocular damage induced by omeprazole seems ill founded. Omeprazole may well affect ATPases in non-gastric tissues, but there is no evidence that ATPases are inhibited in the optic nerve or retina or that such inhibition would have any role in the pathogenesis of anterior ischaemic optic neuropathy. The inhibitory effect of omeprazole on production of cerebrospinal fluid is irrelevant to optic nerve function. Drugs such as carbonic anhydrase inhibitors which reduce cerebrospinal fluid production do not impair optic nerve function and in fact are widely used in the long term to prevent optic nerve damage in glaucoma.

Simmons Lessell *Professor of ophthalmology*
Department of Ophthalmology, Harvard Medical
School, 243 Charles Street, Boston, MA 02114,
USA

1 Schönhöfer P, Werner B, Tröger U. Ocular damage associated with proton pump inhibitors. *BMJ* 1997;314:1805. (21 June.)

Lack of causality holds true

EDITOR—Schönhöfer et al discuss two new cases (cases 6 and 7) and attempt to link anterior ischaemic optic neuropathy with oral dosing of omeprazole.¹ Two other patients (cases 8 and 9) had, they suggest,