Published in final edited form as:

Depress Anxiety. 2009 ; 26(4): 309–315. doi:10.1002/da.20528.

RGS2 and Generalized Anxiety Disorder in an Epidemiologic Sample of Hurricane-Exposed Adults

Karestan C. Koenen, Ananda B. Amstadter, Kenneth J. Ruggiero, Ron Acierno, Sandro Galea, Dean G. Kilpatrick, and Joel Gelernter

From the Departments of Society, Human Development and Health and Epidemiology, Harvard School of Public Health, Boston, MA(KCK); Department of Psychiatry, Medical University of South Carolina, Charleston, SC (AA, KJR, RA, DGK); Auburn University, Auburn, AL (AA); Department of Epidemiology, University of Michigan School of Public Health, Ann Arbor, MI (SG), Departments of Psychiatry, Genetics, and Neurobiology, Yale University School of Medicine, New Haven, CT (JG).

Abstract

Background—Generalized anxiety disorder (GAD) is a common and sometimes disabling condition often associated with stressful life events that involve significant loss or danger. The disorder appears moderately heritable. Polymorphisms in the *RGS2* (regulator of G-protein signaling 2) gene were recently associated with anxious behavior in mice and panic disorder and trait anxiety in humans. We examined whether rs4606, a single nucleotide polymorphism (SNP) in the 3' UTR of *RGS2*, was associated with GAD in an epidemiologic sample of adults exposed to the 2004 Florida Hurricanes.

Methods—The sample for the current study is 607 adults from the 2004 Florida Hurricane Study who returned buccal DNA samples via mail. Participants were selected via random digit dial procedures and interviewed via telephone about hurricane exposure, social support, and GAD symptoms. The outcome measure was DSM-IV diagnosis of GAD derived from structured interviews.

Results—*RGS2* SNP rs4606 was significantly associated with GAD in this sample. In logistic regression analyses, each C allele was associated with a 100% ($p=.026$) increased risk of GAD after controlling for age, sex, ancestry, hurricane exposure and social support.

Conclusions—These findings are consistent with a previously published study showing a higher prevalence of the C allele among panic disorder patients than controls. This work points toward a relevant polymorphism for GAD at the 3' end of the *RGS2* gene; and suggest that studying a recently disaster-exposed sample is both feasible and may improve power to find gene-disorder associations.

Keywords

anxiety; *RGS2*; SNP; association analysis; trauma

Address reprint requests to: Karestan C. Koenen, PhD; Department of Society, Human Development, and Health; Harvard School of Public Health, 677 Huntington Avenue, Kresge 613, Boston, MA 02115. Phone: 617-432-4622. Fax: 617-432-3755. Email: kkoenen@hsph.harvard.edu.

The authors declare no conflicting financial or other competing interests.

Introduction

Generalized anxiety disorder (GAD) is characterized by exaggerated and uncontrollable worry or tension about everyday events at a level more severe than warranted by the situation. Symptoms of GAD also include restlessness, being easily fatigued, having difficulty concentrating, irritability, muscle tension, and sleep disturbance (APA, 1994). GAD is common and sometimes debilitating, with a lifetime prevalence of 5.7% and a 12-month prevalence of 3.1% in the United States (Kessler et al. 2005a; Kessler et al. 2005b; Kessler et al. 2001). Family and twin studies support the role of genetic factors in the etiology of GAD; a meta-analysis of twin studies reported the heritability of GAD to be 31% (95% CI: 24–39%; (Hettema et al. 2001)). However, little progress has been made in identifying specific genes that increase risk of GAD.

GAD is common after natural disasters (Kar and Bastia 2006; Shore et al. 1986) and stressful life events, particularly those involving loss and danger, have been particularly associated with increased risk of GAD (Kendler et al. 2003). Hurricanes, having the potential to produce widespread property damage, physical injury, and death, are stressful life events involving both loss and danger, a fact that was dramatically demonstrated by the widespread devastation produced by Hurricanes Katrina and Rita in the Gulf Coast regions of the United States in 2005. Prior genetic association studies of GAD have not included stress-exposed samples. Moffitt, Caspi, and Rutter (2005) recently suggested the exposed-cohort design as an efficient way of testing genotype-environment interaction: “If a good candidate gene is available, such an exposed sample could be used to test the hypothesis that genotype-risk individuals develop psychopathology but genotype-controls do not” (Moffitt et al. 2005).

We hypothesized that rs4606, a polymorphism in the 3' untranslated region of the *RGS2* (regulator of G-protein signaling 2) gene, would be associated with GAD in an epidemiologic sample of older adults who were exposed to one or more hurricanes during the 2004 Florida Hurricane season (Acierno et al. 2007; Acierno et al. 2006; Galea et al. 2006; Kilpatrick et al. 2007). Regulators of G-protein signaling bind to G α subunits and increase their GTPase activity which then attenuates the downstream signaling (Hollinger and Hepler 2002). *RGS2*, a potent regulator that reduces G-protein activity, selectively inhibits Gq α (Heximer 2004), therefore increasing GTPase activity. G-protein coupled receptors comprise one of the largest groups of signaling proteins; nearly 2% of our genes code for these receptors.

The *RGS2* gene has been associated with anxiety in animal models and in human correlational studies. *RGS2* knockout mice, both hetero- and homozygous, have increased hypertension (Heximer et al. 2003), excitability in CA1 neurons in the hippocampus (Oliveira-Dos-Santos et al. 2000), and anxiety-related traits (Yalcin et al. 2004). A targeted genome screen found modest evidence for linkage between markers in a region including *RGS2* and anxiety disorder proneness (Smoller et al. 2001). Recently, Leygraf and others (2006) found that polymorphisms in *RGS2* were associated with panic disorder in humans with the strongest association being observed for a haplotype containing SNPs rs4606 and rs3767488. Specifically, for rs4606, the authors found a higher prevalence of the C allele among panic disorder patients than controls. (Leygraf et al. 2006). A significant association has also been found between rs4606 and anxiety-related temperament, personality, and brain function, however, the G allele was found to be the risk allele (Smoller et al. 2008). Similarly, the distribution of rs4606 genotypes was found to be significantly different in suicide victims versus controls; the G-allele was more prevalent in suicide victims (Cui et al. 2007). Rs4606 is of particular interest because it is functional. Variation in rs4606 is

associated with variation in peripheral blood mononuclear cell and cultured fibroblast *RGS2* mRNA expression (Semplicini et al. 2006).

Materials & Methods

Data Collection and Sample

This article focuses on 607 participants in the 2004 Florida Hurricane Study who completed structured telephone interviews and provided saliva samples through the mail that yielded genotype data for the rs4606 polymorphism.

Verbal consent was obtained from the participants; the participants were sent a letter documenting the elements of verbal consent and providing them with contact information for the principal investigator. Participants who completed the diagnostic interview and returned saliva samples were paid \$20. The institutional review boards at the relevant institutions approved all procedures.

Assessment Procedure

Telephone interviews were conducted with a probability sample of English- and Spanish-speaking adults from telephone households in 38 counties in Florida within 6–9 months of the 2004 Florida hurricane season, between April 5 and June 12, 2005. Older adults (ages 60 and older) were oversampled to address research questions specific to this age group. Sample selection and telephone interviewing was performed by Schulman, Ronca, Bucuvalas, Inc., a national survey research firm with considerable experience conducting diagnostic interviews by telephone (Galea et al. 2002). Random-digit-dial procedures were used to locate households within the sampling frame. Respondents were randomly selected by using the most recent birthday method when multiple eligible adults were present within a household. Assessments were conducted via highly structured interview using computer-assisted telephone interview (CATI) format. The CATI format uses a computer program that projects instructions and questions on a computer screen that are read verbatim by interviewers to participants over the telephone. Interviewers then recorded participants' responses on a computer with closed-ended response options. Supervisors randomly monitored interviews in progress, thereby providing considerable quality control over data collection. Interviews averaged 26.5 minutes in length.

More details about the sampling procedure and methodology for the Florida Hurricane Study are provided elsewhere (Acierno et al. 2007; Acierno et al. 2006; Galea et al. 2006; Kilpatrick et al. 2007).

Presence of GAD since the hurricanes was measured using a slightly modified version of the SCID-IV (First et al. 1995) structured interview questions that correspond directly to DSM-IV criteria using yes/no response options. The diagnosis required excessive and poorly controlled anxiety and worry occurring more days than not for a period of 6–9 months ("since the hurricanes"), as well as three of six hallmark GAD symptoms (restlessness, fatigue, concentration problems, irritability, tension, and sleep disturbance). This scale showed good internal consistency in the current sample among individuals screening into the module (Cronbach's $\alpha = .85$).

PTSD since the hurricanes was assessed with the National Women's Study PTSD module, a widely used measure in population-based epidemiological research originally modified from the Diagnostic Interview Schedule. Research on the National Women's Study PTSD module has provided support for concurrent validity and several forms of reliability (e.g., temporal stability, internal consistency, diagnostic reliability [Ruggiero et al., 2006]). The National Women's Study PTSD module was validated in the DSM-IV PTSD field trial against the

Structured Clinical Interview for DSM (SCID), in which the interrater kappa coefficient was 0.85 for the diagnosis of PTSD and comparisons between scores on the National Women's Study PTSD module and the SCID yielded a kappa coefficient of 0.71 for current and 0.77 for lifetime PTSD (Kipatrick et al., 1998). Research also has found high correspondence between telephone versus in-person administration of the PTSD module as well as the depression module described below (Acierno et al., 2003). We operationalized PTSD diagnosis based on DSM-IV symptom requirements, including functional impairment.

Major depression since the hurricanes was assessed with modified questions from the SCID-IV. Following DSM-IV criteria, the respondents met criteria for major depression if they had five or more depressive symptoms for at least 2 weeks, including endorsement of one or both of the symptoms relating to depressed mood or loss of interest or pleasure. Past-year major depression identified by this measure is associated with lower reported work quality and mental health treatment-seeking after the addition of control for demographic characteristics, assault history, and PTSD diagnoses (Boscarino et al., 2004).

Hurricane exposure and social support were found to be associated with GAD in this sample (Acierno et al. 2007) and are included as covariates in logistic regression models. *Hurricane exposure* was assessed with five important indicators identified in previous research on Hurricanes Hugo and Andrew (Freedy et al. 1994) on the basis of their relation to posthurricane mental health functioning: 1) present during hurricane force winds or major flooding; 2) lack of adequate access to food, water, electricity, telephone, or clothing for a week or longer; 3) two or more hurricane-related losses of furniture, sentimental possessions, automobile, pets, crops, trees, or garden; 4) displacement from home for 1 week or longer; and 5) out-of-pocket losses of \$1,000 or more that were not reimbursed by insurance or other sources. Hurricane exposure was defined as having experienced none, one, two, three or four or more of these five indicators. High hurricane exposure was defined as having experienced two or more of these five indicators.

Social support during the 6 months before the hurricanes was assessed with a modified five-item version of the Medical Outcomes Study module (Sherbourne and Stewart 1991) that assesses emotional, instrumental, and appraisal social support (sample range=0–20; mean=15.9, SD=4.8). Low social support was operationalized as a score of 15 or less based on the cutoff score derived from prior work (Galea et al. 2002). This scale had good reliability (Chronbach's α =0.86).

Collection of DNA samples

Saliva samples were obtained via a mouthwash protocol and returned via mail to the Yale University laboratory where DNA extraction and analyses was conducted. Saliva samples were provided by 651 participants (42.2% response rate). Of these, valid genetic ancestry data were available for 623 cases (95.7%), and valid genotype data were available for 607 cases (93.2%). The likelihood of submitting a saliva sample did not differ in relation to sex, level of hurricane exposure, level of social support, or GAD status. Additional details on response rate and correlates of participation are summarized elsewhere (Galea et al. 2006).

Genotyping

DNA was extracted from saliva using PUREGENE (Gentra Systems, Minneapolis) kits. SNPs were genotyped with a fluorogenic 5'nuclease assay method ("TaqMan") using the ABI PRISM 7900 Sequence Detection System (ABI, Foster City, CA, USA). All genotypes were assayed in duplicate; discordant genotypes were discarded.

In addition, 36 markers were genotyped to provide ancestry information (Stein et al. 2004; Yang et al. 2005a; Yang et al. 2005b). We added one additional highly informative single

nucleotide polymorphism marker, *SLC24A5* (Lamason et al. 2005) to the panel described previously.

Ancestry Proportion Scores

Participants' ancestries were estimated by Bayesian cluster analysis with the marker panel described above on the procedures and STRUCTURE software developed by Pritchard and colleagues (Falush et al. 2003; Pritchard and Rosenberg 1999). For the STRUCTURE analysis, we specified the “admixture” and “allele frequencies correlated” models and used 100,000 burn-in and 100,000 Markov chain Monte Carlo iterations. Ancestry proportion scores were computed and included in analytic models to avoid spurious associations that can result from population stratification.

Statistical Analyses

We used chi-squared analyses to test whether demographic factors, low social support, hurricane exposure, the rs4606 polymorphism in *RGS2*, and post-hurricane major depression and posttraumatic stress disorder were associated with post-hurricane GAD. We then conducted logistic regression analyses to determine whether any observed association between rs4606 and GAD persisted after adjusting for sex, age, ancestral proportion scores, social support and hurricane exposure. To determine whether the association between *RGS2* and GAD was modified by level of stress exposure, we reran the logistic regression analyses including an interaction term for rs4606Xhurricane exposure. Because we had limited power to detect interaction effects, we also ran the logistic regression analyses stratified by hurricane exposure so we could observe whether the odds ratios for the association between rs4606 and GAD were comparable across high and low levels of hurricane exposure.

Results

Prevalence of GAD and RGS2 Genotype

Table 1 presents the demographic, trauma exposure, and clinical correlates of GAD. The prevalence of GAD since the hurricane was 6.8% (n=41). *RGS2* SNP rs4606 genotype frequencies were in Hardy-Weinberg equilibrium. Genotype frequencies were similar to those reported by Leygraf et al (2006). Female sex, age, and self-reported race were not significantly associated with GAD in unadjusted analyses. Both low social support and level of hurricane exposure significantly predicted GAD. Moreover, the chi-squared linear-by-linear association test supported a significant association between rs4606 and GAD. Since self-reported race was not significantly associated with GAD or with genotype frequencies ($\chi^2[2, n=587]=3.53, p=.47$), it is not likely that population stratification is a cause of false positive findings.

Association between GAD and RGS2 Genotype

Figure 1 shows the dose-response association between rs4606 and GAD. Table 2 shows the results of the logistic regression analyses. Both female sex and hurricane exposure were significantly associated with GAD. For rs4606, each ‘C’ allele at rs4606 was associated with a 100% increased risk of being diagnosed with post-hurricane GAD. Comparable results were found when we restricted the sample to Whites only (OR=1.91, 95% CI:1.00, 3.76). The effect of *RGS2* genotype on risk of GAD was similar for those with low (OR=2.61, 95% CI:0.88,7.78) and high (OR=1.87, 95% CI:0.84,4.12) hurricane exposure. Moreover, the interactions term for rs4606 X hurricane exposure was not significant (interaction OR=0.60, 95% CI: 0.32, 1.12).

Discussion

Our results support a significant association between SNP rs4606 in the *RGS2* gene and GAD in this epidemiologic sample of older adults exposed to the 2004 Florida Hurricane Season. These findings are consistent with those of Leygraf et al. (2006) who found a significant association between the same SNP and panic disorder in a clinical sample. In contrast, Smoller et al (2008) also found an association between rs4606 and anxiety-related traits but the G-allele was the risk allele. Taken together, these findings point toward a relevant polymorphism for anxiety-related disorders at the 3' end of the *RGS2* gene. However, the inconsistency with regard to the risk allele among studies raises questions as to whether rs4606 is the causal variant in *RGS2* as related to anxiety it is possible rs4606 is in linkage disequilibrium with some different functional variant, and the direction of LD differs by population. Alternatively, multiple testing, population stratification or some other form of confounding may account for the association between rs4606 and outcomes in some studies.

The biological mechanism via which variation in *RGS2* increases risk of GAD remains to be elucidated. It is noteworthy that suicide victims showed elevated *RGS2* immunoreactivity in postmortem brain tissue from their amygdala and prefrontal cortex (Cui et al. 2007). The neural circuitry connecting the amygdala and prefrontal cortex is thought to regulate emotional behavior, particularly responses to fear and anxiety inducing stimuli (Berkowitz et al. 2007). *RGS2* is expressed in these brain regions (Ingi and Aoki 2002; Ingi et al. 1998). A recent study demonstrated pathological activation the amygdala-prefrontal fear circuit among adolescents with GAD (McClure et al. 2007). Thus, the possibility that *RGS2* is associated with GAD via fear-circuit dysregulation should be explored in future studies.

We identified five potential limitations of the data presented in this article. First, individuals who relocated outside of the 38 hurricane-affected counties or lacked telephone service 6–9 months after the hurricanes were excluded from the study. However, telephone service was restored to the vast majority of Florida households several months before the study, and these hurricanes did not result in the type of massive evacuation outside the affected area that occurred following Hurricane Katrina. Second, the return rate of saliva samples was 42%, which is lower than optimal, but nonparticipation was not significantly related to major study variables of hurricane exposure, social support and GAD. Third, we used a slightly modified version of the SCID-IV administered by lay interviewers over the telephone rather than from in-person clinical interviews. However, there is substantial evidence supporting the reliability and validity of the SCID-IV (Ramirez Basco et al. 2000; Shear et al. 2000; Zanarini and Frankenburg 2001; Zanarini et al. 2000) and the validity of this measure as a lay screener for GAD (Spitzer et al. 2006; Swinson 2006). Moreover, the reliability of this measure in the current study was high (.85). Furthermore, prior research comparing telephone and in-person assessment with similar measures found no significant difference between the two methods (Acierno et al. 2003). Fourth, the study was cross-sectional, and therefore, we are unable to determine with certainty whether posthurricane GAD represents new onsets of the disorder or recurrence among individuals who might have had prior episodes. From an epidemiologic perspective, predisaster psychopathology could be a confounder in studies concerned with relations between risk factors and postdisaster outcomes. However, in the context of a genetic association analysis, we suggest that it is unlikely that predisaster psychopathology could substantially influence the results documented here. Finally, we note also that the number of affected individuals was relatively small, in the context of a reasonably large sample. Thus, we had very limited power to test for interaction effects.

Conclusions

This is the first genetic association study of GAD of which we are aware to examine gene-disorder relations within a sample that was recently exposed to a significant stressor. It is worth noting that the 6-month prevalence of GAD in this sample is 6.8% which is more than twice the 12-month prevalence in the general population (3.1%) and almost twice the lifetime prevalence (3.6%) among adults over aged 60 (Kessler et al. 2005a; Kessler et al. 2005b). The nature of the population makes this finding less vulnerable to ascertainment or recall biases than for a clinical sample or one using lifetime diagnoses. Additional research is needed to replicate our methodology and findings, but our results suggest that studying a recently disaster-exposed sample is both feasible and may improve power to find gene-disorder associations. Moreover, our findings in conjunction with data from mouse models and an emerging literature on *RGS2* and anxiety in humans suggest that pharmacological agents targeting *RGS2* may provide novel treatments for anxiety disorders.

Acknowledgments

This research was supported by NIMH grant MH05220, NIMH and National Institute of Nursing research grant MH07055, and National Institute on Drug Abuse grant K24 DA15105. Dr. Koenen is supported by NIMH grants K08-MH070627 and MH078928.

References

- Acierno R, Resnick H, Kilpatrick D, Stark-Riemer W. Assessing elder victimization--demonstration of a methodology. *Soc Psychiatry Psychiatr Epidemiol.* 2003; 38:644–653. [PubMed: 14614553]
- Acierno R, Ruggiero KJ, Galea S, Resnick HS, Koenen K, Roitzsch J, de Arellano M, Boyle J, Kilpatrick DG. Psychological sequelae resulting from the 2004 Florida hurricanes: implications for postdisaster intervention. *Am J Public Health.* 2007; 97 Suppl 1:S103–S108. [PubMed: 17413067]
- Acierno R, Ruggiero KJ, Kilpatrick DG, Resnick HS, Galea S. Risk and protective factors for psychopathology among older versus younger adults after the 2004 Florida hurricanes. *Am J Geriatr Psychiatry.* 2006; 14:1051–1059. [PubMed: 17035356]
- Berkowitz RL, Coplan JD, Reddy DP, Gorman JM. The human dimension: how the prefrontal cortex modulates the subcortical fear response. *Rev Neurosci.* 2007; 18:191–207. [PubMed: 18019606]
- Boscarino JA, Galea S, Adams RE, Ahern J, Resnick H, Vlahov D. Mental health service and medication use in New York City after the Sept 11, 2001, terrorist attack. *Psychiatr Serv.* 2004; 55:274–283. [PubMed: 15001728]
- Cui H, Nishiguchi N, Ivleva E, Yanagi M, Fukutake M, Nushida H, Ueno Y, Kitamura N, Maeda K, Shirakawa O. Association of *RGS2* Gene Polymorphisms with Suicide and Increased *RGS2* Immunoreactivity in the Postmortem Brain of Suicide Victims. *Neuropsychopharmacology.* 2007
- Falush D, Stephens M, Pritchard JK. Inference of population structure using multilocus genotype data: linked loci and correlated allele frequencies. *Genetics.* 2003; 164:1567–1587. [PubMed: 12930761]
- First, MB.; Spitzer, RL.; Williams, JBW.; Gibbon, M. Structured Clinical Interview for DSM-IV-Patient Edition (SCID-P). Washington, D.C.: American Psychiatric Press; 1995.
- Freedy JR, Saladin ME, Kilpatrick DG, Resnick HS, Saunders BE. Understanding acute psychological distress following natural disaster. *J Trauma Stress.* 1994; 7:257–273. [PubMed: 8012746]
- Galea S, Acierno R, Ruggiero K, Resnick H, Tracy M, Kilpatrick D. Social context and the psychobiology of posttraumatic stress. *Annals of the New York Academy of Sciences.* 2006; 1071:231–241. [PubMed: 16891574]
- Galea S, Ahern J, Resnick H, Kilpatrick D, Bucuvalas M, Gold J, Vlahov D. Psychological sequelae of the september 11 terrorist attacks in new york city. *New England Journal of Medicine.* 2002; 346:982–987. [PubMed: 11919308]
- Hettema JM, Neale MC, Kendler KS. A review and meta-analysis of the genetic epidemiology of anxiety disorders. *Am J Psychiatry.* 2001; 158:1568–1578. [PubMed: 11578982]

- Heximer SP. RGS2-mediated regulation of Gqalpha. *Methods Enzymol.* 2004; 390:65–82. [PubMed: 15488171]
- Heximer SP, Knutsen RH, Sun X, Kaltenbronn KM, Rhee MH, Peng N, Oliveira-dos-Santos A, Penninger JM, Muslin AJ, Steinberg TH, et al. Hypertension and prolonged vasoconstrictor signaling in RGS2-deficient mice. *J Clin Invest.* 2003; 111:445–452. [PubMed: 12588882]
- Hollinger S, Hepler JR. Cellular regulation of RGS proteins: modulators and integrators of G protein signaling. *Pharmacol Rev.* 2002; 54:527–559. [PubMed: 12223533]
- Ingi T, Aoki Y. Expression of RGS2, RGS4 and RGS7 in the developing postnatal brain. *Eur J Neurosci.* 2002; 15:929–936. [PubMed: 11906535]
- Ingi T, Krumins AM, Chidiac P, Brothers GM, Chung S, Snow BE, Barnes CA, Lanahan AA, Siderovski DP, Ross EM, et al. Dynamic regulation of RGS2 suggests a novel mechanism in G-protein signaling and neuronal plasticity. *J Neurosci.* 1998; 18:7178–7188. [PubMed: 9736641]
- Kar N, Bastia BK. Post-traumatic stress disorder, depression and generalised anxiety disorder in adolescents after a natural disaster: a study of comorbidity. *Clin Pract Epidemiol Ment Health.* 2006; 2
- Kendler KS, Hettema JM, Butera F, Gardner CO, Prescott CA. Life event dimensions of loss, humiliation, entrapment, and danger in the prediction of onsets of major depression and generalized anxiety. *Arch Gen Psychiatry.* 2003; 60:789–796. [PubMed: 12912762]
- Kessler RC, Berglund P, Demler O, Jin R, Merikangas KR, Walters EE. Lifetime prevalence and age-of-onset distributions of DSM-IV disorders in the National Comorbidity Survey Replication. *Arch Gen Psychiatry.* 2005a; 62:593–602. [PubMed: 15939837]
- Kessler RC, Chiu WT, Demler O, Merikangas KR, Walters EE. Prevalence, severity, and comorbidity of 12-month DSM-IV disorders in the National Comorbidity Survey Replication. *Arch Gen Psychiatry.* 2005b; 62:617–627. [PubMed: 15939839]
- Kessler RC, Keller MB, Wittchen HU. The epidemiology of generalized anxiety disorder. *Psychiatr Clin North Am.* 2001; 24:19–39. [PubMed: 11225507]
- Kilpatrick DG, Koenen KC, Ruggiero KJ, Acierno R, Galea S, Resnick HS, Roitzsch J, Boyle J, Gelernter J. The serotonin transporter genotype and social support and moderation of posttraumatic stress disorder and depression in hurricane-exposed adults. *Am J Psychiatry.* 2007; 164:1693–1699. [PubMed: 17974934]
- Kilpatrick, DG.; Resnick, HS.; Freedy, JR.; Pelcovitz, D.; Resick, PA.; Roth, S.; van der Kolk, B. The posttraumatic stress disorder field trial: evaluation of the PTSD construct—criteria A through E, in *DSM-IV Sourcebook*. Widiger, T.; Pincus, HA.; First, MB.; Ross, R.; Davis, W., editors. Washington, DC: American Psychiatric Press; 1998. p. 803-844.
- Lamason RL, Mohideen MA, Mest JR, Wong AC, Norton HL, Aros MC, Jurynek MJ, Mao X, Humphreville VR, Humbert JE, et al. SLC24A5, a putative cation exchanger, affects pigmentation in zebrafish and humans. *Science.* 2005; 310:1782–1786. [PubMed: 16357253]
- Leygraf A, Hohoff C, Freitag C, Willis-Owen SA, Krakowitzky P, Fritze J, Franke P, Bandelow B, Fimmers R, Flint J, et al. Rgs 2 gene polymorphisms as modulators of anxiety in humans? *J Neural Transm.* 2006; 113:1921–1925. [PubMed: 16736243]
- McClure EB, Monk CS, Nelson EE, Parrish JM, Adler A, Blair RJ, Fromm S, Charney DS, Leibenluft E, Ernst M, et al. Abnormal attention modulation of fear circuit function in pediatric generalized anxiety disorder. *Arch Gen Psychiatry.* 2007; 64:97–106. [PubMed: 17199059]
- Moffitt TE, Caspi A, Rutter M. Strategy for investigating interactions between measured genes and measured environments. *Arch Gen Psychiatry.* 2005; 62:473–481. [PubMed: 15867100]
- Oliveira-Dos-Santos AJ, Matsumoto G, Snow BE, Bai D, Houston FP, Whishaw IQ, Mariathasan S, Sasaki T, Wakeham A, Ohashi PS, et al. Regulation of T cell activation, anxiety, and male aggression by RGS2. *Proc Natl Acad Sci U S A.* 2000; 97:12272–12277. [PubMed: 11027316]
- Pritchard JK, Rosenberg NA. Use of unlinked genetic markers to detect population stratification in association studies. *Am J Hum Genet.* 1999; 65:220–228. [PubMed: 10364535]
- Ramirez Basco M, Bostic JQ, Davies D, Rush AJ, Witte B, Hendrickse W, Barnett V. Methods to improve diagnostic accuracy in a community mental health setting. *Am J Psychiatry.* 2000; 157:1599–1605. [PubMed: 11007713]

- Ruggiero KJ, Rheingold AA, Resnick HS, Kilpatrick DG, Galea S. Comparison of two widely used PTSD-screening instruments: implications for public mental health planning. *J Trauma Stress*. 2006; 19:699–707. [PubMed: 17075907]
- Semplicini A, Lenzini L, Sartori M, Papparella I, Calo LA, Pagnin E, Strapazzon G, Benna C, Costa R, Avogaro A, et al. Reduced expression of regulator of G-protein signaling 2 (RGS2) in hypertensive patients increases calcium mobilization and ERK1/2 phosphorylation induced by angiotensin II. *J Hypertens*. 2006; 24:1115–1124. [PubMed: 16685212]
- Shear MK, Greeno C, Kang J, Ludewig D, Frank E, Swartz HA, Hanekamp M. Diagnosis of nonpsychotic patients in community clinics. *Am J Psychiatry*. 2000; 157:581–587. [PubMed: 10739417]
- Sherbourne CD, Stewart AL. The MOS Social Support Survey. *Social Science and Medicine*. 1991; 32:705–714. [PubMed: 2035047]
- Shore JH, Tatum EL, Vollmer WM. Psychiatric reactions to disaster: the Mount St. Helens experience. *Am J Psychiatry*. 1986; 143:590–595. [PubMed: 3963245]
- Smoller JW, Acierno JS Jr, Rosenbaum JF, Biederman J, Pollack MH, Meminger S, Pava JA, Chadwick LH, White C, Bulzacchelli M, et al. Targeted genome screen of panic disorder and anxiety disorder proneness using homology to murine QTL regions. *Am J Med Genet*. 2001; 105:195–206. [PubMed: 11304837]
- Smoller JW, Paulus MP, Fagerness JA, Purcell S, Yamaki LH, Hirshfeld-Becker D, Biederman J, Rosenbaum JF, Gelernter J, Stein MB. Influence of RGS2 on anxiety-related temperament, personality, and brain function. *Arch Gen Psychiatry*. 2008; 65:298–308. [PubMed: 18316676]
- Spitzer RL, Kroenke K, Williams JB, Lowe B. A brief measure for assessing generalized anxiety disorder: the GAD-7. *Arch Intern Med*. 2006; 166:1092–1097. [PubMed: 16717171]
- Stein MB, Schork NJ, Gelernter J. A polymorphism of the beta1-adrenergic receptor is associated with low extraversion. *Biol Psychiatry*. 2004; 56:217–224. [PubMed: 15312808]
- Swinson RP. The GAD-7 scale was accurate for diagnosing generalized anxiety disorder. *Evid Based Med*. 2006; 11:184. [PubMed: 17213178]
- Yalcin B, Willis-Owen SA, Fullerton J, Meesaq A, Deacon RM, Rawlins JN, Copley RR, Morris AP, Flint J, Mott R. Genetic dissection of a behavioral quantitative trait locus shows that Rgs2 modulates anxiety in mice. *Nat Genet*. 2004; 36:1197–1202. [PubMed: 15489855]
- Yang BZ, Zhao H, Kranzler HR, Gelernter J. Characterization of a Likelihood Based Method and Effects of Markers Informativeness in Evaluation of Admixture and Population Group Assignment. *BMC Genet*. 2005a; 6:50. [PubMed: 16225681]
- Yang BZ, Zhao H, Kranzler HR, Gelernter J. Practical population group assignment with selected informative markers: Characteristics and properties of Bayesian clustering via STRUCTURE. *Genetic Epidemiology*. 2005b; 28:302–312. [PubMed: 15782414]
- Zanarini MC, Frankenburg FR. Attainment and maintenance of reliability of axis I and II disorders over the course of a longitudinal study. *Compr Psychiatry*. 2001; 42:369–374. [PubMed: 11559863]
- Zanarini MC, Skodol AE, Bender D, Dolan R, Sanislow C, Schaefer E, Morey LC, Grillo CM, Shea MT, McGlashan TH, et al. The collaborative longitudinal personality disorders study: Reliability of axis I and II diagnoses. *Journal of Personality Disorders*. 2000; 14:291–299. [PubMed: 11213787]

Figure 1.
Prevalence of generalized anxiety disorder by rs4606 genotype.

Table 1

Sample Descriptives and Unadjusted Association between Risk Factors and Current GAD Diagnosis

Variable	Overall (n=607) % (N)	Current GAD (n=41) % (N)	No GAD (n=566) % (N)	χ^2 (df=1)	p-value
Female sex	64.9 (394)	82.9 (34)	63.6 (360)	6.27	.12
Age less than 60 years	22.6 (137)	28.6 (11)	22.3 (126)	0.46	.50
Race/ethnicity					
White	90.6 (547)	82.9 (34)	91.1 (513)		
Other ^a	9.4 (57)	17.1 (7)	8.9 (50)	3.00	.08
Low Social Support	37.0 (224)	51.2 (21)	36.0 (203)	3.80	.05
Hurricane exposure				10.39 ^b	.001
0	9.7 (59)	4.9 (2)	10.1 (57)		
1	45.6 (277)	36.6 (15)	46.3 (262)		
2	23.7 (144)	17.1 (7)	24.2 (137)		
3	13.8 (84)	22.0 (9)	13.3 (75)		
4	7.1 (43)	19.5 (8)	6.2 (35)		
RGS2 rs4606				4.42 ^b	0.04
G/G	7.7 (47)	2.4 (1)	8.1 (46)		
G/C	36.4	26.8	37.1		

Variable	Overall (n=607) % (N)	Current GAD (n=41) % (N)	No GAD (n=566) % (N)	χ^2 (df=1)	p-value
C/C	(221) 55.8 (339)	(11) 70.7 (29)	(210) 54.8 (310)		
Major Depression	5.9 (36)	31.7 (13)	4.1 (23)	52.37	<.001
Posttraumatic Stress Disorder	3.1 (19)	19.5 (8)	1.9 (11)	38.92	<.001

GAD, generalized anxiety disorder; *RGS2*, regulator of g-protein signaling.

^aOther includes African- American, Hispanic, and Asian.

^b χ^2 linear-by linear association test.

Table 2

Final Logistic Regression Analysis of the Association between *RGS2* genotype and Generalized Anxiety Disorder (GAD)

Variable	GAD Diagnosis		
	Adjusted Odds Ratio	95% CI	p
Female Sex	2.6	1.1–6.1	0.025
Age less than 60 Years	1.1	0.5–2.3	0.878
Ancestral proportion score	0.5	0.62–3.6	0.452
Hurricane exposure	1.8	1.2–2.2	0.001
Low social support	1.9	1.0–3.6	0.065
<i>RGS2</i> rs4606 (C is risk allele)	2.0	1.1–3.8	0.026