Published in final edited form as:

J Eukaryot Microbiol. 2008 ; 55(6): 476–480. doi:10.1111/j.1550-7408.2008.00366.x.

How Epigenomics Contributes to the Understanding of Gene Regulation in *Toxoplasma gondii*¹

MATHIEU GISSOT and KAMI KIM

Albert Einstein College of Medicine, Departments of Medicine (Infectious Diseases) and Microbiology and Immunology, Bronx, New York 10461, USA

Abstract

How apicomplexan parasites regulate their gene expression is poorly understood. The complex life cycle of these parasites implies tight control of gene expression to orchestrate the appropriate expression pattern at the right moment. Recently, several studies have demonstrated the role of epigenetic mechanisms for control of coordinated expression of genes. In this review, we discuss the contribution of epigenomics to the understanding of gene regulation in *Toxoplasma gondii*. Studying the distribution of modified histones on the genome provides chromatin modifications to gene expression or gene repression. In particular, coincident trimethylated lysine 4 on histone H3 (H3K4me3), acetylated lysine 9 on histone H3 (H3K9ac), and acetylated histone H4 (H4ac) mark promoters of actively transcribed genes. However, the presence of these modified histones at some non-expressed genes and other histone modifications at only a subset of active promoters implies the presence of other layers of regulation of chromatin structure in *T. gondii*. Epigenomics analysis provides a powerful tool to characterize the activation state of genomic loci of *T. gondii* and possibly of other Apicomplexa including *Plasmodium* or *Cryptosporidium*. Further, integration of epigenetic data with expression data and other genome-wide datasets facilitates refinement of genome annotation based upon experimental data.

Keywords

Apicomplexa; epigenomics; epigenetics; gene regulation; *Plasmodium*; *Toxoplasma*

THE life cycle of *T. gondii* is complex with multiple differentiation steps that are critical to survival of the parasite in its human and feline hosts (Kim and Weiss 2004). As for other parasitic Apicomplexa, the *Toxoplasma* life cycle can be divided into a sexual cycle and asexual cycle occurring in a feline definitive host and in warm blooded animal intermediate hosts, respectively (Dubey, Miller, and Frenkel 1970). Sexual stages are characterized by the production of macro- and microgametes that will form a zygote and lead to the development of oocysts, which ultimately produce sporozoites. The sporozoites are ingested and transform into tachyzoites in the intermediate host. The rapidly growing tachyzoites convert into the latent bradyzoite, which is thought to persist for the life of the host. As clinical disease is frequently due to reactivation of latent infection, tachyzoite-bradyzoite interconversion is a key transition during the infection of the intermediate host. Ingestion by the cat of tissue cysts containing bradyzoites completes the life cycle.

¹Invited presentation, *Toxoplasma* Symposium: Centenary Celebration of *Toxoplasma* Discovery, May 31, 2008, during the X International Workshops on Opportunistic Protists (IWOP-10), May 28–31, 2008 at the Courtyard Marriott Tremont, Boston, MA.

Corresponding Author: Kami Kim Ullmann, 1225 1300 Morris Park Avenue, Bronx, NY 10461, USA---Telephone number: +1 718 430 2611; FAX number: +1 718 430 8968; e-mail:kkim@aecom.yu.edu.

Transcriptome studies of the various stages of the life cycle of the parasite revealed a complex pattern of expression associated with each form of the parasite. Large scale expression sequence tag (EST) studies demonstrated that some transcripts were specifically expressed at different developmental stages of the life cycle (Ajioka et al. 1998; Li et al. 2004; Li et al. 2003; Manger et al. 1998). Further studies focused on the transcriptome analysis of the tachyzoite to bradyzoite transition, using a library of mutants deficient for this conversion step. cDNA microarray analysis of these bradyzoite mutants suggested that a hierarchical gene expression program may control the bradyzoite differentiation (Singh et al. 2002).

The most comprehensive transcriptome study of the life cycle of the parasite was performed using the sequential analysis of gene expression (SAGE) technique and followed the development of the VEG (type III) in vitro from sporozoites to tachyzoites and finally to bradyzoites. The investigators also studied the tachyzoites of three laboratory-adapted strains (RH, type I; Me49, type II; VEGmsg, type III) (Radke et al. 2005). Approximately 18% of genes were stage specific with co-regulated genes specific to each form of the life cycle. These co-regulated genes were scattered across the genome indicating the importance of the promoter and trans-acting factors for the regulation of each particular gene (Radke et al. 2005).

Unlike the kinetoplastid protozoan parasites, apicomplexan parasites including *T. gondii* seem to greatly rely on transcriptional regulation to control gene expression. Supporting this idea, most of the basal transcription machinery required for the transcriptional control of gene expression appears to be encoded by the *T. gondii* genome (Callebaut et al. 2005; Meissner and Soldati 2005). This basal transcriptional complex is present in most eukaryotes and is composed of RNA polymerase II and its associated general transcription factors (GTF). The activity of this complex is modulated by sequence-specific transcription factors (SSTF), *cis*-promoting elements present in promoters, as well the chromatin context. In addition, co-regulators interact with the chromatin state and the SSTFs connecting those actors to the basal transcription machinery (Thomas and Chiang 2006).

Until recently, SSTFs were regarded as poorly represented in apicomplexan genomes (Coulson, Hall, and Ouzounis 2004), leading to speculation that epigenetic or post-transcriptional mechanisms were the primary mode of gene regulation. However, the presence of unique *cis*-promoting elements in the intergenic sequences is more consistent with the presence of apicomplexan specific SSTFs with unrecognized structural features: the *cis* elements mapped to date appear unrelated to other eukaryotic transcription binding sites (Behnke et al. 2008; Bohne, Wirsing, and Gross 1997; Kibe et al. 2005; Ma et al. 2004; Van Poppel et al. 2006). Reanalysis of apicomplexan genomes led to the discovery of a plant-like family of transcription factors with an Arabidopsis APETALA2 like (AP2) DNA-binding domain (Balaji et al. 2005). Approximately 50 putative AP2 transcription factors have been found in the *T. gondii* genome (Iyer et al. 2008), and most appear to be expressed in *T. gondii* tachyzoites (MG and KK, unpubl. data). Further studies are needed to decipher their role in gene regulation, but the AP2 DNA-binding domains in *Plasmodium* and *Cryptosporidium* were recently reported to bind specific DNA motifs (De Silva et al. 2008).

The conservation of the structural features of chromatin modifiers and remodelers facilitated the identification of a wide array of enzymes able to modify or remodel chromatin (Sullivan and Hakimi 2006). Similarly, a large number of predicted proteins encompass domains able to mediate the interaction with modified histones. Surprisingly, the *T. gondii* genome encodes histone modifier enzymes that are absent in yeast (Sullivan and Hakimi 2006), suggesting an important role for chromatin structure in *T. gondii* gene regulation.

Epigenetics in *Toxoplasma gondii*

Cytosine methylation at CpG dinucleotides mediates silencing of promoter activity in higher eukaryotes (Fazzari and Greally 2004). The presence of methylated cytosine plays a role in gene silencing in unicellular eukaryotes, such as *Dictyostelium* (Katoh et al. 2006) and *Entamoeba* (Fisher, Siman-Tor, and Ankri 2004), but is absent in the *Saccharomyces cerevisiae* genome. The cytosine-5 DNA methyltransferases (DNMT) are enzymes able to catalyze the methylation of cytosine in the DNA context. The DNMT2 family is conserved in lower eukaryotes and catalyzes DNA cytosine methylation in *Dictyostelium* (Katoh et al. 2006) and *Entamoeba* (Fisher et al. 2004), mainly at repetitive elements and retroposons. We discovered a homolog of the DNMT2 family of DNMTs in the *T. gondii* genome, but were unable to find methylated cytosine in the genomic DNA (Gissot et al. 2008). Although, we cannot exclude the presence of methylated cytosine in the genomic DNA of other stages of the *T. gondii* life cycle, the absence of this modified base in *Cryptosporidium* and *Plasmodium*, together with the high sensitivity of the methodology used, led us to conclude that DNA methylation is unlikely to be a major mechanism for regulation of gene expression in Apicomplexa during their asexual phases (Gissot et al. 2008).

Chromatin structure plays a crucial role in number of nuclear processes from DNA repair to transcription. Histones within the nucleosome participate in the packaging of DNA and therefore affect all stages of transcription. Nucleosomes possess dynamic properties that are regulated by the modification and the remodeling of chromatin (Li, Carey, and Workman 2007). Methylation, acetylation, ubiquitinylation, Small Ubiquitinlike Modifier modification (SUMOylation), and phosphorylation are common modifications found in eukaryotic histones. They influence the binding of the histone to the DNA as well as the ability to recruit other proteins (Li et al. 2007).

There are the four canonical histones (H2A, H2B, H3, H4) in *T. gondii* along with variant histones. Histone modifications have been reported at different residues of H3 and H4 of *T. gondii* as assessed by reactivity with commercially available antibodies to conserved histone modifications (Table 1) (Bhatti et al. 2006; Gissot et al. 2007; Saksouk et al. 2005; Smith et al. 2005). The best-characterized modifications of histones include methylation and acetylation of lysines and methylation of arginines or lysines located at the N-termini. Most of these histone modifications are believed to be activation marks in eukaryotes and correlate to gene expression (Gissot et al. 2007). One putative repression mark (i.e. trimethylation of the lysine 20 of H4) was found in *T. gondii*: this mark failed to correlate with promoters of repressed genes but instead localized to repetitive DNA and telomeres (Sautel et al. 2007). Another repressive mark, the trimethylation of lysine 9 on H3 (i.e. H3K9me3), a heterochromatin marker in other eukaryotes, was found in *Plasmodium* in the promoters of inactive *var* genes (Chookajorn et al. 2007; Lopez-Rubio et al. 2007), and was also reported in *T. gondii* (Sautel et al. 2007). However, its distribution does not seem to correlate with promoters in *T. gondii* (Sautel et al. 2007).

Although the repertoire of chromatin remodeling enzymes is similar in *Plasmodium* and *T. gondii* (Sullivan and Hakimi 2006), it is not yet clear how conserved individual histone modifications are among the Apicomplexa or whether these modifications correspond to similar states of gene activation. Mass spectrometry analysis of the *T. gondii* histones would be invaluable for precise mapping of histone modifications that are present in the epigenetic arsenal of the parasite. Further, the biochemical activities and specificity of candidate chromatin remodeling enzymes have been experimentally verified for only a subset of these factors (Bhatti et al. 2006; Saksouk et al. 2005; Sautel et al. 2007; Smith et al. 2005).

Epigenomics and modified histones in gene regulation in *Toxoplasma gondii*

Initial studies investigating the role of modified histones in *T. gondii* focused upon a small number of developmentally regulated genes (Saksouk et al. 2005; Sautel et al. 2007) using a combination of chromatin immunoprecipitation (ChIP) and single locus PCR. The low resolution and sensitivity of PCR on a small number of loci limits the conclusions of those studies.

We investigated the epigenetic organization and the transcription patterns of a contiguous 1% of the *T. gondii* genome using ChIP and hybridization on a tiling microarray (ChIP on chip) (Gissot et al. 2007). This custom oligonucleotide microarray contained 12,995 50mer features tiling a 650-kb region of chromosome 1b of the RH strain, with an average resolution of one oligonucleotide every 50 bp. Using commercial antibodies specific for modified histones, we observed 52 clear, discrete, and coincident peaks of H4ac, H3K9ac, and H3K4me3 within the 650-kb region tiled on the microarray. Interestingly, a fourth modified histone (H3R17me2) was only enriched in a small subset of promoters. Overall, 51 of the 52 regions with a cluster of H3K9ac, H4ac, and H3K4me3 peaks had significant cDNA hybridization signals adjacent to them, demonstrating a strong correlation between the presence of modified histone peaks and gene expression. In contrast, two regions of clustered genes, with stage-specific expression based on EST data and no expression at the tachyzoite stage, were shown to lack these modified histone peaks (Gissot et al. 2007).

One region with modified histones peaks was associated with a gene encoding a putative UV-induced double strand break-repair protein that had no detectable mRNA expression. This gene might be poised for rapid expression as demonstrated in human T cells (Roh et al. 2006). Gene expression can be rapidly regulated at certain loci with the inactive PolII associated with the promoter, ready to induce transcription (Margaritis and Holstege 2008). Up to 20% of the regions with modified histones were not associated with a corresponding expressed gene in a survey of human promoters, suggesting that this may be a common mechanism of gene regulation (Kim et al. 2005).

In yeast, phosphorylation of the serine 5 (S5) of the PolII is a marker of elongation since this post-translational modification occurs upon activation and initiation of elongation (Cho 2007). A survey of the distribution of the PolII S5 phosphorylation across the yeast genome revealed that PolII is poised at the promoters of rapidly expressed genes, where the peaks of H3K9ac and H4ac are, but not in the body of the gene (Radonjic et al. 2005). In eukaryotes, the pausing of PolII is dependant upon negative factors (5,6-dichloro-1-beta-D-ribofuranosylbenzimidazole [DRB] sensitivity-inducing factor and the Negative Elongation factor), and factors able to relieve the inhibition of the PolII activity as the positive transcription-elongation factor-b complex (Margaritis and Holstege 2008). Predicted proteins with low but significant similarity with those factors are present in *T. gondii* and therefore may participate in this regulatory mechanism. However, understanding the extent or usage of this type of regulation in *T. gondii* awaits the results of a genome-wide survey of the distribution of modified histones, the localization of different phosphorylated forms of PolII, and the correlation of both of these with gene expression.

We have now expanded our ChIP-chip studies to the whole genome of *T. gondii* using each of the three archetypic strains (RH, type I; PLK/ME49, type II; CEP/CTG, type III) hybridized to a genome-wide tiled microarray based on the genomic sequence of the ME49 strain (www.toxodb.org). H3K9ac and H3K4me3 ChIP-chip of tachyzoite DNA from each of these strains shows that there is a strong concordance between gene expression and the presence of modified histone peaks for the majority of loci (MG, Fajardo, Weiss, Fiser, KK,

unpubl. data). Further, most genes with concordant H3K9ac and H3K4me3 marks are expressed in all three strains. However, we also identified regions of the genome where H3K9ac peaks were not associated with a H3K4me3 peak (MG, Fajardo, Weiss, Fiser, KK, unpubl. data), whereas in our pilot survey of 1% of the RH strain genome, all H3K9ac peaks mapped to genomic regions with a H3K4me3 peak (Gissot et al. 2007).

The presence of acetylation peaks that are not accompanied by methylation peaks may reflect differences in biological states between the PLK and RH strains, since the RH strain is less prone to biological variation at the tachyzoite stage due to its inability to differentiate. Since H3K4me3 and H3K9ac peaks are concordant in the large majority of cases, it is less likely that this difference is technical and due to higher intrinsic background signal associated with the antibodies used to recognize the H3K9ac modification vs. the H3K4me3 antibodies, but further analysis is ongoing. High basal levels of histone acetylation of bradyzoite-specific promoters has been reported for low passage Type I, II, and III strains that readily differentiate to bradyzoites, supporting the concept of “poised” bradyzoite promoters that have some of the initial chromatin modifications associated with active gene expression (Behnke et al. 2008).

Since the recruitment of histone acetyl transferase is believed to precede the presence of PolII (Bhaumik and Green 2001), the unique H3K9ac marks might represent genes ready to be activated (or genes that are in the process of being repressed). However, the temporal sequence of deposition of histone modifications on *T. gondii* chromatin is not yet clear. Moreover, this string of events may depend upon which histone modifying complex is targeted to the promoter and consequently might be promoter-dependant. Therefore, it is difficult to make definitive conclusions without an in-depth analysis of the nature of the genes associated with histone acetylation and the genes' patterns of expression at different stages of the life cycle. Both histone acetylation and methylation are reversible processes and the regulation of histone deacetylase and demethylase enzyme activities are aspects of chromatin modification that have not been fully explored in the Apicomplexa. Further mapping of histone modifications will determine if there are regions of the genome that represent specialized regions, such as enhancers or insulators that participate in the regulation of genes situated in other parts of the genome (Heintzman et al. 2007).

Epigenomics allows the mapping of active promoters

The regions encompassing the peaks of modified histones were located at the 5'-end of predicted genes and strongly correlated to gene expression. To determine if these regions had promoter function, we cloned genomic regions encompassing the H3K9ac and H4ac peaks, which are undistinguishable one from another, and tested their promoter activity using a luciferase reporter. These regions were able to directionally drive the expression of the reporter (Gissot et al. 2007) illustrating the utility of modified histones to define the location of the active promoters on this portion of the genome.

The size of the peaks is on average 1,550 bp, much longer than the previously well-described promoters driving the expression of genes like *GRA1*, *DHFR* or α -tubulin in *T. gondii* (558, 467, and 530 bp respectively). The ChIP technique relies on the hybridization of DNA fragments of 500--1,000 bp generated by sonication of the genomic DNA. The presence of one modified histone at the extremity of the fragment is theoretically enough to yield a positive signal, artificially increasing the size of regions with positive signal. An alternative ChIP-chip technique, based on the hybridization of Micrococcal nuclease-digested DNA, permits a higher resolution, selecting only the histone-bound DNA, which is protected from digestion (Clark and Shen 2006). The regions defined by ChIP-chip probably

do not represent the minimal promoters and encompass more sequence than is strictly needed to drive the expression of the genes.

Our working model envisions the chromatin and the SSTF as co-regulators of the transcriptional activity of promoters in *T. gondii* (Fig. 1). Whether the SSTF recruits the chromatin modifiers and remodelers (Fig. 1A) or the opposite (Fig. 1B), both types of factors act in a coordinated fashion to stabilize the PolII complex at the transcriptional start site.

Epigenomics allows refined gene annotation

Since modified histones can predict the distribution of active promoters in the genome, we can identify loci with discordance between the placement of active promoters and the localization of predicted genes. We found two mispredicted genes in the portion of the ChrIb we studied (Gissot et al. 2007). Therefore, the mapping of active promoters can inform an experimentally-based refinement of the annotation of the genome. Expansion of our analysis to multiple strains and different stages of the life cycle will also enhance accurate gene annotation on other less explored forms of the parasite.

We have also shown that promoters have directional activity and that the distribution of the H3K4me3 is predictive of the orientation of transcription. Using this feature, we discovered two regions where no gene was predicted for which we detected modified histones and mRNA hybridization (Gissot et al. 2007). One of the regions encoded a putative cytochrome oxidase subunit III gene and the other had no obvious open reading frame and could represent a non-coding RNA. Since the 5'-end of the gene is the most difficult to predict (Bajic et al. 2006), locating the promoter of genes represents valuable experimental data that can be incorporated into algorithms used for gene prediction and genome annotation.

Conclusion

The use of epigenomics will allow a better understanding of gene regulation in Apicomplexa and provide invaluable data for experimentally-based refinement of genome annotation. Although the general picture has been investigated with the first characterization of the *T. gondii* tachyzoite epigenome, preliminary data suggest that the modified histone code is more complex than previously envisioned: there are H3K9ac peaks unlinked to a H3K4me3 peak and only a subset of promoters have associated H3R17me2. Deciphering the *T. gondii* epigenome during developmental transitions, as well as in different strains, is likely to provide further insights into the mechanisms of gene activation and repression in *T. gondii*.

Acknowledgments

M. G. was supported by a Philippe Foundation Fellowship. This work was supported by NIH NIAID RO1 AI060496 to K. K.

LITERATURE CITED

- Ajioka JW, Boothroyd JC, Brunk BP, Hehl A, Hillier L, Manger ID, Marra M, Overton GC, Roos DS, Wan KL, Waterston R, Sibley LD. Gene discovery by EST sequencing in *Toxoplasma gondii* reveals sequences restricted to the Apicomplexa. *Genome Res.* 1998; 8:18–28. [PubMed: 9445484]
- Bajic VB, Brent MR, Brown RH, Frankish A, Harrow J, Ohler U, Solovyev VV, Tan SL. Performance assessment of promoter predictions on ENCODE regions in the EGASP experiment. *Genome Biol.* 2006; 7(S3):1–13.

- Balaji S, Babu MM, Iyer LM, Aravind L. Discovery of the principal specific transcription factors of Apicomplexa and their implication for the evolution of the AP2-integrase DNA binding domains. *Nucleic Acids Res.* 2005; 33:3994–4006. [PubMed: 16040597]
- Behnke MS, Radke JB, Smith AT, Sullivan WJ Jr. White MW. The transcription of bradyzoite genes in *Toxoplasma gondii* is controlled by autonomous promoter elements. *Mol. Microbiol.* 2008; 68:1502–1518. [PubMed: 18433450]
- Bhatti MM, Livingston M, Mullapudi N, Sullivan WJ Jr. Pair of unusual GCN5 histone acetyltransferases and ADA2 homologues in the protozoan parasite *Toxoplasma gondii*. *Eukaryot. Cell.* 2006; 5:62–76. [PubMed: 16400169]
- Bhaumik SR, Green MR. SAGA is an essential in vivo target of the yeast acidic activator Gal4p. *Genes Dev.* 2001; 15:1935–1945. [PubMed: 11485988]
- Bohne W, Wirsing A, Gross U. Bradyzoite-specific gene expression in *Toxoplasma gondii* requires minimal genomic elements. *Mol. Biochem. Parasitol.* 1997; 85:89–98. [PubMed: 9108551]
- Callebaut I, Prat K, Meurice E, Mornon JP, Tomavo S. Prediction of the general transcription factors associated with RNA polymerase II in *Plasmodium falciparum*: conserved features and differences relative to other eukaryotes. *BMC Genomics.* 2005; 6:100. [PubMed: 16042788]
- Cho EJ. RNA polymerase II carboxy-terminal domain with multiple connections. *Exp. Mol. Med.* 2007; 39:247–254. [PubMed: 17603278]
- Chookajorn T, Dzikiowski R, Frank M, Li F, Jiwani AZ, Hartl DL, Deitsch KW. Epigenetic memory at malaria virulence genes. *Proc. Natl. Acad. Sci. USA.* 2007; 104:899–902. [PubMed: 17209011]
- Clark DJ, Shen CH. Mapping histone modifications by nucleosome immunoprecipitation. *Methods Enzymol.* 2006; 410:416–430. [PubMed: 16938564]
- Coulson RM, Hall N, Ouzounis CA. Comparative genomics of transcriptional control in the human malaria parasite *Plasmodium falciparum*. *Genome Res.* 2004; 14:1548–1554. [PubMed: 15256513]
- De Silva EK, Gehrke AR, Olszewski K, Leon I, Chahal JS, Bulyk ML, Llinas M. Specific DNA-binding by apicomplexan AP2 transcription factors. *Proc. Natl. Acad. Sci. USA.* 2008; 105:8393–8401. [PubMed: 18541913]
- Dubey JP, Miller NL, Frenkel JK. *Toxoplasma gondii* life cycle in cats. *J. Am. Vet. Med. Assoc.* 1970; 157:1767–1770. [PubMed: 5530378]
- Fazzari MJ, Grealley JM. Epigenomics: beyond CpG islands. *Nat. Rev. Genet.* 2004; 5:446–455. [PubMed: 15153997]
- Fisher O, Siman-Tov R, Ankri S. Characterization of cytosine methylated regions and 5-cytosine DNA methyltransferase (Ehmet) in the protozoan parasite *Entamoeba histolytica*. *Nucleic Acids Res.* 2004; 32:287–297. [PubMed: 14715927]
- Gissot M, Choi SW, Thompson RF, Grealley JM, Kim K. *Toxoplasma gondii* and *Cryptosporidium parvum* lack detectable DNA cytosine methylation. *Eukaryot. Cell.* 2008 Epub ahead of publication 4 January 2008.
- Gissot M, Kelly KA, Ajioka JW, Grealley JM, Kim K. Epigenomic modifications predict active promoters and gene structure in *Toxoplasma gondii*. *PLoS Pathog.* 2007; 3:e77. [PubMed: 17559302]
- Heintzman ND, Stuart RK, Hon G, Fu Y, Ching CW, Hawkins RD, Barrera LO, Van Calcar S, Qu C, Ching KA, Wang W, Weng Z, Green RD, Crawford GE, Ren B. Distinct and predictive chromatin signatures of transcriptional promoters and enhancers in the human genome. *Nat. Genet.* 2007; 39:311–318. [PubMed: 17277777]
- Iyer LM, Anantharaman V, Wolf MY, Aravind L. Comparative genomics of transcription factors and chromatin proteins in parasitic protists and other eukaryotes. *Int. J. Parasitol.* 2008; 38:1–31. [PubMed: 17949725]
- Katoh M, Curk T, Xu Q, Zupan B, Kuspa A, Shaulsky G. Developmentally regulated DNA methylation in *Dictyostelium discoideum*. *Eukaryot. Cell.* 2006; 5:18–25. [PubMed: 16400165]
- Kibe MK, Coppin A, Dendouga N, Oria G, Meurice E, Mortuaire M, Madec E, Tomavo S. Transcriptional regulation of two stage-specifically expressed genes in the protozoan parasite *Toxoplasma gondii*. *Nucleic Acids Res.* 2005; 33:1722–1736. [PubMed: 15784612]

- Kim K, Weiss LM. *Toxoplasma gondii*: the model apicomplexan. *Int. J. Parasitol.* 2004; 34:423–432. [PubMed: 15003501]
- Kim TH, Barrera LO, Zheng M, Qu C, Singer MA, Richmond TA, Wu Y, Green RD, Ren B. A high-resolution map of active promoters in the human genome. *Nature.* 2005; 436:876–880. [PubMed: 15988478]
- Li B, Carey M, Workman JL. The role of chromatin during transcription. *Cell.* 2007; 128:707–719. [PubMed: 17320508]
- Li L, Crabtree J, Fischer S, Pinney D, Stoeckert CJ Jr. Sibley LD, Roos DS. ApiEST-DB: analyzing clustered EST data of the apicomplexan parasites. *Nucleic Acids Res.* 2004; 32 Database issue:D326–328. [PubMed: 14681425]
- Li L, Brunk BP, Kissinger JC, Pape D, Tang K, Cole RH, Martin J, Wylie T, Dante M, Fogarty SJ, Howe DK, Liberator P, Diaz C, Anderson J, White M, Jerome ME, Johnson EA, Radke JA, Stoeckert CJ Jr. Waterston RH, Clifton SW, Roos DS, Sibley LD. Gene discovery in the Apicomplexa as revealed by EST sequencing and assembly of a comparative gene database. *Genome Res.* 2003; 13:443–454. [PubMed: 12618375]
- Lopez-Rubio JJ, Gontijo AM, Nunes MC, Issar N, Hernandez Rivas R, Scherf A. 5' flanking region of var genes nucleate histone modification patterns linked to phenotypic inheritance of virulence traits in malaria parasites. *Mol. Microbiol.* 2007; 66:1296–1305. [PubMed: 18028313]
- Ma YF, Zhang Y, Kim K, Weiss LM. Identification and characterisation of a regulatory region in the *Toxoplasma gondii* hsp70 genomic locus. *Int. J. Parasitol.* 2004; 34:333–346. [PubMed: 15003494]
- Manger ID, Hehl A, Parmley S, Sibley LD, Marra M, Hillier L, Waterston R, Boothroyd JC. Expressed sequence tag analysis of the bradyzoite stage of *Toxoplasma gondii*: identification of developmentally regulated genes. *Infect. Immun.* 1998; 66:1632–1637. [PubMed: 9529091]
- Margaritis T, Holstege FC. Poised RNA polymerase II gives pause for thought. *Cell.* 2008; 133:581–584. [PubMed: 18485867]
- Meissner M, Soldati D. The transcription machinery and the molecular toolbox to control gene expression in *Toxoplasma gondii* and other protozoan parasites. *Microbes Infect.* 2005; 7:1376–1384. [PubMed: 16087378]
- Radke JR, Behnke MS, Mackey AJ, Radke JB, Roos DS, White MW. The transcriptome of *Toxoplasma gondii*. *BMC Biol.* 2005; 3:26. [PubMed: 16324218]
- Radonjic M, Andrau JC, Lijnzaad P, Kemmeren P, Kockelkorn TT, van Leenen D, van Berkum NL, Holstege FC. Genome-wide analyses reveal RNA polymerase II located upstream of genes poised for rapid response upon *S. cerevisiae* stationary phase exit. *Mol. Cell.* 2005; 18:171–183. [PubMed: 15837421]
- Roh TY, Cuddapah S, Cui K, Zhao K. The genomic landscape of histone modifications in human T cells. *Proc. Natl. Acad. Sci. USA.* 2006; 103:15782–15787. [PubMed: 17043231]
- Saksouk N, Bhatti MM, Kieffer S, Smith AT, Musset K, Garin J, Sullivan WJ Jr. Cesbron-Delauw MF, Hakimi MA. Histone-modifying complexes regulate gene expression pertinent to the differentiation of the protozoan parasite *Toxoplasma gondii*. *Mol. Cell Biol.* 2005; 25:10301–10314. [PubMed: 16287846]
- Sautel CF, Cannella D, Bastien O, Kieffer S, Aldebert D, Garin J, Tardieux I, Belrhali H, Hakimi MA. SET8-mediated methylations of histone H4 lysine 20 mark silent heterochromatic domains in apicomplexan genomes. *Mol. Cell Biol.* 2007; 27:5711–5724. [PubMed: 17562855]
- Singh U, Brewer JL, Boothroyd JC. Genetic analysis of tachyzoite to bradyzoite differentiation mutants in *Toxoplasma gondii* reveals a hierarchy of gene induction. *Mol. Microbiol.* 2002; 44:721–733. [PubMed: 11994153]
- Smith AT, Tucker-Samaras SD, Fairlamb AH, Sullivan WJ Jr. MYST family histone acetyltransferases in the protozoan parasite *Toxoplasma gondii*. *Eukaryot. Cell.* 2005; 4:2057–2065. [PubMed: 16339723]
- Sullivan WJ Jr. Hakimi MA. Histone mediated gene activation in *Toxoplasma gondii*. *Mol. Biochem. Parasitol.* 2006; 148:109–116. [PubMed: 16644030]
- Thomas MC, Chiang CM. The general transcription machinery and general cofactors. *Crit. Rev. Biochem. Mol. Biol.* 2006; 41:105–178. [PubMed: 16858867]

Van Poppel NF, Welagen J, Vermeulen AN, Schaap D. The complete set of *Toxoplasma gondii* ribosomal protein genes contains two conserved promoter elements. *Parasitology*. 2006; 133:19–31. [PubMed: 16674839]

Fig. 1.

Hypothetical model representing the major actors of transcriptional control in the apicomplexan *Toxoplasma gondii*. Two alternative, but not mutually exclusive, models are presented: **(A)** Sequence specific transcription factors (SSTF) are recruited to the promoter and interact with the chromatin remodeling and modification machinery to stabilize the PolII complex at the transcriptional start site (TSS). **(B)** Chromatin remodelers and modifiers are targeted to the promoters and promote the binding of SSTF to their binding site, therefore enabling the stabilization of the PolII complex at the TSS. In both cases, chromatin remodeling complexes, the PolII complex, and specific transcription factors act in concert to regulate gene expression. For different loci the components of the chromatin remodeling complex or SSTF complex may vary, allowing for further layers of regulation.

Table 1

Modified histones investigated in *T. gondii* using Chromatin Immunoprecipitation

Modified Histone	ChIP	ChIP-chip	ChIP-chip peaks	Localization of the peaks	Correlation to gene expression
H3K4me1	⁺ _{(a)(b)}	⁺ _(a)	-	NA	NA
H3K4me2	⁺ _{(a)(b)}	⁺ _(a)	-	NA	NA
H3K4me3	⁺ _{(a)(b)}	⁺ _(a)	+	Promoters ^{(a)(b)}	+
H3K9ac	⁺ _{(a)(b)}	⁺ _(a)	+	Promoters ^{(a)(b)}	+
H3K9me3	⁺ _(c)	⁺ _(d)	NA	Telomeres and repetitive sequences ^(c)	-
H3R17me2	⁺ _{(a)(b)}	⁺ _(a)	+	Promoters ^{(a)(b)}	⁺ _{((a) - subset only)}
H3K18ac	+	ND	NA	NA	+
H4ac	+	+	+	Promoters ^{(a)(b)}	+
H4K20me3	⁺ _(c)	⁺ _(d)	NA	Telomeres and repetitive sequences ^(c)	-
H4K20me1	⁺ _(c)	⁺ _(d)	NA	NA	-

Histone modifications reported in the literature for *T. gondii* include methylation and acetylation of histone H3 and histone H4 with studies reported using chromatin immunoprecipitation followed by PCR amplification of genomic loci of interest (ChIP) or chromatin immunoprecipitation followed by hybridization to a tiled genomic array (ChIP-chip)

NA: not applicable

ND: not determined

Studies reporting the histone modification are:

^(a) (Gissot et al. 2007)

^(b) (Saksouk et al. 2005)

^(c) (Sautel et al. 2007), localization of peaks inferred from results with SET8-myc

^(d) MG and KK, unpublished results