Published in final edited form as:

Invest Ophthalmol Vis Sci. 2008 November ; 49(11): 5048–5054. doi:10.1167/iovs.08-1877.

Quantitative optical coherence tomography findings in various subtypes of neovascular age-related macular degeneration

Sandra Liakopoulos, M.D.^{1,2}, Sharel Ongchin, B.S.¹, Alok Bansal, M.D.¹, Sandeep Msutta¹, Alexander C. Walsh, M.D.¹, Paul G. Updike, M.S.¹, and Srinivas R. Sadda, M.D.¹

¹Doheny Image Reading Center, Doheny Eye Institute, Keck School of Medicine of the University of Southern California, Los Angeles, CA

²Department for vitreoretinal surgery, Center of Ophthalmology, University of Cologne, Germany

Abstract

Purpose—To compare the volume of various spaces visible on optical coherence tomography (OCT) images in different angiographic lesion subtypes of neovascular age-related macular degeneration (AMD).

Methods—Sixty-six cases of previously un-treated, active subfoveal choroidal neovascularization (CNV) associated with AMD were retrospectively collected. CNV lesions were classified as occult with no classic CNV, minimally classic CNV, predominantly classic CNV or CNV lesions with associated retinal angiomatous proliferation (RAP). Corresponding StratusOCT image sets were analyzed by trained graders using previously validated custom software (OCTOR) which allows manual placement of boundaries on OCT B-scans. Spaces delineated by these boundaries included the neurosensory retina, subretinal fluid, subretinal tissue, and pigment epithelial detachments (PEDs). Volume measurements were calculated by the software and compared between groups.

Results—Minimally and predominantly classic CNV membranes demonstrated subretinal tissue on OCT in all cases and appeared to show a significantly greater volume of subretinal tissue than occult membranes. Subretinal fluid was present in all of the predominantly classic cases. A PED was visible in all of the occult CNV cases in our study, demonstrating less retinal thickening and significantly greater PED volumes than minimally and predominantly classic CNV lesions. Lesions associated with RAP showed the highest percentage of cystoid spaces.

Conclusions—OCT and angiography provide complementary information regarding CNV lesions. Quantitative analysis of OCT images allows for an improved understanding of the anatomic characteristics of angiographically-defined CNV lesion subtypes.

Keywords

Optical coherence tomography; quantitative image analysis; choroidal neovascularization; age-related macular degeneration; classic; occult

Introduction

Age-related macular degeneration (AMD) is the most common cause of blindness in North America and Europe.¹ In neovascular AMD, two different patterns of choroidal neovascularization (CNV) have been defined according to their appearance on fluorescein

angiography (FA): classic CNV and occult CNV.² Occult CNV has been further subdivided into fibrovascular pigment epithelial detachment (FVPED) and late leakage of undetermined source (LLUS). Recently, the presence of retinal angiomatous proliferation (RAP) or retinal anastomosis to the lesion (RAL) has been indentified in some cases with CNV.³

Classic CNV, occult CNV, and RAP are believed to have different characteristics with respect to the progression of the disease over time and the impact on visual function. Large randomized clinical trials have demonstrated significant differences in treatment effects in minimally classic CNV (less than 50% of the lesion occupied by classic CNV), predominantly classic CNV (more than 50% of the lesion occupied by classic CNV) and occult with no classic CNV.^{2,4,5} The presence of RAP also seems to have an impact of the course of the disease as well as the treatment outcome.³

Although FA remains an important tool in the diagnosis of CNV lesions, optical coherence tomography (OCT) has become an increasingly important component for the management of patients with neovascular AMD. Retinal thickening, subretinal fluid, subretinal tissue and pigment epithelial detachments (PEDs) are all well visualized on OCT B-scans, and have permitted the study of the anatomic features of CNV lesions. Classic CNV is believed to frequently correspond to histological type 2 CNV (CNV between retinal pigment epithelium (RPE) and retina),^{6,7} which may appear on OCT as subretinal tissue.⁸⁻¹² Occult CNV is believed to correspond to histological type 1 CNV (between Bruch's membrane and RPE),^{6,7} which may in FVPED (but not in LLUS) appear as a PED on OCT.⁸⁻¹² Leakage on FA caused by active CNV usually appears as intraretinal, subretinal or sub-RPE fluid accumulation on OCT.⁹⁻¹²

To date, the appearance of the different lesion types on OCT are only qualitatively described.⁹⁻¹² In this study, we analyzed OCT images of eyes with neovascular AMD using computer assisted manual grading software to quantify various spaces visible on OCT images for a more precise comparison of the characteristics of different angiographic CNV lesion subtypes.

Methods

Data collection

Data from 66 eyes of 66 consecutive patients who were about to initiate therapy for neovascular AMD were retrospectively collected. To be eligible for the study, eyes were required to have previously untreated, active subfoveal CNV due to AMD and StratusOCT imaging as well as FA imaging performed on the same date. In addition, the entire CNV lesion as defined on FA had to fall within a 6 mm - diameter circle centered on the fovea..Active CNV was defined as the presence of hemorrhage, or evidence of vision loss (of at least one line of Snellen acuity) or lesion growth within the prior three months.

Approval for the collection and analysis of the data was obtained from the Institutional Review Board of the University of Southern California. The research adhered to the tenets set forth in the Declaration of Helsinki.

Color fundus photographs and FA images were obtained using a Topcon 50IX camera with an 11 megapixel sensor (Topcon Medical Systems, Paramus, New Jersey). The standard imaging protocol of the Doheny Imaging Unit was followed in all cases, consisting of color stereo pairs of the ETDRS fields 1 and 2, followed by multiple angiographic frames of field 2 during the transit phase, and stereo pairs of Field 2 in the mid (<3 min) and late phases (5-10 min) of the angiogram.

OCT images were obtained using the “Radial Lines” protocol in which 6 high-resolution 6 mm B-scans (transverse resolution of 512 A-scans per B-scan) were obtained using a single StratusOCT machine (Carl Zeiss Meditec, Dublin, CA, USA). Although many published reports on OCT have utilized the “Fast Macular Scan” protocol (with a lower transverse resolution of 128 A-scans per B-scan), the high-resolution “Radial Lines” protocol has been the standard method used in the Doheny ophthalmic imaging unit due to the enhanced morphologic detail provided. The Fast Macular Scan protocol was used only when photographers were unable to obtain adequate high-resolution images, most commonly in patients with unstable fixation or poor cooperation. Photographers attempted to center the OCT B-scans on the fovea in all cases.

Data for each case were exported to disk using the export feature available in the StratusOCT version 4.0 analysis software.

Computer-assisted OCT grading software

The software used for OCT analysis (termed OCTOR) was written by Doheny Image Reading Center (DIRC) software engineers to facilitate viewing and manual grading.¹³ The OCTOR software is freely accessible at <https://www.diesel.la>, and has been described and validated in previous reports.^{13,14} The software imports data exported from the StratusOCT machine, and allows the grader to use a computer mouse to draw various boundaries in the retinal cross-sectional images.

After the grader draws the required layers in each of the six OCT B-scans, the software calculates the distance in pixels between the manually drawn boundary lines for each of the various defined spaces. Using the dimensions of the B-scan image, the calculated pixels are converted into microns to yield a thickness measurement at each location. The thickness at all unsampled locations between the radial lines is then interpolated to generate a thickness or volume map analogous to the Stratus OCT output. The interpolation algorithm and inter-grader reproducibility have previously been validated.^{13,14}

Analogous to the StratusOCT software, the OCTOR software provides a report showing the calculated thickness and volume values for the 9 Early Treatment Diabetic Retinopathy Study (ETDRS) macular subfields, and the mean and standard deviation of the foveal center point (FCP) thickness (Figure 1). In contrast to the Stratus output, the OCTOR software provides separate maps for the various macular spaces (e.g. retina, subretinal fluid, subretinal tissue, PEDs).

Grading procedure

FA images and OCT scans were analyzed by certified graders (SL, SRS) at the DIRC.^{2,14} FA's were viewed and graded using DIRC FA grading software equipped with standard planimetric tools to allow calculation of the area and greatest linear dimension (GLD) of any closed-loop figure/structure drawn by the grader using a computer mouse. CNV lesions were graded according to the modified Macular Photocoagulation Study (MPS) grading protocol utilized in the TAP and VIP studies.² Eyes were classified into four groups according to their appearance on FA: 1) predominantly classic CNV, 2) minimally classic CNV, 3) occult with no classic CNV, and 4) CNV associated with RAP.² The area (measured in MPS disc areas) and GLD (measured in mm) of the CNV lesions were calculated. Additionally, the presence of hemorrhage and hard exudates on color fundus photography was noted.

OCT B-scans were analyzed using computer-assisted manual grading software OCTOR.¹⁴ Graders were masked as to the angiographic lesion subtype when analyzing the OCT image set. Boundaries drawn in each of the six OCT B-scans included the internal limiting membrane

(ILM), outer border of the photoreceptors, borders of subretinal fluid and subretinal tissue (if present), inner surface of the RPE, and the estimated normal position of the RPE layer (in cases with RPE elevation). All boundaries were drawn in accordance with the standard OCT grading protocol of the DIRC.¹⁴ The presence of cystoid spaces in the retina was also noted.

After completion of the grading, the OCTOR software was used to calculate the volume for the various spaces: retina, subretinal fluid, subretinal tissue, and PED. In addition, a combined parameter, the inner retinal surface height from the choroid (IHC), was also calculated as the combination of all four individual spaces.

Statistical methods

Volume (mm^3) values for subfields 1-9 were collected for each space and each case, and mean and standard deviation values were calculated across all cases. The area of the CNV lesion was measured using TAP grading criteria in each case. The volume values for each case (and for every space) were computed as a ratio by dividing the volume by the area measurement, effectively normalizing the values with respect to lesion area. Values were compared between groups. Statistical analysis was performed using commercially available software (SigmaPlot 2004 for Windows, ver. 9.01; Systat Software Inc., Erkrath, Germany). For the comparison between groups, ANOVA on ranks test (for comparison between more than 2 groups), the t-test or Mann-Whitney rank sum test (for comparison between 2 groups) was used, depending on whether the data matched the pattern expected in a population with a normal distribution. P values less than 0.05 were considered statistically significant.

Results

Predominantly classic CNV was diagnosed on fluorescein angiography in 11 cases, minimally classic CNV in 23 cases, occult with no classic CNV in 24 cases, and CNV associated with RAP in 8 cases.

OCT measurements for all subtypes are summarized in tables 1 to 2. The different CNV lesion subtypes showed the following characteristics on OCT

Predominantly classic CNV was associated with retinal thickening (defined as total retinal volume greater than 7mm^3) and subretinal fluid accumulation on OCT in all 11 cases (100%, mean retinal volume $8.47 \pm 1.78 \text{ mm}^3$, mean volume of subretinal fluid $0.24 \pm 0.20 \text{ mm}^3$). Cystoid spaces were present in 9 cases (82%). Subretinal tissue was detected on OCT in all cases (100%, mean subretinal tissue volume $0.56 \pm 0.55 \text{ mm}^3$). Five eyes also demonstrated additionally PEDs (45%, mean PED volume $0.38 \pm 0.97 \text{ mm}^3$). In 3 of those 5 cases with PED on OCT (60%), occult CNV could be identified on FA as a lesion component, whereas none of the eyes without PED on OCT showed occult components on FA.

Minimally classic CNV (Figure 2) was associated with retinal thickening in 20 out of 23 cases (87%, mean retinal volume $7.86 \pm 0.91 \text{ mm}^3$), retinal cystoid spaces in 17 eyes (74%) and subretinal fluid accumulation in 21 cases (91%, mean volume $0.5 \pm 0.6 \text{ mm}^3$). All 23 eyes showed subretinal tissue on OCT (100%, mean volume $0.37 \pm 0.41 \text{ mm}^3$) with 16 cases additionally demonstrating PEDs on OCT as well as occult components on FA (70%, mean volume $0.32 \pm 0.36 \text{ mm}^3$). In 7 cases (33%), no PED was present on OCT; however, LLUS could be detected on FA in 4 of those 7 cases (57%) indicating occult CNV without RPE elevation.

Occult with no classic CNV (Figure 3) revealed retinal thickening in 21 out of 24 cases (88%, mean retinal volume $7.62 \pm 0.72 \text{ mm}^3$) and cystoid spaces in 14 cases (58%). Subretinal fluid accumulation was present in 19 eyes (79%, mean volume $0.2 \pm 0.5 \text{ mm}^3$). In all cases, PEDs were seen on OCT (100%, mean volume $0.86 \pm 0.73 \text{ mm}^3$). Subretinal tissue was additionally

present on OCT in 7 eyes (29%, mean volume $0.07 \pm 0.14 \text{ mm}^3$). In 3 of those 7 eyes, thick subretinal hemorrhage was seen on color photographs. In 4 eyes, no obvious correlation for the subretinal tissue seen on OCT could be identified on FA or color photographs despite blocked fluorescence possibly indicating pigment or other material.

CNV lesions with RAP (Figure 4) showed significant retinal thickening with retinal cystoid spaces in all 8 cases (100%, mean retinal volume $8.54 \pm 1.76 \text{ mm}^3$). Subretinal fluid was present in 4 eyes (50%, mean volume $0.42 \pm 0.69 \text{ mm}^3$), subretinal tissue in 7 eyes (88%, mean volume $0.11 \pm 0.1 \text{ mm}^3$) and PEDs in 7 eyes (88%, mean volume $0.59 \pm 0.62 \text{ mm}^3$).

Comparison between the different lesion types (Figures 5 and 6 and Tables 2 and 3)

Area measurements of the total CNV lesion size on fluorescein angiography did not reveal statistically significant differences between lesion type groups (mean area 3.68 ± 2.69 MPS disc areas for predominantly classic CNV, 3.58 ± 3.01 MPS disc areas for minimally classic CNV, 4.22 ± 2.49 MPS disc areas for occult without classic CNV, and 2.96 ± 2.14 MPS disc areas for CNV with RAP lesions).²

On OCT, predominantly and minimally classic CNV lesions showed a statistically greater volume of subretinal tissue than occult CNV membranes ($p < 0.001$). Occult with no classic CNV lesions showed a statistically greater PED volume than minimally or predominantly classic cases ($p < 0.001$). Subretinal fluid was most frequently present in predominantly classic CNV; however, there was no statistically significant difference in the volume of subretinal fluid between the different angiographic lesion subtypes. No difference was found between groups regarding total retinal volume and volume of the IHC. However, when comparing the group of occult with no classic CNV cases with eyes containing some classic CNV (predominantly and minimally classic cases grouped together), occult with no classic CNV lesions were found to have a significantly lower retinal volume.

Discussion

AMD is a multifactorial disease which presents with a variety of different phenotypes.^{1,2,3,15} CNV in neovascular AMD has been classified into classic and occult components based on angiographic characteristics.² Classic CNV is believed to correspond histologically to type 2 CNV which is located in the subretinal space,^{6,7} and therefore appears early and sharply demarcated on FA.² Leakage of classic CNV progresses throughout the angiogram, as leaking fluid can easily accumulate in the subretinal space (a potential space in the eye) or diffuse through the retinal layers. Occult CNV is believed to correspond histologically to type 1 CNV, which is located in the sub-RPE space.^{6,7} The fibrovascular tissue may create disturbances in the overlying RPE, causing occult CNV to often appear as areas of stippled hyperfluorescence on FA.² If the outer retinal barrier is disrupted by the disease process, leakage from type 1 CNV may pass through the RPE layer and spread into the subretinal space and/or neurosensory retina. On FA, CNV lesions are classified as predominantly classic CNV, minimally classic CNV, and occult with no classic CNV based on the proportions of the lesion components classic and occult CNV, thick hemorrhage, blocked fluorescence and serous PED.²

Angiography and OCT represent complementary imaging modalities, both of which display different information. FA reveals information about the integrity of the inner and outer retinal barrier and the presence of pathological vessels. OCT visualizes the arrangement and reflectivity of retinal and subretinal layers as well as the axial distribution of pathological fluid or tissue within the retina, the subretinal or sub-RPE space. Interpretation of both imaging modalities together allows for a more precise understanding of pathology. For example, a CNV lesion is defined as minimally classic CNV according to the TAP criteria², if classic CNV is visible on FA but other lesion components occupy more than 50% of the total lesion size. Those

other lesion components include not only occult CNV, but also blocked fluorescence, thick hemorrhage or serous PED, because these features could possibly obscure the boundaries of CNV. In these cases, OCT could potentially be helpful in confirming or refuting the presence of additional “hidden” CNV, such as the presence of an underlying fibrovascular PED (Figure 2).

Vice versa, FA may be helpful in the interpretation of subretinal tissue on OCT, as subretinal hyperreflective material on OCT does not necessarily correspond to type 2 CNV, but may be attributed to blood, lipid, fibrin, or pigment, which may appear as blocked fluorescence on FA. As it may not be possible to distinguish CNV from other lesion components of similar reflectivity based on OCT characteristics alone, one OCT grading convention in this study was to assign the generic label “subretinal tissue” to all hyperreflective material identified in the subretinal space. Thus, “subretinal tissue volume” in this study is not equivalent to “type 2 CNV volume”.

This study aims to quantitatively analyze and compare the appearance of various angiographic subtypes of active CNV lesions on OCT. Consistent with our hypothesis, eyes with classic CNV components demonstrated the highest percentage and greatest volume of subretinal tissue on OCT images, whereas eyes with occult with no classic CNV showed the highest percentage and greatest volume of PEDs. This is in accordance with qualitative observations in previously published reports.⁹⁻¹² To the authors knowledge, however, this is the first report providing quantitative measurements for comparison of various spaces on OCT other than the neurosensory retina. Interestingly, the mean PED volume in eyes with occult with no classic CNV was higher than in eyes with minimally classic and predominantly classic CNV, although in our series, 20 out of 23 eyes with minimally classic CNV and 3 out of 11 eyes with predominantly classic CNV appeared to have occult components visible on angiography. One possible explanation for this finding is that occult with no classic CNV lesions (with sub-RPE disease alone), were more likely to grow in the sub-RPE space for an extended period of time before manifesting with symptoms or other signs, such as a breakthrough of exudation into the subretinal space, prompting diagnosis and treatment.

Intra- and subretinal fluid accumulation have been observed in eyes with angiographically active CNV lesions and have been assumed to be signs of activity.¹⁶⁻¹⁸ Therefore, some studies proposed OCT guided re-treatment criteria for neovascular AMD based on retinal volume and presence of subretinal fluid.^{17,18} Indeed, in our study, most eyes demonstrated retinal cystoid spaces and / or subretinal fluid accumulation. Four eyes, however, presented with leakage on FA but without subretinal fluid or cystoid spaces on OCT; a finding also described by Eter et al.¹⁹ Three of those 4 eyes were classified as occult with no classic CNV, one as minimally classic CNV. It seems possible that leakage on FA may not be visible as intra- or subretinal fluid on OCT, if the fluid is pumped out by the RPE just as quickly as it leaks or if a healthy outer retinal barrier prevents leakage from type 1 CNV from diffusing through the RPE layer. However, since time domain (StratusOCT) imaging only captures a small area of the retina, it is also possible that SRF or cystoid spaces were in fact present, but not captured by one of the 6 OCT B-scans. Vice versa, intra- and subretinal fluid accumulation on OCT often correlates with leakage or pooling on FA, but such leakage may not be clearly apparent on the commonly used angiographic time frames (up to 10 minutes) if the source of leakage is very small and leakage is slow. Retinal degeneration in old or chronic CNV lesions also may also be associated with loss of neuronal retinal tissue and formation of cystoid spaces without the presence of leakage from active CNV on FA.

In our study, eyes with predominantly classic CNV showed the highest percentage of subretinal fluid. Eyes with occult with no classic CNV demonstrated the lowest percentage of cystoid edema and the smallest volumes of retina and SRF compared to the other groups. However

there was no significant difference between groups regarding the volume of retina and subretinal fluid. This is in accordance with findings from Hughes et al, who did not observe a difference in retinal edema and cystoid spaces between eyes with classic and occult CNV.¹² It seems likely that classic CNV may cause retinal or subretinal fluid accumulation, whereas the outer retinal barrier may prevent leakage from occult CNV from passing through the RPE layer in some cases. Occult CNV may eventually induce a break down of the outer retinal barrier. RAP lesions in contrast are very likely to show cystoid edema, as leakage from the pathological vessels (presumed to be intraretinal) can easily diffuse within the retina. In our study, eyes with RAP demonstrated the highest percentage of cystoid edema.

Limitations of this study include the relatively small sample, and the interpolation of data from 6 radial line StratusOCT B-scans in order to generate volume data. In addition, in eyes with poor fixation, the 6 scans will likely not be centered in one point, causing additional errors in volume calculation. The new Fourier Domain OCT (FDOCT) devices could potentially address some of these concerns. The high scanning speeds achievable with FDOCT devices confer greater resistance to fixation errors and permit a higher sampling density of the scanned region, requiring less interpolation between sampled locations. Unfortunately, automated segmentation algorithms in FDOCT devices have yet to be validated for complex diseases such as neovascular AMD, and grading every B-scan in a dense FDOCT volume acquisition for a large number of patients is not feasible. Manual correction of a more limited set of scans would appear to be a reasonable compromise until the segmentation software associated with this new technology matures.

In conclusion, quantitative analysis of OCT images allows for a more precise analysis of anatomic differences between various angiographic subtypes of CNV lesions in neovascular AMD. Quantitative analysis of changes in retinal volume, subretinal fluid, subretinal tissue or PED volumes over time in clinical trials may improve our understanding about differences between classic CNV, occult CNV and RAP in the natural course of the disease or in response to treatment. These findings may also facilitate the development of a new classification system for CNV lesions which incorporates both angiographic and OCT morphologic characteristics to more precisely characterize lesion subtypes in neovascular AMD.

Acknowledgements

Supported in part by NIH Grant EY03040 and NEI Grant R01 EY014375

References

1. Bressler NM, Bressler SB, Congdon NG, Ferris FL 3rd, Friedman DS, Klein R, et al. Age-Related Eye Disease Study Research Group. Potential public health impact of Age-Related Eye Disease Study results: AREDS report no. 11. *Arch Ophthalmol* 2003;121:1621–1624. [PubMed: 14609922]
2. Barbazetto I, Burdan A, Bressler NM, et al. Treatment of Age-Related Macular Degeneration with Photodynamic Therapy Study Group; Verteporfin in Photodynamic Therapy Study Group. Photodynamic therapy of subfoveal choroidal neovascularization with verteporfin: fluorescein angiographic guidelines for evaluation and treatment--TAP and VIP report No. 2. *Arch Ophthalmol* 2003;121:1253–68. [PubMed: 12963608]
3. Kuhn D, Meunier I, Soubrane G, et al. Imaging of chorioretinal anastomoses in vascularized retinal pigment epithelium detachments. *Arch Ophthalmol* 1995;11:1392–8. [PubMed: 7487600]
4. Brown DM, Kaiser PK, Michels M, et al. ANCHOR Study Group. Ranibizumab versus verteporfin for neovascular age-related macular degeneration. *N Engl J Med* 2006;355:1432–44. [PubMed: 17021319]
5. Rosenfeld PJ, Brown DM, Heier JS, et al. MARINA Study Group. Ranibizumab for neovascular age-related macular degeneration. *N Engl J Med* 2006;355:1419–31. [PubMed: 17021318]

6. Submacular Surgery Trials Research Group. Comparison of 2D reconstructions of surgically excised subfoveal choroidal neovascularization with fluorescein angiographic features: SST report No. 15. *Ophthalmology* 2006;113:279.e1–279.e5. [PubMed: 16406533]
7. Grossniklaus HE, Wilson DJ, Bressler SB, et al. Clinicopathologic studies of eyes that were obtained postmortem from four patients who were enrolled in the submacular surgery trials: SST Report No. 16. *Am J Ophthalmol* 2006;141:93–104. [PubMed: 16386982]
8. Fukuchi T, Takahashi K, Uyama M, Matsumura M. Comparative study of experimental choroidal neovascularization by optical coherence tomography and histopathology. *Jpn J Ophthalmol* 2001;45:252–258. [PubMed: 11369374]
9. Sandhu SS, Talks SJ. Correlation of optical coherence tomography, with or without additional colour fundus photography, with stereo fundus fluorescein angiography in diagnosing choroidal neovascular membranes. *Br J Ophthalmol* 2005;89:967–70. [PubMed: 16024845]
10. Sato T, Iida T, Hagimura N, Kishi S. Correlation of optical coherence tomography with angiography in retinal pigment epithelial detachment associated with age-related macular degeneration. *Retina* 2004;24:910–4. [PubMed: 15579989]
11. Coscas F, Coscas G, Souied E, Tick S, Soubrane G. Optical coherence tomography identification of occult choroidal neovascularization in age-related macular degeneration. *Am J Ophthalmol* 2007;144:592–9. [PubMed: 17698019]
12. Hughes EH, Khan J, Patel N, Kashani S, Chong NV. In vivo demonstration of the anatomic differences between classic and occult choroidal neovascularization using optical coherence tomography. *Am J Ophthalmol* 2005;139:344–6. [PubMed: 15733999]
13. Sadda SR, Joeres S, Wu Z, Romano P, Collins AT, Walsh AC. Error correction and quantitative sub-analysis of optical coherence tomography data using computer-assisted grading. *Invest Ophthalmol Vis Sci* 2007;48:839–48. [PubMed: 17251486]
14. Joeres S, Tsong JW, Updike PG, et al. Reproducibility of quantitative optical coherence tomography sub-analysis in neovascular age-related macular degeneration. *Invest Ophthalmol Vis Sci* 2007;48:4300–7. [PubMed: 17724220]
15. Grossniklaus HE, Green WR. Choroidal neovascularization. *Am J Ophthalmol* 2004;137:496–503. [PubMed: 15013874]
16. Kaiser PK, Blodi BA, Shapiro H, Acharya NR, MARINA Study Group. Angiographic and optical coherence tomographic results of the MARINA study of ranibizumab in neovascular age-related macular degeneration. *Ophthalmology* 2007;114:1868–75. [PubMed: 17628683]
17. Fung AE, Lalwani GA, Rosenfeld PJ, et al. An optical coherence tomography-guided, variable dosing regimen with intravitreal ranibizumab (Lucentis) for neovascular age-related macular degeneration. *Am J Ophthalmol* 2007;143:566–83. [PubMed: 17386270]
18. van Velthoven ME, de Smet MD, Schlingemann RO, Magnani M, Verbraak FD. Added value of OCT in evaluating the presence of leakage in patients with age-related macular degeneration treated with PDT. *Graefes Arch Clin Exp Ophthalmol* 2006;244:1119–23. [PubMed: 16523305]
19. Eter N, Spaide RF. Comparison of fluorescein angiography and optical coherence tomography for patients with choroidal neovascularization after photodynamic therapy. *Retina* 2005;25:691–6. [PubMed: 16141855]

Figure 1.
OCTOR map demonstrating the volume and thickness of all spaces calculated by the OCTOR software after manual grading of all 6 OCT B-scans.

Figure 2.

Fluorescein angiographic (FA) frames [early phase (A), mid phase (B), late phase (C)] of a case with minimally classic CNV. The corresponding OCT B-scan (D, scan direction marked by red arrow in C) demonstrates subretinal fluid, subretinal tissue and a pigment epithelial detachment (PED). Of note, OCT illustrates that the PED (presumably corresponding to Type 1 CNV) continues underneath the subretinal tissue (presumably corresponding to Type 2 CNV), while on FA, only a circular area of stippled hyperfluorescence is visible indicating occult CNV, while the central part of the PED is hidden underneath classic CNV.

Figure 3.

Fluorescein angiographic (FA) frames [early phase (A), mid phase (B), late phase (C)] of a case with occult with no classic choroidal neovascularization (CNV). The corresponding optical coherence tomography (OCT) B-Scan (D, scan direction marked by red arrow in C) shows a pigment epithelial detachment and subretinal fluid.

Figure 4.

Fluorescein angiographic (FA) frames [early phase (A), mid phase (B), late phase (C)] of a case demonstrating a stereoscopically visible retinal angiomatous proliferation (RAP) connected to occult choroidal neovascularization (CNV). The corresponding optical coherence tomography (OCT) B-Scan (C) shows cystoid retinal edema and subretinal tissue. The RAP lesion itself is not captured by the OCT B-scan.

Figure 5.

Comparison between angiographic lesion subtypes regarding the total volume of the different spaces visible on Optical Coherence Tomography (OCT) images. Eyes with retinal angiomatous proliferation (RAP) showed the largest total retinal volume, eyes with classic choroidal neovascularization (CNV) components showed the largest volume of subretinal tissue and eyes with occult with no classic CNV showed the largest volumes of pigment epithelial detachments.

*Statistically significant difference between groups ($p < 0.001$)

A: Total volume values of each space used for analysis

B: Total volume of each space divided by area of the total CNV lesion (volume / area ratio) used for analysis

Figure 6.

Comparison between angiographic lesion subtypes regarding the total volume of the different spaces visible on Optical Coherence Tomography (OCT) images. There was no statistically significant difference between groups regarding the total volume of the retina or the inner retinal height from the choroid (IHC). Eyes with retinal angiomatous proliferation (RAP) showed the largest total retinal volume, eyes with classic choroidal neovascularization (CNV) components showed the largest volume of subretinal tissue and eyes with occult with no classic CNV showed the largest volumes of pigment epithelial detachments.

*Statistically significant difference between groups ($p < 0.001$)

A: Total volume values of each space used for analysis

B: Total volume of each space divided by area of the total CNV lesion (volume / area ratio) used for analysis

Table 1

Number (%) of different optical coherence tomography (OCT) features in different angiographic lesion subtypes

	<i>Occult with no classic (n=24)</i>	<i>Minimally classic (n=23)</i>	<i>Predominantly classic (n=11)</i>	<i>RAP stage III (n=8)</i>
Subretinal Fluid	19 (79%)	21 (91%)	11 (100%)	4 (50%)
Subretinal Tissue	7 (29%)	23 (100%)	11 (100%)	7 (88%)
Pigment Epithelial Detachment	24 (100%)	16 (70%)	5 (45%)	7 (88%)
Cystoid Edema	14 (58%)	17 (74%)	9 (82%)	8 (100%)

Table 2
Volume measurements (mm^3) of various spaces on optical coherence tomography (OCT) B-scans

<i>OCT parameter</i>		<i>Occult with no classic</i> (n=24)	<i>Minimally classic</i> (n=23)	<i>Predominantly classic</i> (n=11)	<i>RAP stage III</i> (n=8)
Retina	Mean $\pm$ SD (Min - Max)	7.62 $\pm$ 0.72 (6.28 - 9.33)	7.86 $\pm$ 0.91 (6.49 - 9.98)	8.47 $\pm$ 1.78 (7.01 - 13.07)	8.54 $\pm$ 1.76 (6.80 - 12.39)
Subretinal Fluid	Mean $\pm$ SD (Min - Max)	0.24 $\pm$ 0.50 (0.00 - 2.39)	0.50 $\pm$ 0.59 (0.00 - 2.41)	0.24 $\pm$ 0.20 (0.01 - 0.58)	0.42 $\pm$ 0.69 (0.00 - 1.87)
Subretinal Tissue	Mean $\pm$ SD (Min - Max)	0.07 $\pm$ 0.14 (0.00 - 0.51)	0.37 $\pm$ 0.41 (0.01 - 1.76)	0.56 $\pm$ 0.55 (0.07 - 1.56)	0.11 $\pm$ 0.10 (0.00 - 0.23)
Pigment Epithelial Detachment	Mean $\pm$ SD (Min - Max)	0.86 $\pm$ 0.73 (0.01 - 2.78)	0.32 $\pm$ 0.36 (0.00 - 1.21)	0.38 $\pm$ 0.97 (0.00 - 3.27)	0.59 $\pm$ 0.62 (0.00 - 1.75)
Inner Retinal Height from Choroid	Mean $\pm$ SD (Min - Max)	8.79 $\pm$ 1.28 (6.72 - 12.27)	9.05 $\pm$ 1.60 (6.87 - 13.33)	9.64 $\pm$ 3.14 (7.51 - 18.46)	9.66 $\pm$ 2.01 (7.70 - 13.57)

Table 3

Volume ratios (mm³ of each space/ MPS DA of the total CNV lesion) of various spaces on optical coherence tomography (OCT) B-scans

<i>OCT parameter</i>		<i>Occult with no classic (n=24)</i>	<i>Minimally classic (n=23)</i>	<i>Predominantly classic (n=11)</i>	<i>RAP stage III (n=8)</i>
Retina	Mean ± SD (Min - Max)	2.58 ± 1.92 0.78 - 9.75	4.04 ± 3.56 0.72 - 14.91	5.37 ± 6.85 1.10 - 24.53	4.74 ± 3.46 1.50 - 10.69
Subretinal Fluid	Mean ± SD (Min - Max)	0.06 ± 0.09 0.00 - 0.30	0.28 ± 0.39 0.00 - 1.25	0.13 ± 0.16 0.00 - 0.57	0.13 ± 0.19 0.00 - 0.48
Subretinal Tissue	Mean ± SD (Min - Max)	0.01 ± 0.03 0.00 - 0.15	0.17 ± 0.19 0.00 - 0.79	0.17 ± 0.10 0.03 - 0.31	0.09 ± 0.10 0.00 - 0.27
Pigment Epithelial Detachment	Mean ± SD (Min - Max)	0.26 ± 0.25 0.00 - 0.91	0.09 ± 0.11 0.00 - 0.43	0.06 ± 0.13 0.00 - 0.44	0.16 ± 0.13 0.00 - 0.39
Inner Retinal Height from Choroid	Mean ± SD (Min - Max)	2.90 ± 2.08 0.93 - 10.56	4.58 ± 3.93 0.89 - 16.54	5.73 ± 7.02 1.44 - 25.37	5.12 ± 3.48 2.00 - 11.04