Published in final edited form as:

Tetrahedron Lett. 2007 November 19; 48(47): 8270–8273. doi:10.1016/j.tetlet.2007.09.129.

Asymmetric Synthesis of (+)-Iso-6-Cassine via Stereoselective Intramolecular Amidomercuration

Satwinder Singh, Om V. Singh, and Hyunsoo Han*

Department of Chemistry, The University of Texas at San Antonio, One UTSA Circle, San Antonio, TX 78249, USA

Abstract

The first asymmetric synthesis of (+)-iso-6-cassine is described. Lipase-catalyzed resolution, enantioselective Overman rearrangement, and diastereoselective intramolecular amidomercuration were used for the installation of the three stereocenters in (+)-iso-6-cassine, and cross-metathesis was employed for the attachment of the side-chain.

Keywords

intramolecular amidomercuration; 2,6-dialkylpiperidines; 2,6-disubstituted 3-piperidinols; double stereodifferentiation; (+)-iso-6-cassine

Substituted piperidine ring systems constitute the core structures of many chemically and biologically important compounds,¹ and thus have been a focus of many synthetic studies.² Particularly, electrophile-induced stereoselective heterocyclization reactions have provided a convenient tool for the construction of such ring systems from the corresponding linear precursors.³ In this context, we recently reported that Hg(II)-mediated tandem Overman rearrangement and intramolecular amidomercuration reactions could be a very efficient method for the stereoselective formation of *cis*- and *trans*- 2,6-dialkyl piperidine ring systems.⁴ The highlight of the report is dominant *trans* stereoselection by *N*-trichloroacetyl (TCA) protection group in the Hg(II)-mediated intramolecular amidomercuration reactions of 5-alkenyl amides (H at the C4 in eq 1).

*Corresponding author. Tel.: +1-210-458-4958; fax: +1-210-458-7428; e-mail: E-mail: hyunsoo.han@utsa.edu..

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

A literature survey revealed that allylic alkoxy/hydroxy groups could also exhibit stereodirecting effects in the Hg(II)-mediated intramolecular amidomercuration reactions.⁵ These prompted us to investigate “matching” and “mismatching” effects (double stereodifferentiation) by allylic *p*-methoxybenzyloxy (PMBO) group and *N*-trichloroacetyl group in the Hg(II)-mediated intramolecular amidomercuration reactions of 4-alkoxy-5-alkenyl amides **I** (PMBO at the C4 in eq 1).⁶ Furthermore, the amidomercuration products of **I** contain 2,6-disubstituted 3-piperidinol structures **II**, which have been found in many bioactive alkaloids such as julifloridine,⁷ cassine,⁸ Bao Gong Teng A,⁹ juliprosine,¹⁰ clavopictines,¹¹ and lepadines¹² (Figure 1). Therefore, the development of the highly diastereoselective amidomercuration reactions of **I** would be of particular use for the stereoselective/asymmetric synthesis of such alkaloids.¹³

The requisite optically pure 4-PMBO-5-alkenyl amides **I** for double stereodifferentiation studies could be accessed from the optically pure allylic alcohols **3** and **4**. As shown in Scheme 1, **3** and **4** were obtained via the lipase-catalyzed kinetic resolution of the readily available racemic allylic alcohol **1**¹⁴ with vinyl acetate as an acyl donor in THF.¹⁵ At 46% conversion, the *ee*'s of the acetate **2** and the remaining alcohol **3** were 97% and 83%, respectively. The optical purity of **3** was further improved to 99% (at 12% conversion) by subjecting **3** to the second round of the resolution. The transesterification reaction of **2** with MeOH in the presence of K₂CO₃ provided the optically pure allylic alcohol **4**.

Scheme 2 depicts how the optically pure allylic alcohols **3** and **4** were converted to the optically pure **I** (= the compounds **9** and **10**). The hydroxy group of **3** was protected by PMBCl and KH in DMF to give the PMB ether **5**. Acidic hydrolysis of **5** followed by the modified Horner-Wadsworth-Emmons olefination of the resulting aldehyde generated the α,β-unsaturated ester **6**.¹⁶ DIBAL reduction of **6** furnished the allylic alcohol **7**, which upon treatment with CCl₃CN and DBU in CH₂Cl₂ at 4 °C transformed into the trichloroacetimidate **8**.¹⁷ Enantioselective Overman rearrangement reaction of the imidate **8** by a chiral cobalt oxazoline palladacycle **III** [(*S*)-(+)-COP-Cl] in CH₂Cl₂ gave rise to the *N*-trichloroacetyl derivative **9** of the optically pure **I**.¹⁸ The *N*-trichloroacetyl derivative **10** of the optically pure **I**, where the stereochemistry of the PMBO group at C4 is opposite to that in **9**, was obtained by applying Scheme 2 to the compound **4**.

Double stereodifferentiation studies were conducted with **9** and **10**, and the results are given in Scheme 3. Compound **9** represented a matched case, since the 2,6-*trans*-directing effect of the *N*-trichloroacetyl group and the 2,3-*cis*-directing effect of the PMBO group were expected to reinforce each other to give the diastereomer **11** in a high diastereoselectivity.^{4,5} Indeed, the amidomercuration reaction of **9** with Hg(TFAO)₂ in the presence of K₂CO₃ (to prevent the

deprotection of the PMB group as well as the establishment of the equilibrating amidomercuration conditions by trifluoroacetic acid generated during the amidomercuration reaction) in nitromethane produced the 2,6-*trans* diastereomer **11** in >20:1 diastereoselectivity, and the trichloroacetyl group was deprotected under the conditions. When **9** was subjected to K₂CO₃ in nitromethane, the trichloroacetyl group was not deprotected even after a prolonged reaction time. Furthermore, the amidomercuration product of **9** (the compound **12**) was independently prepared by treating **11** with trichloroacetyl chloride and Et₃N in CH₂Cl₂. Upon exposure to the amidomercuration conditions, **12** was converted to **11** in almost quantitative reaction yield. These results indicated that the amidomercuration reaction took place on **9**, not on the amine resulting from the trichloroacetyl deprotection of **9**, and the deprotection of the trichloroacetyl group occurred after the amidomercuration to give **11**.

Surprisingly, the amidomercuration reaction of the mismatched diastereomer **10** also proceeded with >20:1 diastereoselectivity in favor of the 2,6-*trans* diastereomer **14**, despite the expectation that the *N*-trichloroacetyl group and the allylic PMBO group should oppose each other. These results indicate that the stereochemical outcome of the amidomercuration reactions is completely governed by the *N*-trichloroacetyl group, and the allylic PMBO group has little effect. In both cases, diastereoselectivities of the amidomercuration reactions were determined by measuring the relative integration of the diastereomeric methyl groups at C-2 after the reductive demercuration of the mercurial compounds **11** and **14** followed by the Cbz protection of the resulting amines to **13** and **15**, respectively.

For the confirmation of the stereochemistry of **13** and a synthetic utility of the developed reactions, compound **13** was elaborated to (+)-iso-6-cassine (Scheme 4). Cross metathesis reaction of the olefin **13** with dodec-11-en-2-one using the 2nd generation Grubbs' catalyst¹⁹ in refluxing CH₂Cl₂ afforded **16** as a mixture of *E* and *Z* isomers. Deprotection of the PMB group by CAN²⁰ followed by hydrogenation gave rise to (+)-iso-6-cassine (**17**), ¹H and ¹³C NMR spectra of which matched those in the literature.²¹

In summary, it has been shown that stereoselection in the Hg(II)-mediated intramolecular amidomercuration reactions of 5-alkenyl amides with a pedant allylic PMBO group is completely governed by *N*-TCA group, and the stereochemistry of the allylic PMBO group has little effect on the cyclization stereochemistry. The developed intramolecular amidomercuration reactions were used for the first asymmetric synthesis of (+)-iso-6-cassine. The origin and generality of the dominant 2,6-*trans* directing effect by the *N*-trichloroacetyl group in the intramolecular amidomercuration reactions are currently under investigation, and the results will be reported in due course

Acknowledgments

Financial support from the National Institutes of Health (GM 08194) and The Welch Foundation (AX-1534) is gratefully acknowledged.

References

1. Pelletier, SW., editor. Alkaloids: Chemical and Biological Perspectives. Vol. Vol. 3. Wiley; New York: 1985. Pelletier, SW., editor. Alkaloids: Chemical and Biological Perspectives. Vol. Vol. 10. Pergamon; Oxford, UK: 1996. (c) Michael JP. Nat. Prod. Rep 2001;18:520–542. [PubMed: 11699884]
2. For recent reviews: (a)Buffat MGP. Tetrahedron 2004;60:1701–1729.1729(b)Weintraub PM, Sabol JS, Kane JM, Borcharding DR. Tetrahedron 2003;59:2953–2989.2989
3. (a) Frederickson M, Grigg R. Org. Prep. Proc. Intl 1997;29:33–62. (b) Frederickson M, Grigg R. Org. Prep. Proc. Intl 1997;29:63–116. (c) Cardillo G, Orena M. Tetrahedron 1990;46:3321–3408.Harding, KE.; Tiner, TH. Comprehensive Organic Synthesis. Trost, B., editor. Pergamon Press; London: 1991.
4. Singh OV, Han H. Org. Lett 2004;6:3067–3070. [PubMed: 15330589]

5. (a)Takacs JM, Helle MA, Sanyal BJ, Eberspacher TA. *Tetrahedron Lett* 1990;31:6765–6768.6768(b) Martin OR, Xie F, Liu L. *Tetrahedron Lett* 1995;36:4027–4030.4030(c)Bernotas RC, Pezzone MA, Ganem B. *Carbohydrate Res* 1987;167:305–311.311(d)Takahata H, Banba Y, Tajima M, Momose T. *J. Org. Chem* 1991;56:240–245.245(e)Hugel HM, Hughes AB, Khalil K. *Aust. J. Chem* 1998;51:1149–1155.1155(f)Harding KE, Hollingsworth DR. *Tetrahedron Lett* 1988;29:3789–3792.3792(g)Chikkanna D, Han H. *Synlett* 2004:2311–2314.2314For a model to predict the stereochemistry of the allylic oxygen-directed cyclization reactions: (h)Chamberlin AR, Mulholland RL Jr. Kahn SD, Hehre WJ. *J. Am. Chem. Soc* 1987;109:672–677.677
6. Masamune S, Choy W, Petersen JS, Sita LR. *Angew. Chem. Int. Ed. Engl* 1985;24:1–30.
7. Recent selective syntheses of julifloridine: (a)Zhai H, Parvez M, Back TG. *J. Org. Chem* 2007;72:3853–3858.3858 [PubMed: 17439181](b)Lemire A, Charette AB. *Org. Lett* 2005;7:2747–2750.2750 [PubMed: 15957937](c)Kiguchi T, Shirakawa M, Honda R, Ninomiya I, Naito T. *Tetrahedron* 1998;54:15589–15606.15606
8. Recent selective syntheses of cassine: (a)Kim G, Kim N. *Tetrahedron Lett* 2007;48:4481–4483.4483 (b)Leverett CA, Cassidy MP, Padwa A. *J. Org. Chem* 2006;71:8591–8601.8601 [PubMed: 17064038] (c)Herdeis C, Kuepper P, Ple S. *Org. Biomol. Chem* 2006;4:524–529.529 [PubMed: 16446811](d) Tsai M-R, Chen B-F, Cheng C-C, Chang N-C. *J. Org. Chem* 2005;70:1780–1785.1785 [PubMed: 15730301](e)Makabe H, Kong LK, Hirota M. *Org. Lett* 2003;5:27–29.29 [PubMed: 12509882](f) Oetting J, Holzkamp J, Meyer HH, Pahl A. *Tetrahedron: Asymmetry* 1997;8:477–484.484(g)Toyooka N, Yoshida Y, Momose T. *Tetrahedron Lett* 1995;36:3715–3718.3718(h)Momose T, Toyooka N. *Tetrahedron Lett* 1993;34:5785–5786.5786
9. Recent selective syntheses of Bao Gong Teng A: (a)Zhang Y, Liebeskind LS. *J. Am. Chem. Soc* 2006;128:465–472.472 [PubMed: 16402833](b)Pham VC, Charlton JL. *J. Org. Chem* 1995;60:8051–8055.8055(c)Jung ME, Zeng L, Peng T, Zeng H, Le Y, Su J. *J. Org. Chem* 1992;57:3528–3530.3530
10. For a recent synthesis: Snider BB, Neubert BJ. *Org. Lett* 2005;7:2715–2718.2718 [PubMed: 15957929]
11. Recent selective syntheses of clavopictines: (a)Yu S, Pu X, Cheng T, Wang R, Ma D. *Org. Lett* 2006;8:3179–3182.3182 [PubMed: 16836360](b)Agami C, Couty F, Evano G, Darro F, Kiss R. *Eur. J. Org. Chem* 2003:2062–2070.2070(c)Ha JD, Cha JK. *J. Am. Chem. Soc* 1999;121:10012–10020.10020(d)Toyooka N, Yotsui Y, Yoshida Y, Momose T. *J. Org. Chem* 1996;61:4882–4883.4883For the isolation of clavopictines A and B: (e)Raub MF, Cardellina JH II, Choudhary MI, Ni CZ, Clardy J, Alley MC. *J. Am. Chem. Soc* 1991;113:3178–80.80
12. Recent selective syntheses of lepadines: (a)Pu X, Ma D. *J. Org. Chem* 2006;71:6562–6572.6572 [PubMed: 16901144](b)Pu X, Ma D. *Angew. Chem. Intl. Ed* 2004;43:4222–4225.4225(c)Kalai C, Tate E, Zard SZ. *Chem. Commun* 2002:1430–1431.1431(d)Ozawa T, Aoyagi S, Kibayashi C. *J. Org. Chem* 2001;66:3338–3347.3347 [PubMed: 11348115](e)Toyooka N, Okumura M, Takahata H. *J. Org. Chem* 1999;64:2182–2183.2183For the isolation of lepadines: (f)Wright AD, Goclik E, Konig GM, Kaminsky R. *J. Med. Chem* 2002;45:3067–3072.3072 [PubMed: 12086492](g)Davis RA, Carroll AR, Quinn RJ. *J. Nat. Prod* 2002;65:454–457.457 [PubMed: 11975478](h)Steffan B. *Tetrahedron* 1991;47:8729–8732.8732
13. Numata, A.; Ibuka, T. *The Alkaloids*. Brossi, A., editor. Vol. 31. Academic Press; New York: 1987. Strunz, GM.; Findlay, JA. *The Alkaloids*. Brossi, A., editor. Vol. 26. Academic Press; New York: 1985. p. 89-174.
14. Sasmal PK, Maier ME. *Org. Lett* 2002;4:1271–1274. [PubMed: 11950340]
15. (a) Ghanem A, Schurig V. *Tetrahedron: Asymmetry* 2003;14:57–62. (b) Langille NF, Panek JS. *Org. Lett* 2004;6:3203–3206. [PubMed: 15330623]
16. Blanchette MA, Choy W, Davis JT, Essensfeld AP, Masamune S, Roush WR, Sakai T. *Tetrahedron Lett* 1984;25:2183–2186.
17. Nishikawa T, Asai M, Ohyabu N, Isobe M. *J. Org. Chem* 1998;63:188–192. [PubMed: 11674062]
18. (a) Kirsch SF, Overman LE, Watson MP. *J. Org. Chem* 2004;69:8101–8104. [PubMed: 15527296] (b) Anderson CE, Overman LE. *J. Am. Chem. Soc* 2003;125:12412–12413. [PubMed: 14531676] (c) Overman LE, Owen CE, Pavan MM, Richards CJ. *Org. Lett* 2003;5:1809–1812. [PubMed: 12762658]
19. Scholl M, Ding S, Lee CW, Grubbs RH. *Org. Lett* 1999;1:953–956. [PubMed: 10823227]
20. Johansson R, Samuelsson B. *J. Chem. Soc. Perkin Trans I* 1984:2371–2374.

21. (a) Kamo T, Maehara K, Sato K, Hirota M. *Heterocycles* 2003;60:1303–1306. (b) Toyooka N, Yoshida Y, Yotsui Y, Momose T. *J. Org. Chem* 1999;64:4914–4919. [PubMed: 11674570] (c) Christofidis I, Welter A, Jadot J. *Tetrahedron* 1977;33:977–979.

Figure 1.
Various 2,6-disubstituted 3-piperidinol alkaloids

Scheme 1.
Synthesis of optically pure allylic alcohols.

Scheme 2.
Synthesis of substrates for the amidomercuration.

Scheme 3.

Double stereodifferentiation in the amidomercuration.

Scheme 4.
Asymmetric synthesis of (+)-iso-6-cassine