Correspondence

Clinical significance of postoperative CA-125 level after primary cytoreductive surgery in ovarian cancer

To the editor: The current study is meaningful in that the decline in serum CA-125 levels after primary cytoreductive surgery may affect progression-free survival (PFS) in patients with epithelial ovarian cancer (EOC). Moreover, similar studies have also shown that the serum CA-125 level may be a prognostic factor for the prediction of survival in patients with EOC although the time to evaluate serum CA-125 levels was different.¹⁻³

However, some questions should be resolved for understanding the role of serum CA-125 levels in EOC. Firstly, histological type of EOC may affect serum CA-125 levels. Especially, serum CA-125 levels are known to be lower in clear cell or mucinous carcinoma than in serous adenocarcinoma.⁴ Thus, a higher rate of non-serous type ovarian cancer may affect serum CA-125 levels and interrupt the reflection of tumor burden by serum CA-125 levels. Thus, the rate of non-serous type ovarian cancer might affect no correlation between postoperative decline in serum CA-125 levels and residual disease. Secondly, various types of chemotherapeutic regimen may affect response and survival in patients with EOC. Although all patients received platinum-based chemotherapy in the current study, the authors have not shown how many patients received taxane- and platinum-based chemotherapy. In some studies of Gynecologic Oncology Group (GOG), taxane- and platinum-based chemotherapy has been reported to be the most efficient for the treatment of EOC.^{5,6} Thus, if there were different chemotherapeutic regimens in the current study, PFS might be influenced by type of chemotherapeutic regimen as well as serum CA-125 levels. Thirdly, retrospective design may affect the results by some bias.

Conclusively, serum CA-125 levels may be helpful for the prediction of clinical outcomes in patients with EOC who received staging laparotomy followed by adjuvant chemotherapy. However, prospective clinical trials by controlling some bias such as various histological types and different types of chemotherapeutic regimen are needed for evaluating the definite role of serum CA-125 levels.

REFERENCES

1. Juretzka MM, Barakat RR, Chi DS, Iasonos A, Dupont J, Aburustum NR, et al. CA125 level as a predictor of progression-free

- survival and overall survival in ovarian cancer patients with surgically defined disease status prior to the initiation of intraperitoneal consolidation therapy. *Gynecol Oncol* 2007; 104: 176-80.
2. Crawford SM, Peace J. Does the nadir CA125 concentration predict a long-term outcome after chemotherapy for carcinoma of the ovary? *Ann Oncol* 2005; 16: 47-50.
3. Kim HS, Park NH, Chung HH, Kim JW, Song YS, Kang SB. Serum CA-125 level after 6 cycles of primary adjuvant chemotherapy is a useful prognostic factor for complete responders' survival in patients with advanced epithelial ovarian cancer. *Onkologie* 2008; 31: 315-20.
4. Eltabbakh GH, Mount SL, Beatty B, Simmons-Arnold L, Cooper K. Clinical and molecular differences between clear cell and papillary serous ovarian carcinoma. *J Surg Oncol* 2006; 93: 379-86.
5. Muggia FM, Braly PS, Brady MF, Sutton G, Niemann TH, Lentz SL, et al. Phase III randomized study of cisplatin versus paclitaxel versus cisplatin and paclitaxel in patients with suboptimal stage III or IV ovarian cancer: A gynecologic oncology group study. *J Clin Oncol* 2000; 18: 106-15.
6. Piccart MJ, Bertelsen K, James K, Cassidy J, Mangioni C, Simonsen E, et al. Randomized intergroup trial of cisplatin-paclitaxel versus cisplatin-cyclophosphamide in women with advanced epithelial ovarian cancer: Three-year results. *J Natl Cancer Inst* 2000; 92: 699-708.

Hee Seung Kim, Noh Hyun Park

Department of Obstetrics and Gynecology, College of Medicine, Seoul National University, Seoul, Korea

Address correspondence and reprint requests to: **Noh Hyun Park**

Department of Obstetrics and Gynecology, College of Medicine, Seoul National University, 28, Yeongeon-dong, Jongno-gu, Seoul 110-744, Korea

Tel: 82-2-2072-3166, Fax: 82-2-762-3599

E-mail: pnkhkr@snu.ac.kr

In reply: We are grateful to Dr. Kim for their interest in our article. In our study, we showed that postoperative decline of CA-125 may be associated with progression free survival in patients with advanced stage ovarian cancer.

First, Dr. Kim said that non-serous malignant tumors should be excluded from the study group. But practically speaking, there is no better tumor marker for malignant mucinous tumors other than CA-125, and indeed there are many other articles in the literature that have included CA-125 measurements in patients with malignant tumors other than serous types including mucinous ones.¹⁻³ Second, we agree that platinum based chemotherapy without paclitaxel is an acceptable choice of treatment but may be not the best regimen. In our study, the number of patients who received chemotherapy without paclitaxel was 17 among 42 patients in the <75% decline group, and it was 14 among 39 of the ≥75% group, and there was no statistical difference ($p=0.819$).

There is substantial literature concerning CA-125 levels and ovarian cancer. Unfortunately, there have been no prospective

trials assessing the relationship between survival and post-operative CA-125. Well-designed prospective trials on more specific patients would yield more worthy outcome as we already mentioned in our study.

REFERENCES

1. Markman M, Liu PY, Rothenberg ML, Monk BJ, Brady M, Alberts DS. Pretreatment CA-125 and risk of relapse in advanced ovarian cancer. *J Clin Oncol* 2006; 24: 1454-8.
2. Obeidat B, Latimer J, Crawford R. Can optimal primary cytoreduction be predicted in advanced stage epithelial ovarian cancer? Role of preoperative serum CA-125 level. *Gynecol Obstet Invest* 2004; 57: 153-6.
3. Menczer J, Chetrit A, Sadetzki S, Golan A, Levy T. Follow-up of

ovarian and primary peritoneal carcinoma: The value of physical examination in patients with pretreatment elevated CA125 levels. *Gynecol Oncol* 2006; 103: 137-40.

Seung-Chul Yoo, Suk-Joon Chang

*Department of Obstetrics and Gynecology,
Ajou University School of Medicine, Suwon, Korea*

Address correspondence and reprint requests to: **Suk-Joon Chang**
Department of Obstetrics and Gynecology, Ajou University School of Medicine, San 5, Woncheon-dong, Yeongtong-gu, Suwon 443-749, Korea
Tel: 82-31-219-5251, Fax: 82-31-219-5245
E-mail: drchang@ajou.ac.kr