Published in final edited form as:

ACS Chem Biol. 2009 March 20; 4(3): 209–220. doi:10.1021/cb800304d.

The Molecular Basis of Species-Specific Ligand Activation of Trace Amine-Associated Receptor 1 (TAAR₁)

Edwin S. Tan^{a,e}, John C. Naylor^d, Eli S. Groban^b, James R. Bunzow^d, Matthew P. Jacobson^c, David K. Grandy^d, and Thomas S. Scanlan^{c,d,f}

^aChemistry and Chemical Biology Graduate Program, University of California, San Francisco, San Francisco, CA 94158-2280

^bGraduate Group in Biophysics, University of California, San Francisco, San Francisco CA 94158-2240

^cDepartment of Pharmaceutical Chemistry, University of California, San Francisco, San Francisco, CA 94158, USA

^dDepartment of Physiology and Pharmacology, Oregon Health and Science University, Portland, OR 97239, USA

Abstract

The trace amine-associated receptor **1** (TAAR₁) is an aminergic G protein-coupled receptor (GPCR) potentially activated by 3-iodothyronamine (**1**), an endogenous derivative of thyroid hormone. Structure activity relationship studies on **1** and related agonists showed that the rat and mouse species of TAAR₁ accommodated structural modifications and functional groups on the ethylamine portion and the biaryl ether moiety of the molecule. However, the two receptors clearly exhibited distinct, species-specific ligand preferences despite being remarkably similar with 93% sequence similarity. In this study, we generated single and double mutants of rat and mouse TAAR₁ to probe the molecular recognition of agonists and the underlying basis for the ligand selectivity of rat and mouse TAAR₁. Key, non-conserved specificity determinant residues in transmembranes helices 4 and 7 within the ligand binding site appear to be the primary source of a number of the observed ligand preferences. Residue 7.39 in transmembrane 7 dictated the preference for a β -phenyl ring while residue 4.56 in transmembrane 4 was partially responsible for the lower potency of **1** and **tyramine** for the mouse receptor. Additionally, **1** and **tyramine** were found to have the same binding mode in rat TAAR₁ despite structure activity relationship data suggesting the possibility of each molecule having different binding orientations. These findings provide valuable insights into the critical binding site residues involved in the ligand-receptor interaction that can influence compound selectivity and functional activity of aminergic GPCRs.

INTRODUCTION

3-Iodothyronamine (**1**, T₁AM; Fig. 1) is an endogenous derivative of the thyroid hormone thyroxine (T₄; Fig. 1) detected in the brain, heart, liver, and blood (1,2). It has profound pharmacological effects *in vivo* that are mostly opposite to those of T₄ (3). Compound **1** induces anergia, hypothermia, bradycardia, hyperglycemia, and hyperinsulinemia when administered to mice and rapidly reduces cardiac output in an *ex vivo* working rat heart preparation (2,4–

^fCorrespondence should be addressed to T.S.S. (E-mail: scanlant@ohsu.edu; phone (503) 494-9292; fax (503) 494-9275).

^ePresent Address: Department of Systems Biology, Harvard Medical School, 200 Longwood Avenue, WA 536, Boston, MA 02115, USA

Supporting Information Available: This material is available free of charge via the Internet.

8). Additionally, **1** can increase food intake and influence energy metabolism by favoring lipid utilization over carbohydrate consumption (9,10).

These physiological responses to **1** may be mediated by more than one molecular target. *In vitro*, **1** can activate aminergic G protein-protein coupled receptors (GPCRs) in the biogenic amine subfamily; stimulating the production of cAMP (adenosine 3',5'-cyclic monophosphate) via the trace amine-associated receptor 1 (TAAR₁) and the degradation of cAMP via the α_{2A} adrenergic receptor (2,7,11,12). Additionally, **1** has a neuromodulatory role, inhibiting neurotransmitter reuptake by the dopamine (DAT) and norepinephrine transporter (NET), and inhibiting vesicular packaging by the vesicular monoamine transporter 2 (VMAT2) (13).

To date, **1** is the most potent endogenous molecule that can activate rat and mouse TAAR₁. TAAR₁ is a member of the trace amine-associated receptor family of orphan GPCRs, which consists of 19 rat, 16 mouse, and 9 human subtypes (14–18). It is homologous to the dopamine, adrenergic, and serotonin receptors and expressed in multiple tissues including the heart, kidney, liver, spleen, and pancreas. Although a biological function for any TAAR subtype has yet to be defined, rodent and human species of TAAR₁ are potently activated by endogenous biogenic amines such as β -phenethylamine as well as the psychostimulants amphetamine and methamphetamine in a species-dependent manner (12,14,15,19–21). In addition, other members of the rodent TAAR family were recently implicated as a secondary class of chemosensory receptors expressed in the olfactory epithelium (22).

In an effort to determine the role of TAAR₁ in mediating the effects of **1**, we have previously explored the structure activity relationship (SAR) of **1** and developed a number of analogs with improved agonist activity (11,23). These SAR studies showed rat and mouse TAAR₁ (rTAAR₁ and mTAAR₁, respectively) could tolerate structural modifications and substituents on the ethylamine portion and biaryl ether moiety of **1**. However the two receptors clearly have distinct structural preferences. In the ethylamine chain, for example, rTAAR₁ favored unsaturated hydrocarbon substituents while mTAAR₁ preferred polar groups and hydrogen bond acceptors (23). Compound **1** and **tyramine** (Table 1), an endogenous phenethylamine metabolite, also exhibited some degree of species variability, being ~7 to 10-fold less potent for mTAAR₁ compared to rTAAR₁ (11). These distinct ligand selectivities of rat and mouse TAAR₁ are unusual given that the two receptors are 93% similar in primary sequence. Since the rodent receptors are only 83%–85% similar to human TAAR₁, understanding the molecular basis of species variability will undoubtedly have important implications in the development of activators and inhibitors for human TAAR₁. In this study, we explored the molecular recognition of **1** and related agonists by TAAR₁ and identified specificity determining residues that give rise to the disparate ligand preferences between rat and mouse TAAR₁.

RESULTS AND DISCUSSION

Our initial structure activity relationship study identified the β -phenylphenoxy-phenethylamine (**2** & **3**) and phenoxy-naphthethylamine (**4**) as promising scaffolds for the development of small molecule regulators of TAAR₁ (Fig. 1) (23). Compounds **2** and **3** were both rTAAR₁ selective, being 357- to 526-fold more potent for rTAAR₁ compared to mTAAR₁ (Table 1). Compound **4**, on the other hand, was less discriminating between rat and mouse TAAR₁ with a disparity in potency of 5-fold between the two receptors. Using the toggle switch model of aminergic GPCR activation as a guideline, **2** was further developed to give superagonists (agonists that are more potent and/or efficacious than **1**) **5**, **6**, and **7** (Table 1) (24). These superagonists remained rTAAR₁ selective exhibiting potencies that were $\geq$ 1600- to 2500-fold better at rTAAR₁ versus mTAAR₁.

The poor agonist activity of the β -phenylphenoxyphenethylamines at mTAAR₁ can be attributed to the β -phenyl group of the molecule and not its outer ring moiety (Fig. 1). Removing the outer rings of **2** and **5** (**8** and **9**, respectively) did not increase potency at mTAAR₁ (Table 1). Both **8** and **9** still activated mTAAR₁ poorly with a potency $\geq 10\mu\text{M}$. In contrast, eliminating the β -phenyl ring of **2**, **5**, and **7** to give **1** or **10** improved the potency at mTAAR₁ ~24- to 32-fold and decreased the rTAAR₁ selectivity from ≥ 350 - to 2500-fold down to ~7–10-fold.

Aminergic GPCRs are heptahelical transmembrane proteins with an extracellular amino terminus and an intracellular carboxy terminus (Fig. 2a). The ligand binding site of aminergic GPCRs is located within the transmembrane (TM) region of the receptor and is predominantly composed of residues from TM 3, 5, 6 and 7 (25–28). Based on pharmacological and mutagenesis studies, epinephrine is proposed to bind to the β_2 -adrenergic receptor ($\beta_2\text{AR}$) with aspartic acid 3.32 (D3.32) acting as the counterion for the charged amine, serine residues 5.42, 5.43, and 5.46 (S5.42, S5.43, and S5.46, respectively) interacting with the catechol hydroxyls, phenylalanines 6.51 and 6.52 (F6.51 and F6.52) interacting with the catechol ring, and asparagines 6.55 (N6.55) as the partner for the β -hydroxy group (Fig. 2b) (see experimental procedures for a description of the residue indexing system) (29–36). In the recently solved crystal structure of an engineered human $\beta_2\text{AR}$, all of these residues were found to line the ligand binding site and interact with the inverse agonist that co-crystallized with the receptor (26,27). By analogy to the catecholamine receptors (dopamine, epinephrine, and norepinephrine) we have previously deduced **2** to bind to rTAAR₁ with the charged amine forming a salt bridge interaction with D3.32 and the biaryl ether oxygen hydrogen bonding to S5.46 (Fig. 2c) (24). In this binding orientation, our homology model of rTAAR₁ showed the β -phenyl ring to be positioned near the interface between TM 6 and 7 and surrounded by cysteine 6.54, methionine 6.55, and asparagines 7.35 and 7.39 (C6.54, M6.55, N7.35, and N7.39, respectively).

In mTAAR₁, C6.54 and N7.35 are conserved but not M6.55 and N7.39. Instead mTAAR₁ has a threonine and tyrosine at 6.55 and 7.39, respectively (T6.55 and Y7.39) (Fig. 2d). In the $\beta_2\text{AR}$, the residues at 6.55 and 7.39 have been previously shown to interact with the ligand and alter the receptor's ligand specificity (35,37). Since the β -phenyl ring is proposed to be in the vicinity of 6.55 and 7.39, and both residues are not conserved between rat and mouse TAAR₁, we hypothesized that one or both of these residues are specificity elements that influence compatibility with the β -phenyl ring of the β -phenylphenoxyphenethylamine scaffold. We tested this hypothesis by measuring the activity of **3** and **5**, representative β -phenylphenoxyphenethylamines, against rat and mouse TAAR₁ in HEK293 (human embryonic kidney 293) cells stably and heterologously expressing single or double swap mutants at 6.55 and/or 7.39.

Residue 7.39 controls specificity for the β -phenyl ring of TAAR₁ ligands

Swapping residue 6.55 of rat and mouse TAAR₁ had minor effects on the activity and selectivity of **3** and **5**. In the rTAAR₁ 6.55 single mutant [rTAAR₁(M6.55T)] the potency and efficacy of **3** ($\text{EC}_{50} = 55 \pm 20 \text{ nM}$, $E_{\text{max}} = 119 \pm 6\%$) and **5** ($\text{EC}_{50} = 11 \pm 1 \text{ nM}$, $E_{\text{max}} = 117 \pm 20\%$) decreased 2- to 3-fold and $\leq 12\%$, respectively (Table 2). When T6.55 was mutated to a methionine in mTAAR₁ [mTAAR₁(T6.55M)], both compounds were still poor agonists activating the receptor with potencies $>10\mu\text{M}$ and efficacies ranging from 20% to 45%. Despite the mutation at residue 6.55, both **3** and **5** remained considerably rTAAR₁ selective (182- to 909-fold) (Fig. 3).

The single swap mutants at residue 7.39 had opposing effects on the activity of **3** and **5**. The potency of both compounds decreased 167- to 526-fold in the rTAAR₁ 7.39 mutant [rTAAR₁(N7.39Y)] ($\text{EC}_{50} = \sim 10 \mu\text{M}$ and $\sim 1 \mu\text{M}$ for **3** and **5**, respectively) but increased 60- to 100-fold in its mTAAR₁ 7.39 mutant counterpart [mTAAR₁(Y7.39N)] ($\text{EC}_{50} = 102 \pm 21$

nM and 176 ± 32 nM for **3** and **5**, respectively) (Table 2). The efficacy of **3** ($E_{\max} = 95 \pm 6\%$) and **5** ($E_{\max} = 91 \pm 11\%$) both increased 29% to 89% in mTAAR₁(Y7.39N). For rTAAR₁(N7.39Y) the efficacy of **5** ($E_{\max} = 126\%$) increased 12% while that of **3** ($E_{\max} = 85\%$) decreased 46% compared to wild type. Interestingly, swapping the residues at 7.39 converted both compounds to good mTAAR₁ agonists that were now 6- to 98-fold selective for mTAAR₁ over rTAAR₁ (Fig. 3).

The activity profile of **3** and **5** for the double swap mutants was similar to that of the 7.39 single mutants but showed some enhancements in both potency and efficacy. The potency of **3** and **5** decreased 333- to 526-fold for the rTAAR₁ 6.55 and 7.39 double mutant [rTAAR₁(M6.55T/N7.39Y)] but increased 70- to 320-fold for the mTAAR₁ 6.55 and 7.39 double mutant equivalent [mTAAR₁(T6.55M/Y7.39N)] (Table 2). The same trend was also observed with regards to efficacy; decreasing 44% to 85% for rTAAR₁(M6.55T/N7.39Y) and increasing 39% to 95% for mTAAR₁(T6.55M/Y7.39N). Compared to the 7.39 single mutant, the mTAAR₁ selectivity of **3** and **5** in the double mutant was more pronounced (Fig. 3).

The decrease in activity of **3** and **5** for the 6.55 and/or 7.39 single and double mutants in rTAAR₁ cannot be attributed to the introduced mutations compromising the functional competency of the receptors because the activity of the positive controls (**1** and **4**) for the same mutants only changed ≤ 4 -fold and $\leq 16\%$ in terms of potency and efficacy, respectively (Table 2). Likewise, the enhanced activity of **3** and **5** for the mTAAR₁ single and double swap mutants is not a consequence of the mutations rendering the receptors constitutively active and more responsive to agonists because the potency of the positive controls (**1**, **4**, and/or **11**) only changed ≤ 2 -fold and the efficacy were essentially identical compared to the wild type receptor (Table 1 and 2). Compound **11** is a novel agonist for rat and mouse TAAR₁ that can be considered a halogen free analog of **1**.

Swapping residues at 7.39 was sufficient to convert **3** and **5** from a rTAAR₁ into a mTAAR₁ selective agonist. The TAAR₁ binding site appears to be able to accommodate a phenyl ring in the interface between TM 6 and 7 within the binding site near residue 7.39. Compounds **3** and **5** are poor agonists for wild type mTAAR₁ because this phenyl pocket near 7.39 is occupied by the phenol group of Y7.39 from the receptor and unavailable to the β -phenyl rings of **3** and **5**. In contrast, **3** and **5** are excellent agonists for wild type rTAAR₁ because the smaller asparagine residue at 7.39 is less sterically encumbering and does not compete with the β -phenyl rings of **3** and **5** for the phenyl pocket near 7.39 of the receptor.

It should be noted that the substitution pattern of the outer ring does not significantly affect the specificity of TAAR₁ for β -phenyl ring bearing compounds. Regardless of whether the outer ring is at the *meta*- (**3**) or *para*- (**5**) position relative to the ethylamine chain, both **3** and **5** were affected similarly by mutations at 6.55 and/or 7.39 in rat and mouse TAAR₁, indicating that the β -phenyl ring of both molecules occupy the same binding pocket within the binding site. This result supports the assumptions we had proposed regarding the location of the antagonistic groups of the lead rTAAR₁ antagonists previously developed with **3** as the core scaffold (24).

Based on its primary sequence, we predict human TAAR₁ (hTAAR₁) to be able to accommodate a phenyl ring at the β carbon of phenethylamine based ligands because the residue at 7.39 in the human receptor is an isoleucine (Fig. 2e). Since isoleucine is smaller than tyrosine and approximately the same size as asparagine, the phenyl pocket near 7.39 should also be present in hTAAR₁.

Compound **1** and Tyramine have similar binding modes

In addition to being a superagonist for rTAAR₁, **7** ($EC_{50} = 4 \pm 1$ nM, $E_{\max} = 115 \pm 2\%$) is also an interesting molecule because it contains the structures of both **tyramine** ($EC_{50} = 65 \pm 1$

nM, $E_{\max} = 119 \pm 7\%$) and **1** ($EC_{50} = 33 \pm 3$ nM, $E_{\max} = 100 \pm 0\%$) in the same compound (Table 1 and Fig. 4). This hybrid compound potentially explains how two molecules with very different molecular volumes can elicit similar responses. If the β -phenyl group of **7** represents the aromatic ring of **tyramine**, then **tyramine** would occupy the phenyl pocket near 7.39 and thus have a binding mode different from **1** (Fig. 4). On the other hand, if the inner ring of **7** represents the **tyramine** aromatic ring then **tyramine** and **1** would have similar binding modes. The rTAAR₁ agonist activity of **9** ($EC_{50} = 115 \pm 12$ nM, $E_{\max} = 105 \pm 5\%$) supports the feasibility of **tyramine** potentially having two alternate binding modes in the rat receptor (Table 1). Since the tyrosine residue at 7.39 in mTAAR₁ abolished the phenyl pocket near 7.39, **tyramine** must share the same binding mode as **1** in mTAAR₁.

To determine if **tyramine** and **1** have similar or distinct binding modes, we mutated alanine 5.42 in rTAAR₁ to a threonine, leucine, or isoleucine [rTAAR₁(A5.42T), rTAAR₁(A5.42L), and rTAAR₁(A5.42I), respectively] and examined the effects of the mutations on the potency of **tyramine** and **1**. The idea behind these 5.42 single mutants is to perturb the pocket occupied by the outer ring of **1**, **7** and other biaryl ether containing molecules by introducing larger residues than alanine. Threonine was chosen because SAR studies on human TAAR₁, which has a threonine at residue 5.42 (Fig. 2e), showed **1** to be a less potent agonist against hTAAR₁ than rTAAR₁ (**12**).

In the rTAAR₁(A5.42L) mutant, the potency of **1** ($EC_{50} = 58 \pm 16$ nM) and **tyramine** ($EC_{50} = \sim 2$ μ M) decreased 2- and 31- fold, respectively (Table 3a). When A5.42 was mutated to isoleucine, there was a 3- and 154-fold decrease in the potency of **1** ($EC_{50} = 108 \pm 14$ nM) and **tyramine** ($EC_{50} = >10$ μ M), respectively. The efficacy of **tyramine** in the leucine mutant decreased 53% to 70% compared to wild type rTAAR₁. A similar trend was observed for the rTAAR₁(A5.42T) mutant where **1** ($EC_{50} = 88 \pm 13$ nM) and **tyramine** ($EC_{50} = >10$ μ M) were 3- and 154-fold less potent. Additionally, **tyramine** was 50% less efficacious for this receptor than wild type. Since the activity of the positive control (**11**) for all 5.42 mutants were essentially unaffected, the reduced activities of **1** and **tyramine** for these mutants cannot be due to a compromised activation capacity of these receptors.

The observed effects on the activity of **1** and **tyramine** in the rTAAR₁ 5.42 mutants are consistent with both compounds having similar binding modes. If **tyramine** preferentially occupied the phenyl pocket near residue 7.39, its potency would not have been affected by changes to residue 5.42. Since its activity decreased 31- to 154-fold when residue 5.42 was mutated, **tyramine** is probably binding to rTAAR₁ with its hydroxyl group engaged in hydrogen bond interactions with S5.46, the residue one turn below 5.42 in TM5 (Fig. 2d). This binding mode corresponds to the same binding orientation of **1** for rat and mouse TAAR₁.

Although the leucine and isoleucine mutations increased the steric bulk around residue 5.42, it is interesting that the agonist activity of the smaller **tyramine** was more severely affected than the larger **1** or **11**; especially when these mutations were intended to block the binding pocket for the outer ring. These results suggest that despite being bigger than alanine, leucine and isoleucine are not large enough to completely abolish the outer ring binding pocket or are sufficiently flexible to accommodate the outer phenyl ring. The small effects of the mutations on the agonist activity of **1** and **11** can potentially be attributed to their larger number of interactions with the receptor compared to **tyramine**. Like **tyramine**, **1** and **11** should both be anchored in the binding site of rTAAR₁ by a salt bridge interaction between the charged amine and D3.32, and a hydrogen bond interaction with S5.46. However, the extra functional groups present in **1** and **11** (i.e. outer ring, iodine, and naphthyl ring) (Table 1) can make additional interactions with TM 5 and 6 that are not available to **tyramine**. With more contacts to the receptor, **1** and **11** would be less sensitive to structural changes at residue 5.42 than **tyramine**.

Residue 4.56 is partially responsible for the lower potency of **1** for mTAAR₁

Within the thyronamine series, rat and mouse TAAR₁ had the same rank order potency but the potency values of individual compounds for the two receptors were not identical. In general, thyronamines are ~10-fold less potent for mTAAR₁ compared to rTAAR₁ (2,11). Assuming all ligands target the same binding pocket and lead to the same active receptor conformation, a possible explanation for this potency disparity can be attributed to a lower G-protein coupling efficiency for mTAAR₁ versus rTAAR₁. If this were the case, then it would be impossible to have an equipotent agonist for both receptors because mTAAR₁ would be inherently less sensitive to ligand activation than rTAAR₁. Since **12** was found to be an equipotent agonist for rTAAR₁ ($EC_{50} = 65 \pm 6$ nM, $E_{max} = 115 \pm 2\%$) and mTAAR₁ ($EC_{50} = 82 \pm 17$ nM, $E_{max} = 112 \pm 3\%$), the G-protein coupling efficiency of mTAAR₁ is comparable to that of rTAAR₁ (Table 1).

We hypothesized that the potency disparity of thyronamines was brought about by nonconserved amino acid(s) at key specificity determinant residues within the binding site. In particular, we speculated that tyrosine 4.56 (Y4.56) was primarily responsible for the ~10-fold lower potency of **1** for mTAAR₁. This residue was identified through a process of elimination based on the following five points: (1) since the binding sites for most aminergic GPCRs are located within the transmembrane regions of the receptor, all intracellular and extracellular loops as well as the amino- and carboxy-terminus were eliminated (28); (2) amino acid differences in TM 1 and 2 were eliminated because the binding site is primarily composed to TM 3, 4, 5, 6, and 7 (30); (3) since the ethylamine chain of **1** and **12** are exactly the same, non-conserved residues in TM 3, 6 and 7 cannot be responsible; (4) TM5 was eliminated because it is absolutely conserved between the two species; (5) the intracellular half of TM 4 was eliminated because the binding site of GPCRs is located in the extracellular half of the transmembrane region (Fig. 5). The only non-conserved residue remaining was Y4.56. In our homology models of rat and mouse TAAR₁, residue 4.56 was found to be in the vicinity of the purported outer ring binding pocket and could conceivably make contacts with the bound ligand. To test the importance of residue 4.56, we generated rat and mouse TAAR₁ single swap mutants at this location.

When Y4.56 of mTAAR₁ was converted to phenylalanine [mTAAR₁(Y4.56F)], the potency of **1** increased 5-fold from 314 ± 43 nM to 67 ± 17 nM (Table 1 and **3b**). Interestingly, the potency of **1** ($EC_{50} = 38 \pm 11$ nM) for the tyrosine mutant of rTAAR₁ [rTAAR₁(F4.56Y)] was comparable to wild type rTAAR₁ (33 ± 3 nM). The same trend was also observed for **tyramine**; where its potency increased 7-fold in mTAAR₁(Y4.56F) ($EC_{50} = 40 \pm 13$ nM) but remained unaffected in rTAAR₁(F4.56Y) ($EC_{50} = 54 \pm 11$ nM). Since the agonist activity of the positive controls (**4** and **11**) for the mutants was comparable to that of the wild type receptors for both species, the mutations did not compromise the functional capacity of either receptors.

Residue 4.56 appears to play an important role in the lower potency of **1** for mTAAR₁. Swapping this residue with the corresponding residue found in rTAAR₁ increased the potency of **1** in mTAAR₁ to within two-fold of the potency value for wild type rTAAR₁. Interestingly, the potency of **tyramine** for mTAAR₁(Y4.56F) was equivalent to that of wild type rTAAR₁. Although mutating residue 4.56 improved the potency in the mTAAR₁ mutants, the reciprocal effect of **1** and **tyramine** becoming less potent for rTAAR₁(F4.56Y) was not observed. This indicates that 4.56 is only partially responsible and not the sole basis for the observed potency disparity of **1** and **tyramine** between the rat and mouse TAAR₁. Overall, these results implicate residue 4.56 in TM4 to affect ligand interaction and receptor activation.

CONCLUSION AND SIGNIFICANCE

The disparate ligand structural preferences exhibited by rat and mouse TAAR₁ can be attributed to key, non-conserved specificity determining residues within the binding site. Residue 7.39 appears to dictate the specificity for a β -phenyl ring; the results suggest that the bulky tyrosine residue at 7.39 in mTAAR₁ sterically clashed with the β -phenyl ring whereas the smaller asparagine at the same location in rTAAR₁ was more compatible and able to accommodate a β -phenyl moiety. The lower potency of **1** in mTAAR₁ was partly caused by the presence of a tyrosine at residue 4.56 rather than a phenylalanine. Although compound **7** implied the possibility of **1** and **tyramine** having different binding modes in the binding site of rTAAR₁, **1** and **tyramine** appear to have the same binding orientation.

With the recent developments and accomplishments in structure determination of the β_2 adrenergic receptor, the practicality of a structure-based drug design approach towards developing activators and inhibitors for aminergic GPCRs has never been so promising. A critical aspect to the success of this strategy will depend on having insights into the molecular basis of ligand recognition, the mechanism of GPCR activation, and the relationship of how these ligand-receptor interactions are relayed and translated into receptor activation or inhibition. The information presented herein should prove beneficial towards this cause as it provides valuable information regarding the binding site residues involved in ligand-receptor interactions that can influence compound specificity and functional activity of an aminergic GPCR.

METHODS

Residue Indexing Scheme

Residues are labeled relative to the most conserved amino acid in the transmembrane segment in which it is located (38). Asparagine 7.39, for example, is located in transmembrane 7 and precedes the most conserved residue by 11 positions. Proline 6.50 is the most conserved residue in TM6. This system simplifies the identification of corresponding residues in different GPCRs.

TAAR₁ Site-Directed Mutagenesis

TAAR₁ mutants were generated by using the QuikChange Site-Directed Mutagenesis Kit (Stratagene, La Jolla, CA). Primers were designed that coded for the desired mutation flanked with 10–15 base pairs of sequence. Complementary oligonucleotides were then used in PCR using an expression plasmid containing the desired receptor as template. The PCR product was digested with Dpn I and transformed into XL1 Blue competent cells. Colonies were picked and the DNA isolated was sequenced to confirm the mutation.

The DNA for the mutants was then used to transfect HEK293 cells using Fugene (Roche, Indianapolis, IN) and stable cell lines were made for further assays under G418 selection.

Homology Model of rat and mouse TAAR₁

The homology model for rTAAR₁ was created using the Prime software package (Schrödinger Inc.). The model was based on a sequence alignment of rTAAR₁ with the sequence of human β_2 AR, for which a crystal structure was recently solved (Protein Data Bank accession code 2RH1), created using the “align GPCR” module (26,27). The modeling procedure involves side chain rotamer optimization and closure of chain breaks due to gaps in the sequence alignment using a previously published loop building and optimization algorithm (39). After building the complete model, additional side chain rotamer optimization, followed by backbone and side chain energy minimization, was performed on all non-conserved residues. The homology modeling program relies on the OPLS all atom force field (40–42) and a Generalized

Born solvent model (43,44) to evaluate the energy of different conformations and select the lowest energy structure as the final model.

Synthesis

Detailed synthetic procedures and characterization information for novel compounds **8**, **9**, **11**, and **12** are described in the supporting information.

In Vitro cAMP Agonist Activity Assay

Compounds were tested using the Hithunter cAMP XS kit (DiscoverX) as described previously (24). Data were reported relative to **1** and expressed as %T₁AM. The activity of **1** at 10μM was set as 100 %T₁AM. Concentration-response curves were plotted and EC₅₀ values were calculated with Prism software (GraphPad, San Diego, CA). Standard error of the mean was calculated from the EC₅₀ and E_{max} values of each independent triplicate experiment by use of Prism Software.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgements

This work was supported by grants from the National Institutes of Health (Grant DK52798 to T.S.S.), the Portland Methamphetamine Abuse Research Center (D.K.G.), the OHSU President's Fund (D.K.G.), the Alfred P. Sloan Foundation (research fellowship to M.P.J.), and Ikaria, Inc. (T.S.S.). M.P.J. is a member of the Scientific Advisory Board of Schrödinger, Incorporated. The authors would also like to acknowledge thoughtful discussions with Dr. Edmund A. Reese.

References

1. Piehl S, Heberer T, Balizs G, Scanlan TS, Smits R, Koksche B, Kohrle J. Thyronamines are isozyme-specific substrates of deiodinases. *Endocrinology* 2008;149:3037–45. [PubMed: 18339710]
2. Scanlan T, Suchland K, Hart M, Chiellini G, Huang Y, Kruzich P, Frascarelli S, Crossley D, Bunzow J, Ronca-Testoni S, Lin E, Hatton D, Zucchi R, Grandy D. 3-Iodothyronamine is an endogenous and rapid-acting derivative of thyroid hormone. *Nat Med* 2004;10:638–642. [PubMed: 15146179]
3. Scanlan TS. 3-Iodothyronamine (T1AM): A new player on the thyroid endocrine team? . *Endocrinology*. 2008 Dec 30;[Epub ahead of print]
4. Chiellini G, Frascarelli S, Ghelardoni S, Carnicelli V, Tobias S, DeBarber A, Brogioni S, Ronca-Testoni S, Cerbai E, Grandy D, Scanlan T, Zucchi R. Cardiac effects of 3-iodothyronamine: a new aminergic system modulating cardiac function. *FASEB J* 2007;21:1597–1608. [PubMed: 17284482]
5. Doyle K, Suchland K, Ciesielski T, Lessov N, Grandy D, Scanlan T, Stenzel-Poore M. Novel thyroxine derivatives, thyronamine and 3-iodothyronamine, induce transient hypothermia and marked neuroprotection against stroke injury. *Stroke* 2007;38:2569–2576. [PubMed: 17690312]
6. Frascarelli S, Ghelardoni S, Chiellini G, Vargiu R, Ronca-Testoni S, Scanlan TS, Grandy DK, Zucchi R. Cardiac effects of trace amines: pharmacological characterization of trace amine-associated receptors. *Eur J Pharmacol* 2008;587:231–6. [PubMed: 18486124]
7. Regard JB, Kataoka H, Cano DA, Camerer E, Yin L, Zheng YW, Scanlan TS, Hebrok M, Coughlin SR. Probing cell type-specific functions of Gi in vivo identifies GPCR regulators of insulin secretion. *J Clin Invest* 2007;117:4034–43. [PubMed: 17992256]
8. Zucchi R, Ghelardoni S, Chiellini G. Cardiac effects of thyronamines. *Heart Fail Rev*. 2008 Nov 19; [Epub ahead of print]
9. Bräulke LJ, Klingenspor M, DeBarber A, Tobias SC, Grandy DK, Scanlan TS, Heldmaier G. 3-Iodothyronamine: a novel hormone controlling the balance between glucose and lipid utilisation. *Journal of Comparative Physiology B* 2008;178:167–77.

10. Dhillon WS, Bewick GA, White NE, Gardiner JV, Thompson EL, Bataveljic A, Murphy KG, Roy D, Patel NA, Scutt JN, Armstrong A, Ghatei MA, Bloom SR. The thyroid hormone derivative 3-iodothyronamine increases food intake in rodents. *Diabetes Obes Metab*. 2008 Jul 29;[Epub ahead of print]
11. Hart M, Suchland K, Miyakawa M, Bunzow J, Grandy D, Scanlan T. Trace amine-associated receptor agonists: Synthesis and evaluation of thyronamines and related analogues. *J Med Chem* 2006;49:1101–1112. [PubMed: 16451074]
12. Wainwright D, Little S, Yin T, Tu Y, Rocco V, He J, Nelson D. Pharmacologic characterization of the cloned human trace amine-associated receptor1 (TAAR1) and evidence for species differences with the rat TAAR1. *J Pharmacol Exp Ther* 2007;320:475–485. [PubMed: 17038507]
13. Snead A, Santos M, Seal R, Miyakawa M, Edwards R, Scanlan T. Thyronamines inhibit plasma membrane and vesicular monoamine transport. *ACS Chem Biol* 2007;2:390–398. [PubMed: 17530732]
14. Borowsky B, Adham N, Jones K, Raddatz R, Artymyshyn R, Ogozalek K, Durkin M, Lakhani P, Bonini J, Pathirana S, Boyle N, Pu X, Kouranova E, Lichtblau H, Ochoa F, Branchek T, Gerald C. Trace amines: Identification of a family of mammalian G protein-coupled receptors. *Proc Natl Acad Sci U S A* 2001;98:8966–8971. [PubMed: 11459929]
15. Bunzow J, Sonders M, Arttamangkul S, Harrison L, Zhang G, Quigley D, Darland T, Suchland K, Pasumamula S, Kennedy J, Olson S, Magenis R, Amara S, Grandy D. Amphetamine, 3,4-methylenedioxymethamphetamine, lysergic acid diethylamide, and metabolites of the catecholamine neurotransmitters are agonists of a rat trace amine receptor. *Mol Pharmacol* 2001;60:1181–1188. [PubMed: 11723224]
16. Lindemann L, Ebeling M, Kratochwil N, Bunzow J, Grandy D, Hoener M. Trace amine-associated receptors form structurally and functionally distinct subfamilies of novel G protein-coupled receptors. *Genomics* 2005;85:372–385. [PubMed: 15718104]
17. Gloriam DE, Bjarnadottir TK, Schiöth HB, Fredriksson R. High species variation within the repertoire of trace amine receptors. *Ann N Y Acad Sci* 2005;1040:323–7. [PubMed: 15891052]
18. Gloriam DE, Bjarnadottir TK, Yan YL, Postlethwait JH, Schiöth HB, Fredriksson R. The repertoire of trace amine G-protein-coupled receptors: large expansion in zebrafish. *Mol Phylogenet Evol* 2005;35:470–82. [PubMed: 15804416]
19. Grandy D. Trace amine-associated receptor 1-Family archetype or iconoclast? *Pharmacology and Therapeutics* 2007;116:355–90. [PubMed: 17888514]
20. Reese E, Bunzow J, Arttamangkul S, Sonders M, Grandy D. Trace amine-associated receptor 1 displays species-dependent stereoselectivity for isomers of methamphetamine, amphetamine, and para-hydroxyamphetamine. *J Pharmacol Exp Ther* 2007;321:178–186. [PubMed: 17218486]
21. Zucchi R, Chiellini G, Scanlan T, Grandy D. Trace amine-associated receptors and their ligands. *Br J Pharmacol* 2006;149:967–978. [PubMed: 17088868]
22. Liberles S, Buck L. A second class of chemosensory receptors in the olfactory epithelium. *Nature* 2006;442:645–650. [PubMed: 16878137]
23. Tan E, Miyakawa M, Bunzow J, Grandy D, Scanlan T. Exploring the structure-activity relationship of the ethylamine portion of 3-iodothyronamine for rat and mouse trace amine-associated receptor 1. *J Med Chem* 2007;50:2787–2798. [PubMed: 17497842]
24. Tan ES, Groban ES, Jacobson MP, Scanlan TS. Toward deciphering the code to aminergic G protein-coupled receptor drug design. *Chem Biol* 2008;15:343–53. [PubMed: 18420141]
25. Bridges TM, Lindsley CW. G-protein-coupled receptors: from classical modes of modulation to allosteric mechanisms. *ACS Chem Biol* 2008;3:530–41. [PubMed: 18652471]
26. Cherezov V, Rosenbaum DM, Hanson MA, Rasmussen SG, Thian FS, Kobilka TS, Choi HJ, Kuhn P, Weis WI, Kobilka BK, Stevens RC. High-resolution crystal structure of an engineered human beta2-adrenergic G protein-coupled receptor. *Science* 2007;318:1258–65. [PubMed: 17962520]
27. Rosenbaum DM, Cherezov V, Hanson MA, Rasmussen SG, Thian FS, Kobilka TS, Choi HJ, Yao XJ, Weis WI, Stevens RC, Kobilka BK. GPCR engineering yields high-resolution structural insights into beta2-adrenergic receptor function. *Science* 2007;318:1266–73. [PubMed: 17962519]
28. Tota MR, Candelore MR, Dixon RA, Strader CD. Biophysical and genetic analysis of the ligand-binding site of the beta-adrenoceptor. *Trends Pharmacol Sci* 1991;12:4–6. [PubMed: 1848734]

29. Liapakis G, Ballesteros J, Papachristou S, Chan W, Chen X, Javitch J. The forgotten serine - A critical role for Ser-203(5.42) in ligand binding to and activation of the beta(2)-adrenergic receptor. *J Biol Chem* 2000;275:37779–37788. [PubMed: 10964911]
30. Shi L, Javitch J. The binding site of aminergic G protein-coupled receptors: The transmembrane segments and second extracellular loop. *Annu Rev Pharmacol Toxicol* 2002;42:437–467. [PubMed: 11807179]
31. Strader C, Candelore M, Hill W, Dixon R, Sigal I. A single amino-acid substitution in the beta-adrenergic-receptor promotes partial agonist activity from antagonists. *J Biol Chem* 1989;264:16470–16477. [PubMed: 2570781]
32. Strader C, Candelore M, Hill W, Sigal I, Dixon R. Identification of 2 serine residues involved in agonist activation of the beta-adrenergic-receptor. *J Biol Chem* 1989;264:13572–13578. [PubMed: 2547766]
33. Strader C, Fong T, Tota M, Underwood D, Dixon R. Structure and function of G-protein-couple receptors. *Annu Rev Biochem* 1994;63:101–132. [PubMed: 7979235]
34. Strader C, Sigal I, Candelore M, Rands E, Hill W, Dixon R. Conserved aspartic-acid residue-79 and residue-113 of the beta-adrenergic-receptor have different roles in receptor function. *J Biol Chem* 1988;263:10267–10271. [PubMed: 2899076]
35. Wieland K, Zuurmond H, Krasel C, Ijzerman A, Lohse M. Involvement of Asn-293 in stereospecific agonist recognition and in activation of the beta(2)-adrenergic receptor. *Proc Natl Acad Sci U S A* 1996;93:9276–9281. [PubMed: 8799191]
36. Zuurmond H, Hessling J, Bluml K, Lohse M, Ijzerman A. Study of interaction between agonists and Asn293 in helix VI of human beta(2)-adrenergic receptor. *Mol Pharmacol* 1999;56:909–916. [PubMed: 10531394]
37. Suryanarayana S, Daunt D, Von Zastrow M, Kobilka B. A point mutation in the 7th hydrophobic domain of the alpha-2 adrenergic-receptor increases its affinity for a family of beta-receptor-antagonists. *J Biol Chem* 1991;266:15488–15492. [PubMed: 1678390]
38. Ballesteros J, Weinstein H. Integrated methods for the construction of three-dimensional models of structure-function relations in G protein-coupled receptors. *Methods Neuroscience* 1995;25:366–428.
39. Jacobson MP, Pincus DL, Rapp CS, Day TJ, Honig B, Shaw DE, Friesner RA. A hierarchical approach to all-atom protein loop prediction. *Proteins* 2004;55:351–67. [PubMed: 15048827]
40. Jacobson M, Kaminski G, Friesner R, Rapp C. Force field validation using protein side chain prediction. *J Phys Chem B* 2002;106:11673–11680.
41. Jorgensen W, Maxwell D, Tirado-Rives J. Development and testing of the OPLS all-atom force field on conformational energetics and properties of organic liquids. *J Am Chem Soc* 1996;118:11225–11236.
42. Kaminski G, Friesner R, Tirado-Rives J, Jorgensen W. Evaluation and reparametrization of the OPLS-AA force field for proteins via comparison with accurate quantum chemical calculations on peptides. *J Phys Chem B* 2001;105:6474–6487.
43. Gallicchio E, Zhang LY, Levy RM. The SGB/NP hydration free energy model based on the surface generalized born solvent reaction field and novel nonpolar hydration free energy estimators. *J Comput Chem* 2002;23:517–29. [PubMed: 11948578]
44. Ghosh A, Rapp C, Friesner R. Generalized born model based on a surface integral formulation. *J Phys Chem B* 1998;102:10983–10990.

Figure 1.

Thyroid hormone, 3-iodothyronamine, and related analogs. Structures of thyroxine, 3-iodothyronamine (**1**), β-phenylphenoxyphenethylamines (**2** and **3**), and phenoxynaphethylamine (**4**). The A, B, and C rings of **2** and **3** correspond to the outer, inner, and β-phenyl rings, respectively.

Figure 2. Biogenic amine GPCRs. (a) Schematic representation of the helical arrangement of GPCRs viewed from the cell membrane. (b) Binding orientation of (R)-epinephrine in the binding site of the β_2 AR. (c) Proposed binding orientation of **2** in the binding site of rTAAR₁. (d) Binding site of mTAAR₁. (e) Binding site of hTAAR₁. The binding sites of β_2 AR, rTAAR₁, and mTAAR₁ are viewed from the perspective TM4. The rotamer switch residues (white letters), proposed binding and specificity determinant residues are labeled. Non-conserved residues that were mutated are shown in red.

Figure 3.

Selectivity profiles of **3** and **5** on rat and mouse TAAR₁ wild type and mutant receptors. The top and bottom panels show the potency and efficacy, respectively, of **3** and **5**. The EC₅₀ and E_{max} for rTAAR₁ and mTAAR₁ receptors are depicted in solid and hollow symbols, respectively. WT rTAAR₁ [■], rTAAR₁(M6.55T) [▲], rTAAR₁(N7.39Y) [●], rTAAR₁(M6.55T/N7.39Y) [◆], WT mTAAR₁ [□], mTAAR₁(T6.55M) [△], mTAAR₁(Y7.39N) [○], and mTAAR₁(T6.55M/Y7.39N) [◇]. The lines connecting the solid and hollow symbols represent the difference in potency (top panels) and efficacy (bottom panels) between rat and mouse TAAR₁ receptors. The fold difference in potency and percent difference in efficacy are listed above (top panels) and to the left (bottom panels) of the connecting lines, respectively.

Figure 4.

Proposed binding modes of **7**, **1**, and **tyramine** in rTAAR₁. The binding site of rTAAR₁ is viewed from the perspective of TM4 (see Fig. 2a). The rotamer switch residues (white letters), proposed binding and specificity determinant residues are labeled. Residue A5.42 is shown in red. The similar and distinct binding mode models of **1** and **tyramine** are outlined in green and blue dashed lines, respectively.

Figure 5.

Sequence comparison of rat and mouse TAAR₁. Dots represent conserved residues. Amino and carboxy termini (N-terminus and C-terminus, respectively), intracellular loops (IL), extracellular loops (EL), and transmembrane regions (TM) are labeled. The most conserved residue in each TM region is labeled X.50 and residue 4.56 is highlighted in green.

Table 2
Agonist activity of **1**, **3–5**, and **11** on rTAAR₁ TM 6 and/or 7 mutants

rTAAR ₁									
M6.55T			N7.39Y			M6.55T/N7.39Y			
Compd	EC ₅₀ ^a ± SEM (nM)	E _{max} ^b ± SEM (%)	N ^c	EC ₅₀ ^a ± SEM (nM)	E _{max} ^b ± SEM (%)	N ^c	EC ₅₀ ^a ± SEM (nM)	E _{max} ^b ± SEM (%)	N ^c
1	129 ± 43	100 ± 0	2	135	100	1	90	100	1
3	55 ± 20	119 ± 6	2	~10,000	85	1	>10,000	46	1
4	59 ± 0	117 ± 0	2	146	129	1	75	111	1
5	11 ± 1	117 ± 20	2	~1,000	126	1	~2,000	70	1
mTAAR ₁									
T6.55M			Y7.39N			T6.55M/Y7.39N			
Compd	EC ₅₀ ^a ± SEM (nM)	E _{max} ^b ± SEM (%)	N ^c	EC ₅₀ ^a ± SEM (nM)	E _{max} ^b ± SEM (%)	N ^c	EC ₅₀ ^a ± SEM (nM)	E _{max} ^b ± SEM (%)	N ^c
1	1418 ± 592	100 ± 0	2	251 ± 43	100 ± 0	5	208 ± 60	100 ± 0	3
3	>10,000	20 ± 4	2	102 ± 21	95 ± 6	5	43 ± 5	110 ± 8	3
4	350 ± 101	96 ± 13	2	173 ± 41	93 ± 15	2	-	-	-
5	>10,000	45 ± 5	2	176 ± 32	91 ± 11	5	138 ± 24	101 ± 9	3
11	-	-	-	179 ± 21	92 ± 6	3	134 ± 43	101 ± 11	3

^{a-c} See footnotes for Table 1. Compound structures are shown in Table 1.

Table 3Agonist activity of **1**, **4**, **11**, and **tyramine** on TAAR₁ TM 4 or 5 mutantsa. TAAR₁ TM 5 mutants

Compd	rTAAR ₁								
	A5.42L			A5.42I			A5.42T		
	EC ₅₀ ^a	E _{max} ^b	N ^c	EC ₅₀ ^a	E _{max} ^b	N ^c	EC ₅₀ ^a	E _{max} ^b	N ^c
	± SEM (nM)	± SEM (%)		± SEM (nM)	± SEM (%)		± SEM (nM)	± SEM (%)	
	58 ± 16	100 ± 0	3	108 ± 14	100 ± 0	3	88 ± 13	100 ± 0	4
Tyramine	~2,000	102 ± 18	3	>10,000	49 ± 7	3	>10,000	69 ± 5	4
	32 ± 12	101 ± 8	3	29 ± 4	94 ± 2	3	25 ± 3	117 ± 13	4

b. TAAR ₁ TM 4 mutants						
	rTAAR ₁			rTAAR ₁		
	A5.42L			A5.42I		
	EC ₅₀ ^a	E _{max} ^b	N ^c	EC ₅₀ ^a	E _{max} ^b	N ^c
Compd	± SEM (nM)	± SEM (%)		± SEM (nM)	± SEM (%)	
1	38 ± 11	100 ± 0	4	67 ± 17	100 ± 0	6
Tyramine	54 ± 11	108 ± 6	4	40 ± 13	118 ± 8	6
4	22 ± 5	101 ± 14	4	100 ± 16	123 ± 7	6
11	27 ± 15	101 ± 6	4	123 ± 41	105 ± 5	6

^{a-c} See footnotes for Table 1. Compound structures are shown in Table 1.