Published in final edited form as:

J Am Chem Soc. 2009 March 18; 131(10): 3470–3471. doi:10.1021/ja900536d.

Total Synthesis of the Lycopodium Alkaloid (+)-Serratezomine A

Aroop Chandra, Julie A. Pigza, Jeong-Seok Han, Daniel Mutnick, and Jeffrey N. Johnston*

Department of Chemistry & Vanderbilt Institute of Chemical Biology, Vanderbilt University, Nashville, TN 37235-1822

Abstract

key reactions:

- free radical-mediated vinyl amination
- Ce(IV)-promoted oxidative allylation

The first total synthesis of (+)-serratezomine A is described. Key aspects of the synthesis include a) the first deployment of free radical-mediated vinyl amination – an intramolecular alkyne aminostannation – in a complex target synthesis, b) the use of a β -stannyl enamine as the lynchpin for convergent assembly of the natural product backbone, c) the use of an oxidative allylation promoted by cerium(IV) (CAN) to establish the all carbon quaternary chiral center with the proper configuration, and d) an intramolecular substitution reaction to form the sensitive bridging lactone. Overall, 15 steps (longest linear sequence) are required to prepare the natural product from a commercially available aldehyde, and assembly of the contiguous array of six stereocenters is accomplished with high stereocontrol.

The *Lycopodium* alkaloids have provided some of the most intricately structured and stereochemically dense specimens among naturally occurring small molecules.¹ The attendant challenge for chemical synthesis, often combined with their promise as potential therapeutics, has led to broad interest in the total chemical synthesis of these targets.² Recent reports have detailed rather concise preparations of (–)-fawcettimine,³ (+)-fawcettidine,⁴ (–)-lycopodine,⁵ lyconadin A and B,⁶ and (–)-cernuine.^{7,8} Importantly, valuable new chemical methods have been developed in nearly every case to solve specific synthetic challenges while also avoiding synthetic routes that would not address the important issue of supply.

E-mail: jeffrey.n.johnston@vanderbilt.edu.

Supporting Information Available: Preparations, spectral data, and structure elucidation for key intermediates. This material is available free of charge via the Internet at <http://pubs.acs.org>.

Kobayashi isolated (+)-serratezomine A (**1**, as its trifluoroacetic acid salt) as part of a search for new therapeutics from the club moss *L. serratum*, from which the huperzine alkaloids were isolated.⁹ The polycyclic structure merges a 5,6-fused ring system (an indolizidine) with a 6,6-spirocyclic ring and a bridging lactone. Kobayashi found that the lactone is subject to isomerization from O13 to O8, resulting in the thermodynamically favored fused γ -lactone.¹⁰ Among the six contiguous stereogenic carbons, the axial alcohol at C8 – absent from the natural products listed above – sterically impinges upon the spirocyclic piperidine through a 1,3-diaxial interaction. Finally, all but two of the stereochemical elements overlap with a bridgehead or bicyclic union. We anticipated that these challenges might provide an opportunity to develop new chemistry to meet the goal of a convergent synthesis. In this Communication, we describe these developments in the context of the first total synthesis of (+)-serratezomine A.

Our solution to the problem of convergency rested on the hypothesis that an exocyclic enamine of general type **A** (Figure 1), an unsaturated 2-methyl pyrrolidine, could provide a multiply nucleophilic C12 and serve as the synthesis lynchpin. The first step toward annulation would involve acylation of the enamine with a structure of type **B**, and subsequent enamine conjugate addition would complete the cyclohexannulation. Preparation of acid chloride **B** could be similarly convergent through the application of a Brown crotylation to establish the proper relative and absolute configuration at C15/C8. Finally, we intended to annulate the spirocyclic piperidine by alkylation sequentially at carbon and nitrogen with a functional equivalent to **C**.

We targeted β -stannyl enamine **3** (Scheme 1) to serve the role of **A** (Figure 1), as the terminal stannyl group would presumably facilitate the first carbon-carbon bond forming step to regenerate the enamine. Alkynyl imine **2** was prepared by condensation of *para*-methoxy acetophenone with the known 5-amino-1-pentyne.¹¹ Application of radical generating conditions ($n\text{Bu}_3\text{SnH}$, AIBN, 90 °C) provided the desired β -stannyl enamine **3** through a sequence of stannyl radical addition to the terminal alkynyl carbon, followed by addition of the nascent vinyl radical to the nitrogen of the azomethine.¹² Regiocontrol in this addition step is achieved through the use of a ketimine that electronically encourages the formation of the α -amino methyl radical, and sterically retards addition to the azomethine carbon through disubstitution.¹³

Fragment **B** (Figure 1) was prepared from commercially available aldehyde **5**, which can also be obtained directly from ozonolysis of inexpensive ethyl sorbate in 93% yield (Scheme 2).¹⁴ Application of Brown's crotylation reagent,¹⁵ and silylation of the secondary alcohol delivered the unsaturated ester in 76% yield (2 steps). The Mosher ester of **6** was prepared in order to assign the absolute configuration as shown.¹⁶ Hydroboration/oxidation of the terminal olefin provided the primary alcohol in 82% yield, and this was immediately oxidized to **8** using a two step method (92% overall). The acid chloride generated from **8** was then treated with freshly prepared β -stannyl enamine **3**, resulting in the desired vinylogous amide (**9**) isolated in 65% yield (2 steps from acid **8**). Our efforts to coerce this *N*-alkyl vinylogous amide to engage the unsaturated ester were uniformly unsuccessful. However, oxidative dealkylation at nitrogen using CAN ($\text{Ce}(\text{NH}_4)_4(\text{NO}_2)_6$) resulted in the desired cyclization to **10**. We anticipated that the tetrasubstituted alkene would provide sufficient A^{1,3}-strain¹⁷ so as to favor axial orientation of the ethyl acetate ($-\text{CH}_2\text{CO}_2\text{Et}$) substituent at C7.

Indeed, measured coupling constants and NOESY measurements were consistent with the diaxial conformer and *Z*-alkene depicted for *N*-H vinylogous amide **10** in Scheme 2. A range of alkylations were evaluated next in an attempt to establish the proper configuration at the central carbon of the β -imino ketone, as this hindered position is the common link (C12) between the two rings of the spirocycles. Unfortunately, nearly all activation methods simply

promoted intramolecular lactamization with the terminal ester. We found one exception in the form of an oxidative allylation developed by Hwu,¹⁸ in which cerium(IV) and allyl trimethyl silane combine with the vinylogous amide to deliver β -imino ketone **11** in 56% yield and 23:1 dr. Coupling constant and NOESY analysis are consistent with the chair conformer depicted for **11**.¹⁹ From this intermediate containing all backbone carbons necessary for the natural product, a series of final bond forming reactions commenced.

Borohydride reduction of **11** provided the equatorial alcohol **12** in 84% yield (3:1 dr), and was readily converted to its mesylate (**13**) in 98% yield. Hydroboration/oxidation of the terminal olefin was notably sluggish, but led to the desired primary alcohol **14** (51% yield, 86% conversion). This was converted to its mesylate which cyclized slowly upon standing. Treatment of the iminium ion (**15**) with ammonium chloride was necessary to prevent its tautomerization to the corresponding enamine, as the latter undergoes efficient cyclization to the C-acyl enamide (C3, pyrroline numbering). Moreover, this undesired reaction could be avoided almost completely so long as iminium reduction by sodium cyanoborohydride commenced once cyclization to the iminium was complete. Inspection of models suggested that hydride approach to **15** might be less hindered from the β -face, but perhaps more important is that the interactions between the pyrrolidine and cyclohexane rings in **16** are considerably less than in *epi*-**16** (at C4). The final two steps of the synthesis involved a tandem saponification/intramolecular S_N2 cyclization to form the bridging lactone,²⁰ and careful desilylation so as to avoid formation of the fused lactone.^{10,21} Synthetic (+)-serratezomine A, so formed, was identical spectroscopically (¹H & ¹³C NMR)²² to reported values. Optical rotation ([α]_D +9.5 (c 0.3, MeOH)) was also consistent with the literature value ([α]_D +13 (c 0.5, MeOH)), thereby providing further confirmation of absolute configuration.

In summary, the first total synthesis of (+)-serratezomine A is described. The use of β -stannyl enamine **3** as a synthesis lynchpin, the stereocontrolled, a strategically convergent assembly of fragments (**3**, **8**, and allyl trimethyl silane), and the minimal use of protecting groups combine to provide a reasonably short chemical synthesis of one of the most structurally and stereochemically complex members of the alkaloids *lycopodium*. The challenges posed by serratezomine A inspired the development of free radical-mediated vinyl amination (**2**→**3**), and led to the application of cerium(IV) oxidative allylation (**10**→**11**) as a means to establish the congested all carbon quaternary center with the proper configuration needed for serratezomine A.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgements

Financial support was provided by the NIH (GM063557), an Eli Lilly Grantee award, and a Pfizer Diversity Fellowship (JAP). We are grateful to Mr. Matthew Mulder for mass-directed LC purification of the final product.

References

1. Ma XQ, Gang DR. Natural Product Reports 2004;21:752. [PubMed: 15565253] Kobayashi, JI.; Morita, H. Alkaloids. San Diego, CA: United States; 2005. p. 1
2. Selected examples: Kozaka T, Miyakoshi N, Mukai C. J Org Chem 2007;72:10147. [PubMed: 18044932] Paquette LA, Friedrich D, Pinard E, Williams JP, StLaurent D, Roden BA. J Am Chem Soc 1993;115:4377. Hirst GC, Johnson TO Jr, Overman LE. J Am Chem Soc 1993;115:2992. Qian L, Ji R. Tetrahedron Lett 1989;30:2089. Heathcock CH, Kleinman EF, Binkley ES. J Am Chem Soc 1982;104:1054. Harayama T, Takatani M, Inubushi Y. Chem Pharm Bull 1980;28:2394. Kim SW, Bando Y, Takahashi N, Horii Z. Chem Pharm Bull 1978;26:3150. Harayama T, Ohtani M, Oki M,

- Inubushi Y. *J Chem Soc, Chem Commun* 1974:827.Scott WL, Evans DA. *J Am Chem Soc* 1972;94:4779. [PubMed: 5036174]Wiesner K, Poon L. *Tetrahedron Lett* 1967:4937. [PubMed: 6062698]
3. Linghu X, Kennedy-Smith JJ, Toste FD. *Angew Chem Int Ed* 2007;46:7671. See ref. 9–11 in this report for preparations of rac-fawcettimine.
 4. Kozak JA, Dake GR. *Angew Chem Int Ed* 2008;47:4221.
 5. Yang H, Carter RG, Zakharov LN. *J Am Chem Soc.* 2008 See ref. 5 in this report for preparations of rac-lycopodine.
 6. (+)-Lyonadin A and (–)-lyconadin B: Beshore DC, Smith AB. *J Am Chem Soc* 2007;129:4148. [PubMed: 17367142]Beshore DC, Smith AB. *J Am Chem Soc* 2008;130:13778. [PubMed: 18798629] rac-Lyonadin A: Bisai A, West SP, Sarpong R. *J Am Chem Soc* 2008;130:7222. [PubMed: 18479095]
 7. Nishikawa Y, Kitajima M, Takayama H. *Org Lett* 2008;10:1987. [PubMed: 18429621]
 8. See also: Shigeyama T, Katakawa K, Kogure N, Kitajima M, Takayama H. *Org Lett* 2007;9:4069. [PubMed: 17824711]Evans DA, Scheerer JR. *Angew Chem Int Ed* 2005;44:6038.Yen CF, Liao CC. *Angew Chem Int Ed* 2002;41:4090.Sha CK, Lee FK, Chang CJ. *J Am Chem Soc* 1999;121:9875.
 9. Morita H, Arisaka M, Yoshida N, Kobayashi J. *J Org Chem* 2000;65:6241. [PubMed: 10987968]
 10. Morita H, Kobayashi J. *J Org Chem* 2002;67:5378. [PubMed: 12126432]
 11. Li YW, Marks TJ. *J Am Chem Soc* 1996;118:9295.
 12. Nugent BM, Williams AL, Prabhakaran EN, Johnston JN. *Tetrahedron* 2003;59:8877.Prabhakaran EN, Nugent BM, Williams AL, Nailor KE, Johnston JN. *Org Lett* 2002;4:4197. [PubMed: 12443057]
 13. Williams AL, Srinivasan JM, Johnston JN. *Org Lett* 2006;8:6047. [PubMed: 17165926]Srinivasan JM, Burks HE, Smith CR, Viswanathan R, Johnston JN. *Synthesis* 2005:330.Viswanathan R, Prabhakaran EN, Plotkin MA, Johnston JN. *J Am Chem Soc* 2003;125:163. [PubMed: 12515518] Viswanathan R, Mutnick D, Johnston JN. *J Am Chem Soc* 2003;125:7266. [PubMed: 12797800] Johnston JN, Plotkin MA, Viswanathan R, Prabhakaran EN. *Org Lett* 2001;3:1009. [PubMed: 11277782]
 14. Garaas SD, Hunter TJ, O'Doherty GA. *J Org Chem* 2002;67:2682. [PubMed: 11950317]Hunter TJ, O'Doherty GA. *Org Lett* 2001;3:2777. [PubMed: 11506632]
 15. Toshima K, Jyojima T, Miyamoto N, Katohno M, Nakata M, Matsumura S. *J Org Chem* 2001;66:1708. [PubMed: 11262117]Brown HC, Bhat KS. *J Am Chem Soc* 1986;108:293.
 16. Kusumi T, Hamada T, Ishitsuka MO, Ohtani I, Kakisawa H. *J Org Chem* 1992;57:1033.Ohtani I, Kusumi T, Kashman Y, Kakisawa H. *J Am Chem Soc* 1991;113:4092.
 17. Hoffmann RW. *Chem Rev* 1989;89:1841.
 18. β -Diketones: Hwu JR, Chen CN, Shiao SS. *J Org Chem* 1995;60:856.Zhang Y, Raines AJ, Flowers RA. *Org Lett* 2003;5:2363. [PubMed: 12816449]Zhang Y, Raines AJ, Flowers RA. *J Org Chem* 2004;69:6267. [PubMed: 15357585]
 19. The conformers depicted in Scheme 2 are believed to predominate in solution and were assigned using coupling constant analysis and/or 2D NMR techniques in each case. However, the ^1H NMR spectrum for mesylate 16 was severely broadened.
 20. Ziegler FE, Klein SI, Pati UK, Wang TF. *J Am Chem Soc* 1985;107:2730.
 21. Purification of 1 was performed similarly to that described by Kobayashi (ref. 9). However, the nature of mass directed LC leads to some material loss.
 22. See Supporting Information for comparisons.

Figure 1.
(+)-Serratezomine A: A Convergent Construction Strategy

Scheme 1. Use of Free Radical-Mediated Alkyne Aminostannation to Prepare β -Stannyl Enamine **3**

Scheme 2.
Completion of the Total Synthesis of (+)-Serratezomine A