Published in final edited form as:

Tetrahedron Lett. 2009 April 8; 50(14): 1523–1525. doi:10.1016/j.tetlet.2009.01.046.

Thio-mediated two-component coupling reaction of carboxylic acids and isonitriles under mild conditions

Xiangyang Wu^a, Xuechen Li^a, and Samuel J. Danishefsky^{a,b}

^aLaboratory for Bioorganic Chemistry, Sloan-Kettering Institute for Cancer Research, 1275 York Avenue, New York, NY 10065, USA

^bDepartment of Chemistry, Columbia University, 3000 Broadway, New York, NY 10027, USA

Abstract

The coupling reaction between carboxylic acids and isonitriles in the presence of thio phenol as activator under mild conditions is described.

Recently, our laboratory described some conceptually noteworthy and translationally promising results arising from reactions of carboxylic acids (**1**) with isonitriles (**2**).¹ Several notions and terms have been suggested to help facilitate communication in this rapidly emerging chemistry. Equation 1 describes the formation of formimidate carboxylate mixed anhydrides (FCMA) in *E* and/or *Z* form via the merger of **1** and **2** (Figure 1). Equation 2 corresponds to a 1,3-O→N acyl migration to generate an N-formyl amide (**4**). This migration is the key and terminating step in the overall two-component coupling (2CC) process. Depending on how the experiment is structured, the O→N acyl migration within FCMA **3** may be interdicted by a resident nucleophile (NuH) to afford **5**.² Several reports³ in the literature which did not appreciate the high susceptibility of FCMA to nucleophile attack, have been corrected.⁴ We interpret the formation of **5** at room temperature, wherein the formation of 2CC product **4** is *not* observed, to suggest that FCMA **3** (*Z* or *E*) is readily produced, albeit in low concentration. However, the 1,3-O→N acyl transfer requires thermolysis (best done via microwave heating at 150 °C). In earlier papers, we described some of the high “value added” chemistry which accrues from readily achievable transformations of the formyl group of **4** (see Figure 1, product types **6**). Moreover, the 2CC chemistry, generalized above, has been shown to be applicable to the synthesis of N-linked small peptides and glycopeptides.¹

Of course, the main disadvantage in the 1,3-O→N acyl transfer step (cf. **3**→**4**) is the requirement for thermolytic activation. Such conditions may or may not be consistent with maintenance of a sensitive polypeptidic or polypeptidic glycan structure.

The work described herein was motivated by the hope of realizing 2CC reactions, ideally *at room temperature but certainly well below the current microwave (150 °C) conditions*. As is described below, important progress in this regard has been realized.

Correspondence to: Samuel J. Danishefsky.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

That the FCMA is produced, albeit slowly, at room temperature, had been established through interdiction experiments, resulting in the formation of secondary amide **9**, albeit in modest yield (Figure 2).¹

We were also intrigued by the implications of our findings in the context of an intended model experiment for achieving “*serine ligation*.”^{4,5} Remarkably, reaction of L-serine isonitrile **10** with acid **11** afforded **15** (Figure 2). Although the yield is modest, we noted that when the side chain hydroxyl of the serine was protected, no coupling occurred at room temperature. However, the microwave (150 °C) induced 2CC coupling in the –OTMS protected series occurred normally (**16**→**17**→**18**). When the silyl ether of **18** was cleaved, the same formate ester, **15**, was smoothly produced, thus corroborating the various structural assignments.

The mechanistic inferences to be drawn from the “serine” experiments are far from certain. Perhaps the free-hydroxyl group adds to the imino linkage of the FCMA **12** to generate **13**. Following this line of reasoning, one would be obliged to conclude that the acyl transfer in **13** is significantly faster than are its counterparts, **3** (see **3**→**4**) or **17** (see **17**→**18**). Alternatively, the role of the free hydroxyl group in the serine might be that of promoting (via **13**) the formation of FCMA **12-E** required for acyl transfer.⁶ Still another possibility to account for the serine acceleration focuses on the “late stage” of the acyl transfer shown in Figure 2.⁷ It contemplates intervention of a stabilizing hydrogen bond to the emerging imide (at either the nitrogen or the formyl lone pairs) which helps facilitate the culminating O→N acyl transfer of the 2CC reaction (see **12**→**14**).

While a full understanding of the detailed mechanism of the 2CC serine reaction leading to **15** is a work in progress, the already intriguing results reported above prompted us to explore the possibility of achieving a similar outcome through mediation by an *external* nucleophile. Pursuing this admittedly speculative line of thought, we studied the consequences of adding thiols to putative 2CC reactions.

Happily, already in the opening experiment, it was found that in the presence of 0.5 eq of thiophenol, with 1,2-dichloroethane as the solvent, coupling of **19** and **7** took place *at room temperature* to provide **20** and **21** in the yields shown. The yields could be improved (ca 60%) by conducting the reaction at 50° C.

The results of our survey investigations are shown in Table 1. While the concurrent formation of N-formyl amides and their “thioacetals” (see **21**) is obviously awkward, the yields for converting type **21** products to **20** are high (88%) so that in the end the 2CC reaction is quite productive.

As was suggested in the discussion of the serine ligation experiment, we are tempted to propose the formation of the tetrahedral intermediate **B** by combining the FCMA **A** with the thio nucleophile (Scheme 2). As before, the tetrahedral intermediate might undergo rapid acyl transfer or revert to the required *E*-FCMA for the acyl transfer step leading to N-formyl amides. Alternatively, the intermolecular thiol SH bond could provide a hydrogen bonding source (as described above) to facilitate the culminating O→N transfer step.

Hoping to gain further insight into the thiol mediated 2CC reaction, we explored the possibility of generating a related tetrahedral intermediate **B** (Scheme 2) by the reaction of an acid with a thioimide. However, attempted aspartylation via the reaction of aspartic acid with independently synthesized thioimide⁹ led to no observed coupling product (Scheme 2).

In summary, we have described herein an improved method for coupling of isonitriles and carboxylic acids in the presence of thiophenol as an activator under mild conditions. This

coupling method, with appropriate fine tuning, could be utilizable in the preparation of important biomolecules.

Acknowledgments

This work was supported by the NIH (CA28824 to S.J.D.). Special thanks go to Dana Ryan and Rebecca Wilson for assistance with the preparation of the manuscript. We thank Dr. Jianglong Zhu, Dr. Yu Yuan and Dr. Cindy Kan for their helpful discussions. We thank Dr. George Sukenick, Ms. Hui Fang, and Ms. Sylvi Rusli of the Sloan-Kettering Institute's NMR core facility for mass spectral and NMR spectroscopic analysis.

References

1. Li X, Danishefsky SJ. J Am Chem Soc 2008;130:5446. [PubMed: 18370392] Li X, Danishefsky SJ. Nature Protocols 2008;3:1666.
2. Li X, Yuan Y, Kan C, Danishefsky SJ. J Am Chem Soc 2008;130:13225. [PubMed: 18783225]
3. a) Shaabani A, Soleimani E, Rezayan AH. Tetrahedron Lett 2007;48:6137. b) Gloede J, Gross H. Zeitsch fuer Chem 1968;8:219.
4. Li X, Yuan Y, Berkowitz WF, Todaro LJ, Danishefsky SJ. J Am Chem Soc 2008;130:13222. [PubMed: 18783224]
5. Similar chemistry in comparable yield has been achieved with the threonine isonitrile.
6. For this interpretation to explain the serine ligation, one would have to propose that the kinetic product of the original insertion reaction between the carboxylic acid and the isonitrile is the Z-FCMA which does not undergo 1,3 O→N transfer. The formation of tetrahedral intermediate **12** allows for equilibration of the FCMA forms, enabling the E-form to rearrange. This formulation seems counterintuitive but is included as a formal possibility. See Marcelli T, Himo F. Eur J Org Chem 2008:4751.
7. The eventual progression to formate ester **15** provides a thermodynamic advantage but is unlikely to explain the sharply accelerated rate of the serine ligation since it comes after the 1,3-N→O acyl transfer.
8. Attempts to use L- α -amino acid derivatives in the thio mediated 2CC reaction have not been rewarding.
9. Saegusa T, Kobayashi S, Hirota K, Okumura Y, Ito Y. Bulletin Chem Soc Jpn 1968;41:1638.

Figure 1.

Possible Mechanistic Rationales:

Figure 2.

Scheme 1.

Key: (a) $\text{CH}_2\text{ClCH}_2\text{Cl}$, PhSH ; (b) TsOH , CH_2Cl_2 .

Scheme 2.

Table 1

Conditions: (a) 50 °C, 48 h.; (b) 80 °C, 48 h.⁸