Published in final edited form as:

Tetrahedron Lett. 2009 June 3; 50(22): 2693–2696. doi:10.1016/j.tetlet.2009.03.148.

Synthesis of (*R*)-3,4-dihydro-2*H*-pyran-2-carboxaldehyde: application to the synthesis of potent adenosine A_{2A} and A₃ receptor agonist

Prakash G. Jagtap^{a,*}, Zhiyu Chen^a, Karsten Koppetsch^a, Elizabeth Piro^a, Paula Fronce^a, Garry J. Southan^a, and Karl-Norbert Klotz^b

^a Inotek Pharmaceuticals Corporation, 33 Hayden Avenue, Lexington, MA 02421, USA

^b Institut für Pharmakologie und Toxikologie, Universität Würzburg, D-97078 Würzburg, Germany

Abstract

Synthesis of potent adenosine A_{2A} and A₃ receptor agonist from the modification of adenosine-5'-*N*-ethylcarboxamide (NECA) has been reported. Diastereoisomer possessing an (*R*) 3,4-dihydro-2*H*-pyran-2-carboxaldehyde (DHP) moiety exhibited the highest affinity at the A_{2A} and A₃ receptors. The key steps involve the synthesis of (*R*)-3,4-dihydro-2*H*-pyran-2-carboxaldehyde (**7**), which was obtained through the enzyme catalyzed kinetic resolution of (±)-2-acetoxymethyl-3,4-dihydro-2*H*-pyran (**5**).

The metabolite adenosine plays a critical role in the regulation of cell function in many tissues. It mediates its actions via four subtypes of G protein-coupled receptors named A₁, A_{2A}, A_{2B} and A₃.¹ Based on adenosine's profound actions on the heart, kidney, brain, and the immune system, numerous potential therapeutic options have been discussed for adenosine receptor agonists and antagonists with selectivity for the known receptor subtypes.²

It has been demonstrated that a lipophilic linker in the N⁶ position is necessary to gain high affinity for the A₃ receptor, and a polar linker possessing an additional lipophilic group in the C-2 position is necessary to gain high affinity for the A_{2A} receptor.² Thus, the major efforts in the discovery of new adenosine A_{2A} and A₃ agonists have been made with the introduction of appropriate substituents at the C-2 or N⁶ position in combination with the modification of the 5' position of the nucleoside scaffold via various linkers.^{2,3}

The stimulation of both A_{2A} and A₃ receptor subtypes have been shown to mediate antiinflammatory actions,⁴ generating interest in such compounds. Our interest in the synthesis of (*R*)-3,4-dihydro-2*H*-pyran-2-carboxaldehyde [(*R*)-acrolein dimer] arose from the investigation of one of our potent adenosine A_{2A} and A₃ receptor agonists **2**, which possesses a (*R*)-3,4-dihydro-2*H*-pyran (DHP) moiety.⁵ During our structure-activity relationships study of 2-hydrazone-NECA derivatives, we discovered that compound **2** exhibits an interesting pharmacological profile. In a radioligand binding assay, it showed high affinity at both the A_{2A} and A₃ receptors. Compound **2** was initially prepared from racemic aldehyde **7** and 2-hydrazino-NECA **1** (Scheme 1). Due to the interesting properties of **2**, diastereomeric

*Corresponding author. Tel.: +1-781-676-2118; fax: +1-781-676-2155; e-mail: pjagtap@inotekcorp.com.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

separation of isomers **3** and **4** from **2** was then performed via preparative HPLC using reverse phase C-8 or C-18 columns. After the resolution of diastereomers, we found that the molecule possessing the (*R*)-DHP ring (**3**) is more potent than the (*S*)-DHP analogue (**4**). The *R*-isomer (**3**) was also the more selective of the two diastereomers (Table 1).

In order to make compound **3**, we required the acrolein dimer (*R*)-**7** on a large scale. However, the lack of a literature method for the synthesis of enantiomers of acrolein dimer led us to investigate a convenient synthetic route for acrolein dimer (*R*)-**7**. The racemic aldehyde has been synthesized from acrolein under high pressure⁶ or microwave assisted⁷ dimerisation (Figure 1). There is no report on the enantio-selective synthesis of acrolein dimer.

Pederson *et al.*⁸ have reported the parallel kinetic resolution of ($\pm$)-acrolein dimer by reacting with chiral phosphonates using asymmetric Horner-Wadsworth-Emmons reaction. Although resolution of the corresponding diastereoisomeric adducts with chiral auxiliaries has been completed, their conversion to pure enantiomers of (*R*) and (*S*)-aldehydes remains to be achieved. Herein, we report the convenient synthesis of (*R*)-**7** and its application in the synthesis of potent adenosine agonist **3** (Scheme 1 and 2).

There are several syntheses⁹ and synthetic applications^{10,11} of the precursor 3,4-dihydro-2*H*-pyran-2-methanol (**6**) reported in literature. We decided to follow the method reported by Kang *et al.*¹² and Ley *et al.*^{11a} for the source of (*R*)-**5** and (*R*)-**6** (Scheme 2), which describes the enzymatic hydrolysis of ($\pm$)-**5**. Kang *et al.*¹² performed enzymatic hydrolysis of ($\pm$)-**5** using porcine pancreatic lipase (PPL), 0.01 M phosphate buffer and acetone that provided (*S*)-alcohol **3** in >99% e.e. However, the process reported by Kang *et al.*¹² produced our desired product (*R*)-**5** only in 48% e.e. Ley *et al.*^{11a} modified the above method¹² but it requires an additional enzymatic hydrolysis and two purification steps.

In an attempt to simplify the above reported methods,^{11a, 12} we performed the enzyme catalyzed kinetic resolution on ($\pm$)-**5** by extending reaction time to 4 days as well as repeating the addition of PPL after the first 24 hours.¹³ These modifications pleasingly lead us to (*R*)-acetate **5** in >99% e.e. in one step instead of (*S*)-alcohol enriched product as reported by Kang *et al.*¹² The (*R*)-acetate was then hydrolyzed to the corresponding (*R*)-alcohol **5** in the presence of 0.3 equivalent of 1.0 N potassium hydroxide.¹⁴

In order to make compound ($\pm$)-**7**, oxidation of commercially available racemic alcohol **6** was first carried out using various reagents. These results are summarized in Table 2. Our initial attempt of a Swern oxidation reaction¹⁵ of ($\pm$)-**6** produced ($\pm$)-**7** in poor yield. Several other oxidation reactions were attempted, which did not result in any trace of the desired product ($\pm$)-**7**: (1) Tetrapropylammonium perruthenate (TPAP), 4-methylmorpholine *N*-oxide (NMO), CH₂Cl₂; (2) MnO₂, acetone; (3) Pyridinium chlorochromate (PCC), ether; (4) Pyridinium dichromate (PDC), CH₂Cl₂.

The reaction of ($\pm$)-**6** in the presence of Dess-Martin periodinane¹⁶ in CH₂Cl₂ provided aldehyde **7** as the major product but the separation of **7** from impurities was complicated. Further attempts to oxidize ($\pm$)-**6** using iodobenzene diacetate (BAIB) and 2,2,6,6-tetramethyl-1-piperidinyloxy (TEMPO) in CH₂Cl₂ provided ($\pm$)-**7** in better yield.¹⁷ Under similar conditions, the enantiomer (*R*)-**7** was then prepared from (*R*)-**6**.¹⁸ (*R*)-Aldehyde was separated from the crude reaction mixture by column chromatography. Of the various oxidizing reagents studied, the method with BAIB and TEMPO was found to be efficient and clean.

Treatment of (*R*)-**7** with 2-hydrazino-NECA (**1**) produced the diastereoisomer **3** enriched with (*R*)-3,4-dihydro-2*H*-pyran-2-yl (DHP) side chain.¹⁹ The (*R*)-DHP analogue **3** showed highest affinity for both A_{2A} and A₃ receptors [K_i(A_{2A}) = 3.76 and K_i(A₃) = 4.51 nM]. Compound **3** is 7-fold better than reference compound CGS 21680, binds with a K_i(A_{2A}) value of 3.76 nM,

and is 12-fold selective *versus* A₁. Compound 3 also showed 10-fold A₃ selectivity versus the A₁ subtype, which is slightly better than the selectivity of CGS21680 as one of the most potent and selective A_{2A} agonists known (Table 1). The adenylyl cyclase functional assay¹ has shown that it is a full agonist at A_{2A} and A₃ receptors (data not shown).

In summary, a synthesis of potent and selective adenosine A_{2A} and A₃ receptor agonist **3** has been accomplished from the coupling reaction of (*R*)-3,4-dihydro-2*H*-pyran-2-carboxaldehyde (**7**) and 2-hydrazino-NECA (**1**). The enantiopure (*R*)-2-acetoxymethyl-3,4-dihydro-2*H*-pyran (**5**) was prepared on multi-gram scale by modifying the earlier PPL catalyzed resolution methods.^{11a, 12} A method for the synthesis of aldehyde (*R*)-**7** from alcohol (*R*)-**6** has been discussed. The results from *in vivo* studies will be published elsewhere.

Acknowledgments

This work was supported by a grant from the National Institutes of Health (R44AI46167). We thank Sonja Kachler for her expert technical assistance.

References and notes

- Fredholm BB, Jzerman IAP, Jacobson KA, Klotz KN, Linden J. *Pharmacol Rev* 2001;53:1–26. [PubMed: 11171937]
- For adenosine receptor agonists and antagonists reviews see: (a) Jacobson KA, Gao ZG. *Nat Rev Drug Discov* 2006;5:247–64. [PubMed: 16518376] (b) Jacobson KA, van Galen PJM, Williams M. *J Med Chem* 1992;35:407–422. [PubMed: 1738138] (c) DeNinno MP. *Adenosine. Annu Rep Med Chem* 1998;33:111–120. (d) Poulsen SA, Quinn RJ. *Bioorg Med Chem* 1998;6:619–641. [PubMed: 9681130] (e) Appleman JR, Erion MD. *Exp Opin Invest Drugs* 1998;7:225–243. (f) Cristalli G, Lambertucci C, Taffi S, Vittori S, Volpini R. *Curr Top Med Chem* 2003;3:387–401. [PubMed: 12570757] (g) Baraldi PG, Cacciari B, Romagnoli R, Merighi S, Varani K, Borea PA, Spalluto G. *Med Res Rev* 2000;20:103–128. [PubMed: 10723024] (h) Akkari R, Burbiel JC, Hockemeyer J, Müller CE. *Curr Top Med Chem* 2006;6:1375–1399. [PubMed: 16918456]
- (a) Niiya K, Olsson RA, Thompson RD, Silvia SK, Ueeda M. *J Med Chem* 1992;35:4557–4561. [PubMed: 1469687] (b) Niiya K, Thompson RD, Silvia SK, Olsson RA. *J Med Chem* 1992;35:4562–566. [PubMed: 1469688] (c) Viziano M, Ongini E, Conti A, Zocchi C, Seminati M, Pocar D. *J Med Chem* 1995;38:3581–3585. [PubMed: 7658444] (d) van Tilburg EW, von Frijtag Drabbe Kunzel J, de Groote M, IJzerman AP. *J Med Chem* 2002;45:420–429. [PubMed: 11784146]
- (a) Salvatore CA, Tilley SL, Latour AM, Fletcher DS, Koller BH, Jacobson MA. *J Biol Chem* 2000;275:4429–4434. [PubMed: 10660615] (b) Sitkovsky MV, Lukashev D, Apasov S, Kojima H, Koshiba M, Caldwell C, Ohta A, Thiel M. *Annu Rev Immunol* 2004;22:657–682. [PubMed: 15032592]
- Jagtap PG, Salzman AL, Szabo C. *WO* 2006034190.
- (a) Powell, JE.; Barker, MD. US patent 4,388,104. (b) Etzkorn, WG.; Galley, RA.; Snead, TE.; Brockwell, JL.; Young, MA.; Maher, JM.; Knight, B. US patent 6,187,793 B1.
- Hsieh HP, Chen ST, Wang KT. *J Chinese Chem Soc* 1997;44:597–600.
- Pedersen TM, Jensen JF, Humble RE, Rein T, Tanner D, Boddman K, Reiser O. *Org Lett* 2000;2:535–538. [PubMed: 10814370]
- (a) Buffet MF, Dixon DJ, Edwards GL, Ley SV, Tate EW. *J Chem Soc Perkin Trans 1* 2000:1815–1827. (b) Fehr C, Galindo J, Ohloff G. *Helv Chim Acta* 1981:1247–1256. (c) Ibrahim N, Eggimann T, Dixon EA, Wieser H. *Tetrahedron* 1990;46:1503–1514. (d) Brimacombe JS, Hunedy F, Mather AM, Tucker LCN. *Carbohydr Res* 1979;68:231–238. (e) Komada H, Okada M, Sumitomo H. *Macromolecules* 1979;12:5–9.
- (a) Thompson LA, Ellman JA. *Tetrahedron Lett* 1994;35:9333–9936. (b) Kick EK, Ellman JA. *J Med Chem* 1995;38:1427–1430. [PubMed: 7739000]
- (a) Ley SV, Mio S, Mesenguer B. *Synlett* 1996:787–788. (b) Lainé D, Fujita M, Ley SV. *J Chem Soc Perkin Trans 1* 1999:1639–1646.

12. Kang SK, Jeon JH, Yamaguchi T, Hong RK, Ko BS. *Tetrahedron: Asymmetry* 1995;6:97–100.
13. Typical procedure for the enzymatic hydrolysis of 2-acetoxymethyl-3,4-dihydro-2*H*-pyran (5): To the mixture of phosphate buffer (4.2 L, 0.01 M, pH 7.6) was added ($\pm$)-5 (29.5 g, 189.1 mmol) in acetone (145 ml) at rt and stirred for 5 min. Then it was treated with PPL (Sigma, 2.8 g) and the reaction mixture was stirred at rt. During the reaction pH was constantly adjusted to 7.6 with NaOH (3N). Additional PPL (0.7 g) was added after 24 hr. Progress of the reaction was monitored by HPLC using chiral column as shown below. After 4 days, the reaction mixture was extracted with EtOAc (4 X 700 ml) and organic layer was dried on Na₂SO₄. It was filtered and concentrated on rotavaporator and purified on the neutral aluminum oxide using 25% EtOAc – hexane to give (*R*)-5 (8.1 gm, 38%). ¹H NMR (300 MHz, CDCl₃) δ 1.6–2.2 (m, 4 H), 2.1 (s, 3 H), 4.2 (m, 3 H), 4.8 (m, 1 H), 6.4 (m, 1 H). [α]_D²⁵ -76.67 (*c* = 2.58 in CHCl₃). Following this method (*R*)-5 was prepared on multi-gram scale (200 g). The enantiomeric excess was determined by HPLC using Chiralpak AD-RH column from Daicel. Conditions: Isocratic using 45% MeOH, 5% MeCN in H₂O for 35 min, flow rate 0.7 ml/min, and detected at 215 nm. The retention time of the (*R*) and the (*S*) isomers were 16.28 and 18.28 min. respectively.
14. (*R*)-3,4-dihydro-2*H*-pyran-2-methanol (6): [α]_D²⁵ -74.61 (*c* = 2.52 in CHCl₃).
15. (a) Paquette LA, Oplinger JA. *J Org Chem* 1988;53:2953–2959. (b) Bianchi P, Roda G, Riva S, Danieli B, Zabelinskaja-Mackova A, Griengl H. *Tetrahedron* 2001;57:2213–2220.
16. (a) Dess DB, Martin JC. *J Org Chem* 1983;48:4155–4156. (b) Dess DB, Martin JC. *J Am Chem Soc* 1991;113:7277–7287.
17. Mico AD, Margarita R, Parlanti L, Vescovi A, Piancatelli G. *J Org Chem* 1997;62:6974–6977.
18. Typical procedure for the oxidation of (*R*)-3,4-dihydro-2*H*-pyran-2-methanol (6): A mixture of (*R*)-alcohol 6 (14.0 g, 122.8 mmol) and BAIB (59.13 g, 184.2 mmol) in CH₂Cl₂ (105 ml) was stirred at rt for 30 min., and then treated with TEMPO (1.92 g, 12.28 mmol). The reaction mixture was stirred at room temperature for 5 hr. Progress of reaction was monitored by tlc (silica gel, 30% EtOAc – hexane, R_f = approx. 0.65). The reaction mixture was then diluted with CH₂Cl₂ (100 ml) and treated with saturated solution of Na₂S₂O₃ (100 ml) and NaCl (2×50 ml). The biphasic mixture was then treated with dilute solution of Na₂CO₃ (to adjust pH of mixture to 6 – 7). The organic layer was separated and the aqueous layer was extracted in CH₂Cl₂ (8 X 20 ml). The combined organic solution was dried over Na₂SO₄ (10 g) and filtered. Then it was concentrated on rotavaporator at 30 °C under vacuum. Purification of crude material on the silica gel column by eluting first with 100% CH₂Cl₂ followed by the mixture of CH₂Cl₂-MeOH (20:1) furnished aldehyde (*R*)-7 (7.5 g, 54%). ¹H NMR (300 MHz, CDCl₃) δ 1.56 – 2.07 (m, 4H), 4.29 – 4.32 (dd, J = 5.4 and 7.5 Hz, 1H), 4.77 – 4.79 (m, 1H), 6.49 (d, J = 6.3 Hz, 1H), 9.71 (s, 1H). Compound (*R*)-7 was found to be unstable if exposed to heat and light. It was stored in a refrigerator using the dark vials.
19. The HPLC analysis of the final product obtained from (*R*)-aldehyde and 2-hydrazino-NECA showed (*R*)-DHP enriched diastereomer 3 as the major product (98.4%). ¹H NMR (300 MHz, DMSO-*d*₆) δ 1.04 (t, J = 6.6 Hz, 3H), 1.89 – 2.08 (m, 4H), 3.15 – 3.20 (m, 3H), 4.22 – 4.24 (m, 1H), 4.36 (d, J = 3.6 Hz, 1H), 4.55–4.61 (m, 2H), 4.80 (d, J = 5.4 Hz, 1H), 5.69 (s, 2H), 5.91 (d, J = 6.3 Hz, 1H), 6.49 (d, J = 6.3 Hz, 1H), 7.56 (d, J = 5.4 Hz, 1H), 8.25 (t, J = 5.1 Hz, 1H), 8.59 (s, 1H).

Figure 1.

Scheme 1.

a. hydrazine hydrate, neat, rt, 7h, 99%; b. MeOH, 50 °C, 78%; c. HPLC purification on C-8 or C-18 column

Scheme 2.

a. Ac_2O , pyridine, rt, 8h, 75%; b. 0.01 M phosphate buffer, acetone, NaOH, PPL, 4 days, *R*-(-)-acetate 31%; c. MeOH, KOH (1N, 0.3 eq), 0 °C - rt, 1 h, 82%; d. BAIB, TEMPO, CH_2Cl_2 , 5 h, 54%; e. **1**, MeOH, rt, 3h, 93%.

Table 1

of the adenosine analogues **2**, **3** and **4** in radioligand binding assays at human A₁, A_{2A}, and A₃ adenosine receptors

position of NECA and compound number.	A ₁ K _i (nM) ^a	A _{2A} K _i (nM) ^b	A _{2B} EC ₅₀ (nM) ^d	A ₃ K _i (nM) ^c	A ₁ /A _{2A}	A ₁ /A ₃
	34.6	6.64	17,400	6.32	5.2	5.47

position of NECA and compound number.

$A_1 K_i$ (nM) ^a	$A_{2A} K_i$ (nM) ^b	$A_{2B} EC_{50}$ (nM) ^d	$A_3 K_i$ (nM) ^c	A_1/A_{2a}	A_1/A_3
46.5	3.76	13,200	4.51	12.36	10.31
76.4	23.3	31,000	16.0	3.27	4.85
290	27	89,000	67	10.7	4.3

Tetrahedron Lett. Author manuscript; available in PMC 2010 June 3.

f specific [³H]CCPA binding in CHO cells stably transfected with human recombinant A₁ adenosine receptor, expressed as K_i (nM).

^b Displacement of specific [³H]NECA binding in CHO cells stably transfected with human recombinant A_{2A} adenosine receptor, expressed as K_i (nM).

^c Displacement of specific [³H]NECA binding in CHO cells stably transfected with human recombinant A₃ adenosine receptor, expressed as K_i (nM).

^d Activation of adenylyl cyclase in membranes from CHO cells with stably transfected human A_{2B} receptors.

^e Data from reference ¹.

Table 2Oxidation of alcohol **6**

Entry	Conditions	Yield %
1	DMSO, (COCl) ₂ , Et ₃ N, CH ₂ Cl ₂	30%
2	DMSO, SO ₃ -pyridine, Et ₃ N, CH ₂ Cl ₂	22%
3	TPAP, NMO, CH ₂ Cl ₂	0%
4	MnO ₂ , acetone	0%
5	PCC, ether or PCC, CH ₂ Cl ₂	0%
6	Dess-Martin periodinate, CH ₂ Cl ₂	25%
7	BAIB, TEMPO, CH ₂ Cl ₂	54%