Published in final edited form as:

Curr Opin Allergy Clin Immunol. 2009 February ; 9(1): 29–37. doi:10.1097/ACI.0b013e32831f44a6.

Interactions between helminth parasites and allergy

Philip J Cooper

Centre for Infection, St George's University of London, London, UK

Instituto de Microbiologia, Universidad San Francisco de Quito, Quito, Ecuador

Centro de Investigaciones FEPIS, Hospital "Padre Alberto Buffoni", Quinindé, Esmeraldas Province, Ecuador

Abstract

Purpose of review: This article will review the findings of recent human studies of the association between helminth parasite infections and allergy and discuss their potential relevance to public health.

Recent findings: Different helminth parasites may have different effects on allergy that may depend on the timing of the exposure. Infections with *T. trichiura* in early life are associated with a reduced prevalence of allergen skin test reactivity later in life and infants of helminth-infected mothers have been reported to have a reduced prevalence of eczema. Hookworm infection has been associated with a reduced prevalence of asthma in Ethiopia. Several studies have reported that anti-*Ascaris* IgE is an important risk factor for asthma, but this could be explained by an enhanced ability of atopics to produce IgE. *Toxocara* infections may be associated with an increased risk of wheeze in some populations that may be caused by the host response to the parasite or by parasite-enhanced Th2 responses to aeroallergens.

Summary: Although helminth infections can modulate the host inflammatory response directed against the parasite, a causal association between helminths and atopic diseases remains uncertain.

Keywords

Helminths; Geohelminths; Allergy; Asthma

Introduction

There is a large and growing literature of the interaction between parasite infections and allergy derived from observations in humans and experimental animal models. This article will review the more recent findings from human studies of the possible relationship between respect parasite infections and allergy. The findings from experimental animal models have been reviewed extensively elsewhere [1,2]. The review will focus on the findings of the interaction between helminth parasites and allergy and will not discuss studies of protozoal parasite infections such as malaria [3] and giardiasis [4] for which there is very limited data.

The allergy epidemic

The prevalence of asthma and allergic diseases has increased in high-income countries over recent decades and may have reached a peak [5]. Allergic diseases are becoming an

important public health in many low and middle-income countries. Urban centres of Latin America appear to be most affected and have some of the highest reported prevalences of asthma worldwide [6,7]. The prevalence of asthma and allergic diseases appear to be low in many rural areas [8], an observation that has led to the suggestion that common environmental exposures present from an early age [9] in rural areas may be protective against allergy [10].

The hygiene hypothesis

A popular explanation for the increase in the prevalence of allergy is the hygiene hypothesis that attributes the allergy epidemic to a failure to develop appropriate immune regulation because of reduced exposures to microbes and their products in childhood [11]. There is considerable interest in the potential role of helminth infections in reducing allergy prevalence – certainly, helminth infections have strong regulatory effects, are highly prevalent, and first occur in early life in endemic areas.

Helminth parasites

The most common helminth infections are caused by geohelminth parasites (also known as intestinal and soil-transmitted helminths). Geohelminth parasites include *Ascaris lumbricoides*, *Trichuris trichiura*, and hookworm (*Ancylostoma duodenale* and *Necator americanus*) have a worldwide distribution, are estimated to infect a quarter of the World's population [12], and are most prevalent among children living in areas of the rural Tropics with poor access to sanitation and clean water. Other important helminth infections include schistosomiasis and filariasis have a more focal distribution within endemic countries.

Allergic inflammation directed against helminths

The human immune response to helminth infections is associated with elevated levels of IgE, tissue eosinophilia and mastocytosis, and the presence of CD4+ T cells that preferentially produce IL-4, IL-5, and IL-13 [1]. Th2-mediated mechanisms are considered to mediate protective immunity against these parasites [13]. Parasites in the tissues stimulate a strong localized Th2 response, characterized by an eosinophil-rich inflammatory infiltrate. A classic example is the Th2 granuloma that develops around schistosome eggs in the liver or wall of the intestine [14].

Individuals exposed to helminth infection may have allergic inflammatory responses to parasites and parasite antigens (Figure 1). Individuals with limited exposures to helminths such as expatriates or recent migrants often develop allergic-type clinical manifestations (Table 1) [15], a probable host response to isolate and kill the parasites (Figures 1B-D). A classic example is the asthma-like illness, Loeffler's syndrome, caused by the passage of *A. lumbricoides* larvae through the lungs. Helminth parasites in endemic areas tend to cause chronic infections - individual adult parasites may survive for many years in their human host - that are associated with few allergic-type reactions and a more tightly controlled Th2 response. Regulation of the Th2 response may be important for parasite survival and may allow the host to escape potentially damaging inflammation in the tissues.

For example, during infections with the tissue helminth, *Onchocerca volvulus*, the skin may be populated by millions of larval microfilariae and these appear to elicit little in the way of a host inflammatory response (Figure 2A). This state of hyporesponsiveness may be reversed rapidly after the killing of microfilariae by chemotherapy - treated individuals may develop allergic-type reactions (Figure 2C) that are associated with the development of eosinophilic abscesses in the superficial dermis (Figure 2B) within hours after treatment. The onset and severity of these reactions are associated with the release of allergic mediators

such as tryptase and eosinophil degranulation products into the peripheral circulation [16]. The hyporesponsiveness associated with chronic helminth infections appears to be actively regulated and may require the presence of live parasites.

Geohelminth parasites that are confined to the intestinal lumen may be less likely to induce strong systemic immune regulation although the tissue migratory life cycle stages of parasites such as *Ascaris lumbricoides* may induce strong allergic reactions in infected individuals living in regions where transmission of infection is seasonal. The comparative rarity of such reactions in endemic populations with year-round transmission [17] may reflect difficulties in diagnosis or perhaps suppression of the inflammatory response.

Many zoonotic helminth infections cannot develop to maturity in the human host and the helminth larvae may migrate for prolonged periods in the tissues (Table 1). Examples are infections with *Toxocara* spp, *Ascaris suum*, and dog hookworms. Such infections cause allergic type syndromes such as cutaneous (Figure 1B) and visceral larva migrans [18-20]. Tissue damage is caused by allergic inflammation directed against the migrating larvae. During such infections there appears to be a failure of immune regulation probably because host and parasite have not co-evolved.

Factors affecting the effects of helminths on allergy

Four factors may determine the effect of helminths on allergy: 1. *Timing* – the time of first infection and the duration of infection are likely to be important [21,22]. Early and/or long-lasting (chronic) infections may be more likely to induce immune modulatory effects that suppress allergic inflammation caused by parasite and non-parasite allergens while later and/or periodic infections may enhance allergy. The effect of geohelminths in suppressing atopy may be more important in the first years of life and the temporary elimination of infections later in childhood or adulthood may not affect a phenotype that is ‘programmed’ in infancy [21]. 2. *Intensity of infection* – heavy parasite burdens may induce immune down modulation while light infections may be more likely to have the opposite effect – the effects are likely to be stronger for tissue helminth infections than for geohelminth infections. 3. *Host genetics* – the ability to induce specific host immune regulatory mechanisms may be partly determined by host genetics. Individuals that are genetically susceptible to atopic disease may be more likely to develop allergic responses to helminth and non-parasite allergens and may be genetically more resistant to infection [23,24]. 4. *The parasite* – Different helminth parasites may have different effects on the risk of atopy and allergic disease [25].

Association of helminths with allergic diseases?

Helminth antigens stimulate allergic inflammatory responses directed against the parasite in the human host and that this inflammation may be actively suppressed during chronic infection. A distinct question is whether helminth infections may modulate also allergic inflammatory responses directed against non-parasite allergens such as aeroallergens and affect allergic sensitization and the expression of allergic diseases.

Helminths and atopy

Epidemiological studies have shown inverse associations between allergen skin test reactivity and infections with *A. lumbricoides* [26,27] and *T. trichiura* [22,26], hookworm [28], and schistosomiasis [29,30]. Both active and past infections appear to mediate this effect. Infections with *T. trichiura* in the first years of life are associated with a reduced prevalence of allergen skin test reactivity later in childhood independent of later infections [22]. A study of European farm children showed an inverse association between

sensitization to ascariasis (measured by the presence of specific IgG antibodies) and the presence of aeroallergen-specific IgE [31]. Not all studies, however, have shown an inverse association and some have provided evidence for positive associations between the presence of geohelminth infection [32] or *Ascaris*-specific IgE [27,33,34] and allergen skin test reactivity [27,32,34] or elevated allergen-specific IgE [33]. One study showed that allergen skin test reactivity was positively associated with anti-*Ascaris* IgE and negatively associated with active *A. lumbricoides* infection [27]. The possible relationship between atopy and *A. lumbricoides* infection is illustrated in Figure 3 [25-27, 33-45].

If helminth infections can actively suppress allergic inflammation, then anthelmintic treatment would be expected to reverse this effect. Several intervention studies have investigated this: 1) a non-randomized study of 94 children in Venezuela provided evidence that monthly anthelmintic treatment of children for 18 months caused an increase in the prevalence of skin test reactivity to house dust mite [46]. 2) a randomized placebo-controlled study of 165 children in Gabon showed that anthelmintic treatments every 3 months with treated the children with praziquantel and mebendazole every 3 months for 30-months was associated with an increase in the incidence of skin reactivity to house dust mite [47]. 3) a cluster-randomized study of 1,632 children in Ecuador did not show an effect on allergen skin test reactivity of anthelmintic treatment given every 2 months for 12 months [48]. The differences in the findings of these intervention studies may be explained by differences in the period of anthelmintic treatment, different parasites, and selection bias and uncontrolled confounding [21]. It would appear unlikely, however, that protection against allergen skin test reactivity is mediated by any single environmental exposure.

Helminths and asthma

A meta-analysis of many of studies investigating the association between the presence of geohelminth eggs in stool samples and asthma provided some evidence for parasite-specific effects [25]; *A. lumbricoides* eggs was associated with an increased prevalence of asthma, *T. trichiura* with no effect, and hookworm eggs with a reduced prevalence of asthma. All hookworm studies were conducted in Ethiopia and replication in other geographic regions is important.

Ascariasis may contribute to an increased risk of asthma either by causing directly inflammation in the airways (i.e. migrating larvae) or through increased atopy [34] and Th2 inflammatory responses in the airways. Studies investigating the association between the presence of anti-*Ascaris* IgE and asthma or bronchial hyperresponsiveness (BHR) have shown a strong positive association [34,41] that may be independent of endemicity and the presence of active infection (i.e. *A. lumbricoides* eggs in stool samples) [27,33,41]. Studies in urban Brazil have shown positive associations between *A. lumbricoides* infection and recent wheeze [43,49] and BHR [44]. The capacity to produce high levels of IgE on exposure to *A. lumbricoides* infection might simply be explained by a greater capacity to produce IgE on allergen exposure by (genetically susceptible) atopic children [50]. A study in Venezuela showed that the presence of atopy to *Ascaris* (elevated specific IgE and skin test reactivity) was an important risk factor for BHR in rural children but not in urban children in whom BHR was associated with atopy to house dust mite [42]. This could reflect a shift in the dominant allergen exposures to which children with atopic asthma are exposed [42]. The presence of anti-*Ascaris* IgE in asthmatics in Costa Rica has been associated with asthma severity and morbidity [34]. Monthly treatments of children with anthelmintic drugs in Venezuela may reduce BHR [46], symptoms of wheeze and the need for asthma medications [51]. The possible relationship between exposures to *A. lumbricoides* and asthma and BHR is illustrated in Figure 3.

An alternative explanation for the association between elevated *Ascaris*-specific IgE and asthma is infections with other common ascarid worms such as toxocariasis – antigen preparations from both worms show a high degree of immunological cross-reactivity. Toxocariasis is associated with asthma-like symptoms in children with visceral larva migrans [18] and there is some evidence that *Toxocara* may be an important risk factor for asthma in some populations [52,53]. Whether such asthma symptoms are caused directly by the parasite or by Th2 adjuvant effects of parasite antigens on responses to aeroallergens is not clear.

Other zoonotic infections associated with asthma and that may be an important risk factor in some populations or high-risk groups is *Anisakis simplex*. Anisakiasis is caused by the ingestion of live L3 larvae in inadequately cooked seafood or perhaps exposure to *Anisakis* proteins [54], and is considered to be an important cause of food allergy and BHR in Spain and Japan [55].

Helminths and eczema

Recent studies have shown both positive [56] and negative [45,57] associations between geohelminth infection and the prevalence of eczema [45,57]. A small intervention study in Uganda showed that infants of mothers with helminth infection at the time of delivery had a reduced risk of eczema compared to those born of uninfected mothers, and also a non-significant trend of a reduced risk of subsequent atopic dermatitis in the offspring of the mothers given anthelmintic treatment during pregnancy [58].

Immunological mechanisms of helminth-mediated modulation of allergy

Helminth parasites could affect allergic inflammation in three ways:

1) By enhancing or suppressing allergic inflammation directed against the parasite

Chronic helminth infections of humans suppress anti-parasite immune responses through regulatory immune cells such as regulatory T cells and alternatively activated macrophages and mechanisms that include the production of immune modulatory cytokines such as IL-10 and TGF- β [59].

2) Through immunological cross-reactivity between helminth allergens and aeroallergens

Important allergens such as tropomyosin of helminth parasites and invertebrates demonstrate immunological cross-reactivity [60]. A recent study of subjects infected with *A. lumbricoides* and asthmatics sensitized to American cockroach showed that although IgE antibodies from both groups were cross-reactive for American cockroach and *A. lumbricoides* tropomyosin, the cross-reactive IgE did not appear to be clinically relevant – none of the subjects with ascariasis had a positive skin test for American cockroach and none of the cockroach-sensitized asthmatic subject had ascariasis [61].

3) By affecting allergic inflammation directed against aeroallergens through bystander effects in the same tissues such as the lungs

Helminth infections may contribute to ‘immune homeostasis’. Early life exposures in particular could have important long-term effects [21]. Immune homeostatic mechanisms affected might include the production of baseline levels of regulatory cytokines (e.g. IL-10) by immune cells in the tissues that could raise the thresholds for the induction of effector cell responses to aeroallergens. A study of children infected in Cameroon provided some evidence for elevated production of IL-10 and TGF- β 1 by unstimulated peripheral blood leukocytes (PBLs) that was associated positively with geohelminth parasite burden and inversely with for immune reactivity [62]. Regulatory homeostasis may be expected to be

insensitive to short term fluctuations in parasite burdens (e.g single dose anthelmintic treatment) but could be 'reset' by long-term changes in parasite levels (e.g. increases in parasite burdens or repeated anthelmintic treatments). Single anthelmintic treatments for geohelminth infections do not affect cytokine responses to parasite antigen [63,64] but repeated doses over prolonged periods caused an increase in Th2 cytokine responses of PBLs from Ecuadorian children [65]. Long-term anthelmintic treatment did not affect cytokine responses to aeroallergens indicating that the suppressive effect was specific for antiparasite responses [65].

Bystander suppression may be also a more active process that requires the continued presence of the parasite. Active immune regulation may be mediated through mechanisms such as the direct suppressive effects of parasite-secretions and parasite-specific regulatory immune cells in the tissues. Such suppression may be reversed rapidly after anthelmintic treatment or parasite death and appears to be an important survival mechanism for tissue helminth parasites. Such suppression may affect also immunity to aeroallergens and could explain the observation of an inverse association between the production of parasite-antigen induced IL-10 by PBLs and skin test reactivity to house dust mite among children from Gabon living in an endemic area *S. mansoni* [30]. Further, a study in Brazil that compared asthmatics infected with *S. mansoni* and those not infected (from a different population) showed that *D. pteronyssinus*-stimulated PBLs from infected asthmatics produced less Th2 cytokines and more IL-10 compared to controls [66]. However, studies conducted in areas where *A. lumbricoides* is the predominant helminth have not provided evidence for either enhanced IL-10 responses to aeroallergens [67,68] or an increase in the frequency of regulatory T cell populations induced by aeroallergen stimulation of PBLs [68].

Public health importance of parasite-allergy interactions

Helminth infections cause significant morbidity in endemic countries and treatment programmes for a number of these diseases are being implemented as a public health priority. Helminth parasites are presumed to be protective against allergic diseases [1,21] and there is some concern that treatment programmes, by controlling helminth parasites [48], may increase the prevalence of allergy among rural populations where atopic disease are considered to be relatively rare [8]. However, there is no compelling evidence that helminths protect against atopic diseases or that the treatment of endemic populations increases the prevalence of these [21,46,48]. Anthelmintic treatment may increase the prevalence of allergen skin test reactivity in some studies [46,47], but the public health relevance of this is not clear – only a small proportion (~11%) of asthma may be attributable to allergen skin test reactivity in some areas of Latin America [7, 69]. Further, ascariasis and toxocariasis may be important risk factors for asthma in some populations [34,41,43,44,46,52,53], and therefore, anthelmintic treatment programmes may have beneficial effects. Early exposures to parasites such as geohelminth infections may contribute to programming of immune homeostasis and the long-term effects of anthelmintic treatment programmes in endemic populations on public health (i.e. inflammatory diseases) remains to be established. There is some evidence that hookworm infection may protect against asthma in Ethiopia [25] and clinical studies of experimental infections with hookworm in individuals with atopic diseases are in progress [70]. Helminth infections produce powerful modulators of the host inflammatory response that protect them against killing or expulsion by the host. The identification of these factors and an understanding of the mechanisms by which they work may lead to the development of novel anti-inflammatory treatments.

Conclusion

Helminth infections have strong modulatory effects on anti-parasite inflammatory responses in the human host but it is not clear if helminths can affect allergic inflammatory responses to aeroallergens. Helminth infections have been associated with both a reduced prevalence and increased prevalence of atopy and atopic disease in different populations. The immune regulatory effects of tissue helminths are likely to be stronger than those of geohelminths. Further research in prospective observational and intervention studies is required to address the question of causality. An understanding of the mechanisms by which helminth parasites modulate the host allergic inflammatory response may lead to the development of novel anti-inflammatory interventions. The demonstration of a causal association between some helminth parasites (particularly geohelminths and toxocariasis that have a worldwide distribution) and an increased risk of asthma could lead to anthelmintic treatment programmes in populations considered to be at high risk.

References

1. Fallon PG, Mangan NE. Suppression of Th2-type allergic reactions by helminth infections. *Nature Rev Immunol.* 2007; 7:220–230. [PubMed: 17318233] [Excellent review of immunology of parasite-allergy interactions in experimental animal models]
2. van Riet E, Hartgers FC, Yazdanbakhsh M. Chronic helminth infections induce immunomodulation: consequences and mechanisms. *Immunobiol.* 2007; 212:475–490.
3. van den Bigelaar AHJ, Lopuhaa C, van Ree R, et al. The prevalence of parasite infestation and house dust mite sensitisation in Gabonese schoolchildren. *Int Arch Allergy Immunol.* 2001; 126:231–238. [PubMed: 11752881]
4. Bayraktar MR, Mehmet N, Durmaz R. Serum cytokine changes in Turkish children infected with *Giardia lamblia* with and without allergy: Effect of metronidazole treatment. *Acta Trop.* 2005; 95:116–122. [PubMed: 15979046]
5. Asher MI, Montefort S, Björkstén B, et al. Worldwide time trends in the prevalence of symptoms of asthma, allergic rhinoconjunctivitis, and eczema in childhood: ISAAC Phases One and Three repeat multicountry cross-sectional surveys. *Lancet.* 2006; 368:733–743. [PubMed: 16935684]
6. Pearce N, Ait-Khaled N, Beasley R, et al. Worldwide trends in the prevalence of asthma symptoms: phase III of the International Study of Asthma and Allergies in Childhood (ISAAC). *Thorax.* 2007; 62:758–766. [PubMed: 17504817]
7. Cooper PJ, Rodrigues LC, Cruz AA, Barreto ML. Asthma in Latin America: a public health challenge and research opportunity. *Allergy.* 2008 in press.
8. Nicolaou N, Siddique N, Custovic A. Allergic disease in urban and rural populations: increasing prevalence with increasing urbanization. *Allergy.* 2005; 60:1357–1360. [PubMed: 16197466]
9. Ege MJ, Herzum I, Büchele G, et al. Prenatal exposure to a farm environment modifies atopic sensitization at birth. *J Allergy Clin Immunol.* 2008; 122:407–412. [PubMed: 18678343]
10. von Mutius E. Asthma and allergies in rural areas of Europe. *Proc Am Thorac Soc.* 2007; 4:212–216. [PubMed: 17607001]
10. Ege MJ, Herzum I, Büchele G, et al. Prenatal exposure to a farm environment modifies atopic sensitization at birth. *J Allergy Clin Immunol.* 2008; 122:407–412. [PubMed: 18678343]
11. Garn H, Renz H. Epidemiological and immunological evidence for the hygiene hypothesis. *Immunobiology.* 2007; 212:441–452. [PubMed: 17544829]
12. Bethony J, Brooker S, Albonico M, et al. Soil-transmitted helminth infections: ascariasis, trichuriasis, and hookworm. *Lancet.* 2006; 367:1521–1532. [PubMed: 16679166]
13. Anthony RM, Rutitzky LI, Urban JF Jr, et al. Protective immune mechanisms in helminth infection. *Nat Rev Immunol.* 2007; 7:975–987. [PubMed: 18007680]
14. Wilson MS, Mentink-Kane MM, Pesce JT, et al. Immunopathology of schistosomiasis. *Immunol Cell Biol.* 2007; 85:148–154. [PubMed: 17160074]

15. Cooper, PJ.; Nutman, TB. IgE and its role in parasitic helminth infection: implications for anti-IgE based therapies. In: Fick, RB.; Jardieu, P., editors. *IgE and anti-IgE therapy in asthma and allergic disease, Lung Biology in Health and Disease*. Vol. 164. New York: Marcel Dekker; 2002. p. 409-425.
16. Cooper PJ, Schwartz L, Awadzi K, et al. Plasma tryptase, a marker of mast cell degranulation, is associated with the development of adverse reactions following the treatment of onchocerciasis with ivermectin. *J Infect Dis*. 2002; 186:1307–1313. [PubMed: 12402200]
17. Spillmann RK. Pulmonary ascariasis in tropical communities. *Am J Trop Med Hyg*. 1975; 24:791–800. [PubMed: 1238026]
18. Nash, TE. Visceral larva migrans and other unusual helminth infections. In: Mandell, GR.; Bennett, JE.; Dolin, R., editors. *Principles and Practice of Infectious Diseases*. 6th Edition. Philadelphia: Elsevier Churchill Livingstone; 2005. p. 3293-99.
19. Maruyama H, Nawa Y, Noda S, et al. An outbreak of visceral larva migrans due to *Ascaris suum* in Kyushu Japan. *Lancet*. 1996; 347:1766–1767. [PubMed: 8656926]
20. Heukelbach J, Feldmeier H. Epidemiological and clinical characteristics of hookworm-related cutaneous larva migrans. *Lancet Infect Dis*. 2008; 8:302–309. [PubMed: 18471775]
21. Cooper PJ, Barreto M, Rodrigues LC. Human allergy and intestinal helminth infections: a review of the literature and discussion of a conceptual model to investigate the possible causal association. *Br Med Bull*. 2006; 79-80:203–218. [PubMed: 17204485]
22. Rodrigues LC, Newland P, Cunha SS, et al. Early infections with intestinal helminths reduce the risk of atopy later in childhood. *Clin Exp Allergy*. Jun 10.2008 [First paper showing that early geohelminth infections may affect the later development of allergen skin test reactivity]
23. Peisong G, Mao X-Q, Enomoto T, et al. An asthma-associated genetic variant of STAT6 predicts low burden of ascaris worm infestation. *Genes Immun*. 2005; 5:58–62. [PubMed: 14735150]
24. Moller M, Gravenor MB, Roberts SE, et al. Genetic haplotypes of Th-2 immune signaling link allergy to enhanced protection to parasitic worms. *Hum Mol Gen*. 2007; 16:1828–1836. [PubMed: 17519224]
25. Leonardi-Bee J, Pritchard D, Britton J. Asthma and current intestinal parasite infection: systematic review and meta-analysis. *Am J Respir Crit Care Med*. 2006; 174:514–523. [PubMed: 16778161]
26. Cooper PJ, Chico ME, Rodrigues LC, et al. Reduced risk of atopy among school age children infected with geohelminth parasites in a rural area of the tropics. *J Allergy Clin Immunol*. 2003; 111:995–1000. [PubMed: 12743563]
27. Obihara CC, Beyers N, Gie RP, et al. Respiratory atopic disease, Ascaris-immunoglobulin E and tuberculin testing in urban South African children. *Clin Exp Allergy*. 2006; 36:640–648. [PubMed: 16650050]
28. Flohr C, Tuyen LN, Lewis S, et al. Poor sanitation and helminth infection protect against skin sensitization in Vietnamese children: a cross-sectional study. *J Allergy Clin Immunol*. 2006; 118:1305–1311. [PubMed: 17157661]
29. Araujo MI, de Carvalho EM. Human schistosomiasis decreases immune responses to allergens and clinical manifestations of asthma. *Chem Immunol Allergy*. 2006; 90:29–44. [PubMed: 16210901]
30. van den Biggelaar AH, van Ree R, Rodrigues LC, et al. Decreased atopy in children infected with *Schistosoma haematobium*: a role for parasite-induced interleukin-10. *Lancet*. 2000; 356:1723–1727. [PubMed: 11095260]
31. Karadag N, Ege M, Bradley JE, et al. The role of parasitic infections in atopic diseases in rural schoolchildren. *Allergy*. 2006; 61:996–1001. [PubMed: 16867054]
32. Palmer LJ, Celedon JC, Weiss ST, et al. *Ascaris lumbricoides* infection is associated with increased risk of childhood asthma and atopy in rural China. *Am J Respir Crit Care Med*. 2002; 165:1489–1493. [PubMed: 12045121]
33. Dold S, Heinrich J, Wichmann HE, Wjst M. *Ascaris*-specific IgE and allergic sensitization in a cohort of school children in the former East Germany. *J Allergy Clin Immunol*. 1998; 102:414–420. [PubMed: 9768582]
34. Hunninghake GM, Soto-Quiros ME, Avila L, et al. Sensitization to *Ascaris lumbricoides* and severity of childhood asthma in Costa Rica. *J Allergy Clin Immunol*. 2007; 119:654–661. [PubMed: 17336615]

35. Cooper PJ, Chico ME, Sandoval C, et al. Human infection with *Ascaris lumbricoides* is associated with a polarized cytokine response. *J Infect Dis.* 2000; 182:1207–1213. [PubMed: 10979919]
36. Cooper PJ, Chico ME, Griffin GE, Nutman TB. Allergy symptoms, atopy, and geohelminth infections in a rural area of Ecuador. *Am J Resp Crit Care Med.* 2003; 168:313–317. [PubMed: 12714349]
37. Dagoye D, Bekele Z, Woldemichael K, et al. Wheezing, allergy and parasite infection in children in urban and rural Ethiopia. *Am J Resp Crit Care Med.* 2003; 167:1369–1373. [PubMed: 12738598]
38. Davey G, Venn A, Belete H, et al. Wheeze, allergic sensitization and geohelminth infection in Butajira, Ethiopia. *Clin Exp Allergy.* 2005; 35:301–307. [PubMed: 15784107]
39. Turner JD, Faulkner H, Kamgno J, et al. Allergen-specific IgE and IgG4 are markers of resistance and susceptibility in a human intestinal nematode infection. *Microbes Infect.* 2005; 7:990–996. [PubMed: 15961339]
40. Steinman HA, Donson H, Kawalski M, et al. Bronchial hyperresponsiveness and atopy in urban, peri-urban, and rural South African children. *Pediatr Allergy Immunol.* 2003; 14:383–393. [PubMed: 14641609]
41. Takeuchi H, Zaman K, Takahashi J, et al. High titre of anti-*Ascaris* immunoglobulin E associated with bronchial asthma symptoms in 5-year-old rural Bangladeshi children. *Clin Exp Allergy.* 2008; 38:276–282. [PubMed: 18070165]
42. Hagel I, Cabrera M, Hurtado MA, et al. Infection by *Ascaris lumbricoides* and bronchial hyper reactivity: an outstanding association in Venezuelan school children from endemic areas. *Acta Trop.* 2007; 103:231–241. [PubMed: 17698018]
43. Pereira MU, Sly PD, Pitrez PM, et al. Nonatopic asthma is associated with helminth infections and bronchiolitis in poor children. *Eur Respir J.* 2007; 29:1154–1160. [PubMed: 17331964]
44. da Silva ER, Sly PD, de Pereira MU, et al. Intestinal helminth infestation is associated with increased bronchial responsiveness in children. *Pediatr Pulmonol.* 2008; 43:662–665. [PubMed: 18484663]
45. Wordemann M, Diaz RJ, Heredia LM, et al. Association of atopy, asthma, atopic dermatitis and intestinal helminth infections in Cuban children. *Trop Med Int Health.* 2008; 13:180–186. [PubMed: 18304263]
46. Lynch NR, Hagel I, Perez M, et al. Effect of anthelmintic treatment on the allergic reactivity of children in a tropical slum. *J Allergy Clin Immunol.* 1993; 92:404–411. [PubMed: 8360391]
47. van den Biggelaar AHJ, Rodrigues LC, van Ree R, et al. Long-term treatment of intestinal helminths increases mite skin-test reactivity in Gabonese schoolchildren. *J Infect Dis.* 2004; 189:892–900. [PubMed: 14976607]
48. Cooper PJ, Chico ME, Vaca M, et al. Impact of bimonthly treatment of geohelminth-infected children with albendazole on atopy prevalence: a cluster-randomized trial. *Lancet.* 2006; 367:1598–1603. [PubMed: 16698413]
49. Camara AA, Silva JM, Ferriani VPL, et al. Risk factors for wheezing in a subtropical environment: role of respiratory viruses and allergen sensitization. *J Allergy Clin Immunol.* 2004; 113:551–557. [PubMed: 15007360]
50. Lynch NR, Hagel IA, Palenque ME, et al. Relationship between helminthic infection and IgE response in atopic and nonatopic children in a tropical environment. *J Allergy Clin Immunol.* 1998; 101:217–221. [PubMed: 9500755]
51. Lynch NR, Palenque M, Hagel I, DiPrisco MC. Clinical improvement of asthma after anthelmintic treatment in a tropical situation. *Am J Resp Crit Care Med.* 1997; 156:50–54. [PubMed: 9230725]
52. Cooper PJ. *Toxocara canis* infection: an important and neglected risk factor for asthma? *Clin Exp Allergy.* 2008; 38:551–553. [PubMed: 18241245]
53. Ferreira MU, Rubinsky-Elefant G, Gontijo de Castro T, et al. Bottle feeding and exposure to *Toxocara* as risk factors for wheezing illness among under-five Amazonian children: a population-based cross-sectional study. *J Trop Ped.* 2007; 53:119–124.

54. Nieuwenhuizen N, Lopata AL, Jeebhay MF, et al. Exposure to the fish parasite *Anisakis* causes allergic airway hyperreactivity and dermatitis. *J Allergy Clin Immunol.* 2006; 117:1098–1105. [PubMed: 16675338]
55. Audicana MT, Kennedy MW. *Anisakis* simplex: from obscure infectious worm to inducer of immune hypersensitivity. *Clin Microbiol Rev.* 2008; 21:360–379. [PubMed: 18400801]
56. Haileamlak A, Dagoye D, Williams H, et al. Early life risk factors for atopic dermatitis in Ethiopian children. *J Allergy Clin Immunol.* 2005; 115:370–376. [PubMed: 15696097]
57. Schafer T, Meyer T, Ring J, et al. Worm infestation and the negative association with eczema (atopic/nonatopic) and allergic sensitization. *Allergy.* 2005; 60:1014–1020. [PubMed: 15969681]
58. Elliott AM, Mpairwe H, Quigley MA, et al. Helminth infection during pregnancy and development of infantile eczema. *JAMA.* 2005; 294:2032–2034. [PubMed: 16249415]
59. Hoerauf A, Satoguina J, Saeftel M, Specht S. Immunomodulation by filarial nematodes. *Parasite Immunol.* 2005; 27:417–29. [PubMed: 16179035]
60. Arruda LK, Santos AB. Immunologic responses to common antigens in helminthic infections and allergic disease. *Curr Opin Allergy Clin Immunol.* 2005; 5:399–402. [PubMed: 16131913]
61. Santos ABR, Rocha GM, Oliver C, et al. Cross-reactive IgE antibody responses to tropomyosins from *Ascaris lumbricoides* and cockroach. *J Allergy Clin Immunol.* 2008; 121:1040–1046. [PubMed: 18275995] [Paper showing immunological cross-reactivity between *A. lumbricoides* and American cockroach tropomyosin may not be clinically important]
62. Turner JD, Jackson JA, Faulkner H, et al. Intensity of intestinal infection with multiple worm species is related to regulatory cytokine output and immune hyporesponsiveness. *J Infect Dis.* 2008; 197:1204–1212. [PubMed: 18462167] [Paper provides some evidence for the importance of geohelminth infections in immune regulation]
63. Cooper PJ, Chico M, Sandoval C, et al. Human infection with *Ascaris lumbricoides* is associated with suppression of the interleukin-2 response to recombinant cholera toxin B subunit following vaccination with the live oral cholera vaccine CVD 103-HgR. *Infect Immun.* 2001; 69:1574–1580. [PubMed: 11179329]
64. Geiger SM, Massara CL, Bethony J, et al. Cellular responses and cytokine production in post-treatment hookworm patients from an endemic area in Brazil. *Clin Exp Immunol.* 2004; 136:334–340. [PubMed: 15086399]
65. Cooper PJ, Mitre E, Moncayo AL, et al. *Ascaris lumbricoides*-induced interleukin-10 is not associated with atopy in schoolchildren in a rural area of the tropics. *J Infect Dis.* 2008; 197:1333–1340. [PubMed: 18422446]
66. Araujo MI, Hoppe B, Medeiros M Jr, et al. Impaired T helper 2 response to aeroallergen in helminth-infected patients with asthma. *J Infect Dis.* 2004; 190:1797–1803. [PubMed: 15499536]
67. Ponte EV, Lima F, Araujo MI, et al. Skin test reactivity and Der p-induced interleukin 10 production in patients with asthma or rhinitis infected with *Ascaris*. *Ann Allergy Asthma Immunol.* 2006; 96:713–718.
68. Cooper PJ, Moncayo AL, Guadalupe I, et al. Repeated treatments with albendazole enhance Th2 responses to *Ascaris lumbricoides*, but not to aeroallergens, in children from rural communities in the Tropics. *J Infect Dis.* Aug 26, 2008
69. Weinmayr G, Weiland SK, Björkstén B, et al. Atopic sensitization and the international variation of asthma symptom prevalence in children. *Am J Respir Crit Care Med.* 2007; 176:565–574. [PubMed: 17575099] [Important paper from ISAAC phase II studies showing international variations in the association between atopy and asthma]
70. Falcone FH, Pritchard DI. Parasite role reversal: worms on trial. *Trends Parasitol.* 2005; 21:157–160. [PubMed: 15780835]

Figure 1.

Examples of allergic-type reactions to helminth parasites. A. Immediate hypersensitivity reaction to *Paragonimus mexicanus* antigen extract injected into the forearm of child. B. Cutaneous larva migrans showing serpiginous track of dog hookworm larvae in foot. C. Punctate keratitis in young child infected with *Onchocerca volvulus*. The punctate opacities are sites of microfilarial death in the cornea and are composed of eosinophil-rich inflammatory infiltrates. The photograph was taken after removal of an onchocercal nodule (bandage). D. Acute papular onchodermatitis in woman infected with *O. volvulus*. Localized popular eruptions appear periodically in infected individuals representing the sites of 'spontaneous' microfilarial death.

Figure 2.

Allergic-type inflammatory reactions to *Onchocerca volvulus* microfilariae in the skin. The Figure shows effect of treatment with the microfilaricidal drug diethylcarbamazine. Pre-treatment skin biopsy (A) shows microfilariae in the dermis with few associated inflammatory cells. After treatment with a single dose of DEC, the dead microfilariae become the focus of an intense inflammatory reaction composed of eosinophilic abscesses in the superficial dermis (B). Clinical appearance of the skin after DEC treatment (C).

Figure 3.

Possible effects of *Ascaris lumbricoides* infection on atopy and asthma. The human response to exposure to *A. lumbricoides* is likely to be modified by host genetic factors and other factors including the intensity of infection and the age at which first infections occur. Exposure to *A. lumbricoides* could affect the development of allergen skin test reactivity in 4 ways (shown by paths 1-2-3, 1-2-4-7 and 1-4). Pathway 9-10 represents the effect of chronic infections - a high IgG4/low IgE response to *A. lumbricoides* or modified Th2 response [reference 1] develops and may lead to the suppression of allergen skin test reactivity. The pathways leading from step1 would be associated with a high IgE/low IgG4 response to *A. lumbricoides*. Pathway 7 represents the enhancement of allergen skin test reactivity by an unknown mechanism. *Exposure to A. lumbricoides could cause or increase asthma symptoms and bronchial hyper-reactivity* via enhanced allergen skin test reactivity (paths 7-6, 1-2-3-6, and 1-4-6) or be independent of allergen skin test reactivity (paths 1-5 and 8) or independent of both anti-*Ascaris* IgE and allergen skin test reactivity (8). Suppression of asthma symptoms and or BHR could occur via path 9-11, although there is very limited evidence in support of this. Shown are published associations from cross-sectional studies and presumed causal directions (arrows). Reference numbers are shown in brackets.

Table 1

Allergic-type reactions associated with human helminth parasites and possible associations between helminth infections and atopic diseases.

Helminth infection	Examples of allergic-type reactions and syndromes	Treatment reactions*	Atopic diseases
Intestinal helminths			
<i>Ascaris lumbricoides</i>	'Asthma-like' syndrome	No	Asthma
<i>Trichiura trichiura</i>	Tropical dysentery syndrome	No	No
Hookworm	Ground itch/allergic enteritis	No	Decreased asthma?
<i>Strongyloides stercoralis</i>	Larva currens/urticaria/'asthma-like' syndrome	No	No
<i>Enterobius vermicularis</i>	Itchy bum	No	Decreased allergic rhinitis?
Schistosomiasis			
<i>S. mansoni</i>	Cercarial dermatitis/acute schistosomiasis/urticaria/'asthma-like' syndrome	Yes	Milder asthma?
<i>S. haematobium</i>		Yes	No
<i>S. japonicum</i>		Yes	No
Filariasis			
<i>Wuchereria bancrofti</i>	Tropical pulmonary eosinophilia/acute lymphangitis	Yes	No
<i>Onchocerca volvulus</i>	Sowda/acute popular onchodermatitis/punctate keratitis	Yes	No
<i>Loa loa</i>	Calabar swellings	Yes	No
Others			
<i>Toxocara</i> spp.	Visceral larva migrans/'asthma-like' syndrome	No	Asthma
<i>Anisakis</i> spp.	'gastroallergic'/asthma-like syndrome/urticaria/anaphylaxis	No	Asthma
<i>Paragonimus</i> spp.	Asthma-like syndrome	No	No
<i>Trichinella spiralis</i>	Acute trichinosis	No	No
<i>Echinococcus granulosus</i>	Acute anaphylaxis associated with rupture of cyst	No	No
<i>Ancylostoma braziliense</i>	Cutaneous larva migrans	No	No

* Treatment reactions are more commonly observed for tissue parasites associated with very high parasite burdens.