CORRESPONDENCE

Fulminant Unilateral Pulmonary Edema After Insertion of a Chest Tube: a Complication After a Primary Spontaneous Pneumothorax

by Nina Schmidt-Horlohé, Dr. med. Chadwick T. Azvedo,
Priv.-Doz. Dr. med. Lothar Rudig, Dr. med. Michael Habekost in volume 50/2008

Undiagnosed Takotsubo Cardiomyopathy?

It is remarkable and difficult to explain in terms of pathophysiology that in an otherwise healthy, young individual with unilateral pulmonary impairment, pulmonary edema required intubation and administration of catecholamines. Increasingly often, even in pneumology patients (1), instances have been reported of transient, stress associated cardiomyopathy with complete reversibility (takotsubo cardiomyopathy). The descriptions of both pathologies have shown parallels: the rapid manifestation after an event/intervention, the short duration of the impairment, and the generally good prognosis. The clientele that is affected differs; takotsubo has been described more often for women beyond their reproductive years. In the radiographs of the acute episode that were reproduced with the article, the transverse cardiac diameter appears widened on the account of the left ventricle; however, it has to be borne in mind that the images were taken with the patient in a supine position. Were you perhaps able to undertake echocardiography in the acute situation?

The unilateral development of pulmonary edema during the crisis could actually be explained with a reduction of the surfactant in the affected lung, even if the complaint did have a cardiac cause.

Perhaps the differential diagnosis of takotsubo cardiomyopathy might be followed in the scenario of a re-expansion edema, and an echocardiogram could be included in the diagnostic work-up for this complication.

DOI: 10.3238/arztebl.2009.0223a

REFERENCES

1. Cardiovascular collapse after therapeutic bronchoscopy. Ernst A, Brookline, USA at the ERS Annual Congress, Berlin 2008.
2. Schmidt-Horlohé N, Azvedo CT, Rudig L et al.: Case report: Fulminant unilateral pulmonary edema after insertion of a chest tube: a complication after a primary spontaneous pneumothorax. *Dtsch Arztebl Int* 2008; 105(50): 878–81.

Dr. med. Barbara Khanavkar
Krankenhaus Salzwedel
Brunnenstr. 1, 29410 Salzwedel, Germany
B.Khanavkar@altmark-klinikum.de

In Reply:

The fact that an otherwise healthy, young patient's cardiopulmonary system decompensates in such a manner owing to unilateral impairment is indeed surprising. The literature cited with our article, however, reports

several such instances. The pathophysiological examination remains unclear—the shift of larger volumes of intravascular fluids into the thorax (interstitial and intra-alveolar) has been discussed as a possible mechanism (1). The need for catecholamines in the patient in our case report existed for a short period of time only, before sedation was stepped down, and the required dosage was low.

The suggested differential diagnosis of stress associated takotsubo cardiomyopathy is an interesting one. Unfortunately, we cannot refer to valid parameters in the sense of valid diagnostic tests as neither electrocardiography nor echocardiography were undertaken, nor were the respective laboratory results available. Repeated work-up of the documentation, however, does not hint at such an entity. The clinical symptoms described for takotsubo cardiomyopathy, similar to those of cardiac ischemia, were not present before intubation, after extubation, and retrospectively in the patient's further inpatient and outpatient progress (2). Isolated unilateral pulmonary edema as a result of a transient cardiac pump failure owing to the "myocardial stunning" described in the context of this disorder seems to us rather untypical. Also, female patients of an older age are regarded as having a higher risk (3). Without objective findings, takotsubo cardiomyopathy can certainly not be excluded, but to us it seems rather unlikely. If any clinical signs are present then this differential diagnosis should probably be excluded by means of suitable diagnostic tests.

DOI: 10.3238/arztebl.2009.0223b

REFERENCES

1. Pavlin DJ, Raghu G, Rogers TR et al.: Reexpansion hypotension. A complication of rapid evacuation of prolonged pneumothorax. *Chest* 1986; 89: 70–4.
2. von Korn H, Yu J, Lotze U et al.: Tako-Tsubo-like cardiomyopathy: specific ECG findings, characterization and clinical findings in an European single center. *Cardiology* 2009; 112: 42–8.
3. Vizzardi E, D'Aloia A, Zanini G et al.: Tako-tsubo-like left ventricular dysfunction: transient left ventricular apical ballooning syndrome. *Int J Clin Pract* 2008 (published online).
4. Schmidt-Horlohé N, Azvedo CT, Rudig L et al.: Case report: Fulminant unilateral pulmonary edema after insertion of a chest tube: a complication after a primary spontaneous pneumothorax. *Dtsch Arztebl Int* 2008; 105(50): 878–81.

Nina Schmidt-Horlohé
Klinik für Unfall-, Hand- und Wiederherstellungschirurgie
GPR Klinikum Rüsselsheim, August-Bebel-Str. 59, 65428 Rüsselsheim, Germany
schmidt-horlohe@gp-ruesselsheim.de

Conflict of interest statement

The authors of both letters declare that no conflict of interest exists according to the guidelines of the International Committee of Medical Journal Editors.