Published in final edited form as:

Calcif Tissue Int. 2009 February ; 84(2): 75–84. doi:10.1007/s00223-008-9197-3.

Quantitative trait locus on chromosome 1q influences bone loss in young Mexican American adults

John R. Shaffer¹, Candace M. Kammerer¹, Jan M. Bruder², Shelley A. Cole³, Thomas D. Dyer³, Laura Almasy³, Jean W. MacCluer³, John Blangero³, Richard L. Bauer², and Braxton D. Mitchell⁴

¹Department of Human Genetics, University of Pittsburgh Graduate School of Public Health, Pittsburgh, PA, USA

²University of Texas Health Science Center, San Antonio, TX, USA

³Southwest Foundation for Biomedical Research, San Antonio, TX, USA

⁴Division of Endocrinology, Diabetes and Nutrition, University of Maryland School of Medicine, Baltimore, MD, USA

Abstract

Introduction—Bone loss occurs as early as the third decade and its cumulative effect throughout adulthood may impact risk for osteoporosis in later life, however the genes and environmental factors influencing early bone loss are largely unknown. We investigated the role of genes in the change in bone mineral density (BMD) in participants of the San Antonio Family Osteoporosis Study.

Materials and Methods—BMD change in 327 Mexican Americans (ages 25–45 years) from 32 extended pedigrees was calculated from DXA measurements at baseline and follow-up (3.5 to 8.9 years later). Family-based likelihood methods were used to estimate heritability (h^2) and perform autosome-wide linkage analysis for BMD change of the proximal femur and forearm, and estimate heritability for BMD change of lumbar spine.

Results—BMD change was significantly heritable for total hip, ultradistal radius and 33% radius ($h^2 = 0.34, 0.34, 0.27$, respectively, $p < 0.03$ for all), modestly heritable for femoral neck ($h^2 = 0.22$, $p = 0.06$) and not heritable for spine BMD. Covariates associated with BMD change included age, sex, baseline BMD, menopause, body mass index, and interim BMI change, and accounted for 6% to 24% of phenotype variation. A significant quantitative trait locus (LOD = 3.6) for femoral neck BMD change was observed on chromosome 1q23.

Conclusions—We observed that change in BMD in young adults is heritable, and performed one of the first linkage studies for BMD change. Linkage to chromosome 1q23 suggests this region may harbor one or more genes involved in regulating early BMD change of the femoral neck.

Keywords

Bone Mineral Density; Bone Loss; Genetics; Linkage; Heritability

Introduction

Peak bone mass and rates of bone loss following menopause in women and later in life in men are commonly acknowledged to be important risk factors for osteoporosis. In fact, Hui et al. suggest that the contributions of peak bone mass acquisition and bone loss after menopause are roughly equal in determining bone mass in women 70 years of age [1]. Over the past three decades, some of the factors that influence peak bone mass in adults, as well as bone loss among older men and women, have been identified [2–8]. In addition to environmental factors, numerous studies have documented that peak bone mineral density (BMD) is highly heritable [9–14] and many quantitative trait loci (QTLs, i.e. implicated chromosomal regions potentially harboring regulatory genes) have been reported [15–34]. More recently, studies have clearly demonstrated that genetic factors regulate BMD change at several skeletal sites in women and older men. Specifically, the heritability of change in BMD at the femoral neck, in premenopausal women [35], lumbar spine, forearm, and whole body (but not hip) in peri- and postmenopausal women [36], and lumbar spine, hip, and forearm (but not whole body) in older men and women (>45 years) [37], has been demonstrated. Studies to identify specific genes influencing peak bone mass and bone loss in older individuals are ongoing.

Although many studies have investigated genetic and environmental factors influencing peak bone mass and bone loss among older individuals, few studies have looked at the factors affecting early change in BMD among young adults, particularly the role of genetics in bone loss [9,38]. While most cross-sectional studies project that minimal bone loss occurs among individuals <50 years of age [39], more recent longitudinal studies have revealed significant rates of bone loss occurring as early as the third decade in women [35,40–43] and men [43, 44]. Furthermore, results from cross-sectional and longitudinal studies indicate that the onset of trabecular bone loss may begin in early adulthood, whereas the onset of cortical bone loss is delayed until mid-life in women and later in life in men [10,43]. Thus, bone health later in life may depend upon early bone loss, in addition to peak bone mass and advanced age-related bone loss. Furthermore, the importance of early bone loss with regard to later bone health may depend on the trabecular and cortical content of a particular skeletal site. One study has reported that bone loss of the femoral neck is heritable in premenopausal Caucasian women [35], but to date, no study has assessed the genetic contribution to early bone loss at other skeletal sites or in men (younger than 45 years of age). Furthermore, no specific QTLs for longitudinal BMD change have been identified in any population.

As part of the San Antonio Family Osteoporosis Study, we sought to investigate the role of genetic factors on bone loss in younger individuals at a number of skeletal sites. Specifically, we have estimated the heritability of 5-year change in BMD in 327 Mexican American men and women (ages 25 to 45) from large multigenerational kinships, and performed autosomal-wide linkage analysis in search of QTLs influencing common variation in early BMD change.

Experimental Subjects

Recruitment and data collection for the baseline and follow-up phases of the San Antonio Family Osteoporosis Study have been fully described previously [11,37]. In brief, participants from 34 multigenerational families were recruited from a low-income neighborhood via a house-to-house recruitment protocol. Probands meeting eligibility criteria (i.e. aged 40 to 60 years and having large families in the San Antonio area) and all first, second, and third degree relatives and spouses were invited to participate irrespective of current health outcomes. Participating families represent a fundamentally unselected, population-based sample of Mexican American kinships, for which longitudinal data are available on 724 individuals. All participants provided informed consent and all research was conducted under approval of respective institutional review boards.

Materials and Methods

Data collection

Anthropometric, medical, and body composition data were collected during medical examinations at baseline (from 1997 to 2000) and follow-up (3.5 to 8.9 years later; mean = 5.6 years; from 2003 to 2006). Lifestyle, medical history, and reproductive history data were concomitantly assessed at both times via questionnaire. Approximately 81% of the original study participants were re-enrolled for follow-up.

The aim of the present study was to assess environmental and genetic influences on early BMD change; therefore analysis was performed in the subset of participants aged 25 to 45 years at baseline ($n = 327$). This cohort comprised participants from 32 kinships, including 1,434 relative pairs (206 sibling pairs, 96 avuncular relationships, 484 first cousin pairs, and 648 other relationships).

BMD measurements of femoral neck, total hip, total lumbar spine (L1-L4), ultradistal radius, and 33% radius (measured at 33% of total length from the distal end) were obtained by dual-energy x-ray absorptiometry (DXA) at both baseline and follow-up. Different sites of radius are included because they represent different proportions of trabecular and cortical bone: ultradistal radius is largely trabecular, whereas 33% radius is mostly cortical bone. During the interim between baseline and follow-up clinic visits, DXA equipment was upgraded from the Hologic 1500W model to the 4500W model (Hologic Inc., Bedford, MA), along with a software update to ensure comparability of scoring algorithms. Cross-calibration of absorptiometers showed near-perfect agreement on 10 test subjects ($R^2 = 99.9\%$, 99.8% , and 99.9% for spine, total hip, and femoral neck sites, respectively; $p < 10^{-13}$ for all). No mean difference between absorptiometers was detected (paired t-test $p > 0.1$ for all sites). Precision of Hologic 1500W was 0.009 g/cm^2 for spine, 0.007 g/cm^2 for total hip, and 0.002 g/cm^2 for the manufacturer's spine phantom. Precision of Hologic 4500W was 0.006 g/cm^2 for spine, 0.007 g/cm^2 for hip, and 0.002 g/cm^2 for radius. For quality control, all DXA readings were performed by the same trained technician, measurement drift was prevented by calibrating equipment daily on the phantom, and baseline and follow-up scans were evaluated by the same reviewer ensuring comparability of regions of interest.

To prevent any unknown inter-machine differences from affecting our assessment of rates of BMD change, measurements of BMD were standardized (mean = 0, SD = 1) independently at baseline and follow-up. Annual change in standardized BMD was calculated as the difference divided by exact elapsed time between standardized baseline and follow-up measurements. This process retains the information of individual measurements relative to the sample, but not of the absolute magnitude of measurements, thus avoiding potential unknown machine differences that could invalidate direct inter-machine comparisons. Rates of BMD change are expressed as percent of a standard deviation per year (%SD/year). The formula for calculating this metric of BMD change is

$$y_i = \frac{100}{t_i} \cdot \left(\frac{y_{fi} - \bar{y}_f}{\sigma_f} - \frac{y_{bi} - \bar{y}_b}{\sigma_b} \right)$$

where y_i is the rate of BMD change for the i th individual, y_{bi} is the baseline BMD for the i th individual, $\bar{y}_b$ is the mean baseline BMD, σ_b is the standard deviation of baseline BMD, y_{fi} is the follow-up BMD for the i th individual, $\bar{y}_f$ is the mean follow-up BMD, σ_f is the standard deviation of follow-up BMD, and t_i is the elapsed time between baseline and follow-up measurements.

Based on the precision of our DXA equipment, the least significant change (LSC; detectable with 95% confidence) was ± 2.22 %SD/year for hip, ± 2.54 %SD/year for spine, and ± 1.53 %SD/year for forearm. Observed rates of change less than the LSC values were not significantly different than zero (i.e. no observed change). Of the 327 participants included in this study, 65%, 73%, and 81% experienced significant rates of change in BMD (i.e. greater than the LSC) at the hip, spine, and radius, respectively.

Covariates included in our analysis were sex, baseline age (years), site-specific baseline BMD (g/cm^2), baseline body mass index (BMI; kg/m^2), yearly interim change in BMI ($\text{kg}/\text{m}^2/\text{year}$; calculated as difference between baseline and follow-up measurements divided by the exact elapsed time), and interim entrance into menopause (yes or no; defined as surgical menopause or 1 or more years since most recent menstrual cycle). Covariate measurements were collected identically at baseline and follow-up clinic visits, as previously described [11].

Genotypes

Automated genotyping of the San Antonio Family Osteoporosis Study participants has previously been described [19]. DNA from lymphocytes was amplified via polymerase chain reaction using fluorescently-tagged primers (MapPairs Human Screening Set Versions 6 and 8; Research Genetics, Huntsville, AL, USA) to detect repeat alleles at highly polymorphic microsatellite markers. Aliquots of amplified DNA were genotyped with Applied Biosystems Model 377 DNA Sequencers and analyzed with GeneScan and Genotyper DNA Fragment Analysis software (Perkin Elmer, Foster City, CA, USA). CRI-MAP [45] was used to assemble 460 microsatellite markers across chromosomes 1 to 22 into a genetic map, for which all marker positions were confirmed by deCODE (deCODE genetics, Reykjavik, Iceland). Mean inter-marker distance was 7.6 cM, ranging from <0.1 cM to 15.7 cM (Haldane).

Statistical analyses

Distributions of BMD (at baseline and follow-up) and covariates were assessed, and outliers greater than 4 SD from trait means were excluded (0 to 3 observations removed per trait). Because all traits were relatively normally distributed, no transformations were necessary. Heritability of rate of BMD change was estimated in a variance components framework, which models phenotypic variance as a function of effects due to measured covariates, additive polygenic (based on average allele-sharing between relative pairs), and error components. The

general form of this model is $y_i = \mu + \sum_{j=1}^n \beta_j X_{ij} + g_i + e_i$, where y_i is the rate of BMD change for the i th individual, μ is the sample mean rate of BMD change, X_{ij} is the j th covariate for the i th individual, β_j is the corresponding regression coefficient, g_i is the additive polygenic effect, and e_i is the residual error effect. Model parameters were estimated using pedigree-based maximum likelihood methods, from which residual heritability (h^2_r , i.e. the proportion of phenotype variance attributable to the additive genetic component after removing variation due to covariates) was estimated. The significance of covariate and heritable components were tested via the likelihood ratio test, which compares the likelihood of models including and excluding each component. This test follows a chi-squared distribution with 1 degree of freedom for testing covariates, and a 50:50 mixture of a point mass at zero and a chi-squared distribution with 1 degree of freedom for testing heritability. Only covariates with significant effects at $\alpha = 0.1$ were retained in final models for each skeletal site. The proportion of variance explained by covariates was determined by comparing models including and excluding retained covariates. Models of rates of BMD change for each skeletal site were restricted to participants with data on all retained covariates. Power to detect true heritability of 0.25, 0.35, and 0.45 was 70%, 85%, and 95%, respectively, at a significance threshold of $\alpha = 0.05$.

Multipoint linkage analysis was performed by extending the variance components model described above to include the effect of a theoretical QTL, σ_m^2 , as a component of genetic variance. Multipoint identical-by-descent (IBD) probabilities across chromosomes 1 to 22 were estimated from genotype data of relatives via a Markov chain Monte Carlo algorithm implemented in Loki [46,47]. Maximum likelihood methods were used to estimate σ_m^2 based on the expected covariance due to IBD probabilities between relatives at each locus along the chromosomes. Significance of σ_m^2 was assessed by the likelihood ratio test, which compared the QTL model (i.e. including σ_m^2 as a component of variance) to the polygenic model (i.e. $\sigma_m^2=0$), and expressed as a logarithm of the odds (LOD) score ($\log_{10}$ of the likelihood ratio). This test follows a 50:50 mixed distribution of a zero point mass and a 1-degree of freedom chi-squared distribution. Empirical LOD score adjustment based on 10,000 simulated unlinked markers was used to guard against inflated LOD scores, which can occur due to deviations of the phenotype distribution from normality [48]. All LOD scores (and corresponding p-values) presented herein are adjusted. Power to detect linkage was low: approximately 25% and 40% at thresholds of LOD = 2.0 and 1.5, respectively, for a QTL explaining 25% of variance in rates of BMD change at any skeletal site.

Genetic modeling was performed in the set of all participants aged 25 to 45 years ($n = 327$) as well as the subset of age-eligible men and premenopausal women ($n = 292$). To assess whether differences in linkage results obtained from the total sample and premenopausal subset were due to diminished sample size (null hypothesis), or alternatively, due to inclusion of post-menopausal women, 100 random subsets of 292 individuals (sampled without regard to menopausal status) were used to generate an empirical distribution of the effect of reduced sample size [49]. Genetic analyses were performed using the Sequential Oligogenetic Linkage Analysis Routines (SOLAR) software [50]; data management, summary statistics, outliers, and figures were done in R (R Foundation for Statistical Computing, Vienna, Austria).

Results

Characteristics of the 327 participants included in this study are presented in Table 1. Participants returning for follow-up measurements did not differ in any characteristics from those not available for follow-up. On average, adiposity in this sample was high (BMI > 30) in both men and women, and weight increased (0.82 kg/year in women and 0.45 kg/year in men) over 5.6 years of follow-up (range = 3.5 to 8.9 years). Distributions of rate of BMD change (%SD/year) are shown in Figure 1. As shown, rate of BMD change (unadjusted for covariate effects) varied widely among young men and women. Change in BMD was calculated as the yearly difference between standardized (mean = 0, SD = 1) BMD measurements at baseline and follow-up in order to mitigate the effects of differences between baseline and follow-up measurements.

Environmental and genetic influences on rate of BMD change for participants aged 25 to 45 years are shown in Table 2 and Table 3. As expected, women who entered menopause during the interim between visits experienced significantly faster bone loss at the femoral neck, total hip, and total spine; however, this relationship was not observed for sites of the forearm (Table 2). Interestingly, women experienced slower rates of bone loss than did men (i.e. female sex was positively correlated with rate of BMD change) for all sites. Bone loss increased with increasing age only at the radius sites. The cumulative amount of variation attributable to measured covariates differed greatly between skeletal sites, ranging from 6% for the 33% radius to 24% for the femoral neck.

Significant residual heritability of rate of BMD change (Table 3) was observed for the total hip ($h^2_r = 0.34$, $p = 0.01$), ultradistal radius ($h^2_r = 0.34$, $p = 0.004$), and 33% radius ($h^2_r = 0.27$,

$p = 0.03$). Modest residual heritability was observed for the femoral neck ($h^2_r = 0.22$, $p = 0.06$). In contrast, rate of BMD change of the total spine was not heritable.

Because menopause is widely acknowledged to have profound effects on bone loss, genetic and environmental influences on rate of BMD change were also assessed after excluding 35 women who reported having undergone menopause during the years of follow-up. Residual heritability estimates in this subset were greater than those of the total group for the femoral neck ($h^2_r = 0.29$) and total hip ($h^2_r = 0.49$), similar for the ultradistal radius ($h^2_r = 0.34$), 33% radius ($h^2_r = 0.25$) and spine ($h^2_r = 0.0$). Standard errors and cumulative variance due to covariates (excluding the effect of menopause) were essentially unchanged.

Figure 2 depicts autosome-wide multipoint linkage scans for rate of BMD change of the femoral neck, total hip, ultradistal radius, and 33% radius (but not spine because change at this site was not heritable). Significant evidence for linkage was observed on chromosome 1q23 at 151 cM for rate of femoral neck BMD change (LOD = 3.6, adjusted $p = 0.00004$, Figure 3). No other chromosomal regions exhibited evidence of linkage at LOD > 1.5 in the entire sample. In the subset of men and premenopausal women, linkage was diminished for the femoral neck QTL (LOD = 2.9 on chromosome 1) and an additional suggestive QTL was identified for ultradistal radius (LOD = 2.9 on chromosome 6q26–27 at 189 cM, adjusted $p = 0.0003$) (results not shown).

To verify the robustness of our chromosome 1 QTL and determine whether the decrease in LOD score in the subset of men and premenopausal women was due to reduced sample size, or due to heterogeneity specific to postmenopausal women, we generated an empirical distribution for 100 subsets in which 35 (10.7%) individuals were excluded at random. The mean maximum LOD score across all leave-35-out subsets was 3.0 (at position 151 cM), which is not significantly different ($p = 0.45$) from our observed LOD = 2.9 in the subset of individuals that included only men and pre-menopausal women. Thus, the observed decrease in evidence of linkage in the subset is indistinguishable from the LOD-score decrease due to a reduction in sample size alone, and therefore we have no evidence that heterogeneity is contributing to our linkage signal.

Discussion

Cross-sectional and longitudinal studies have shown that BMD declines in young men and women of European [41,43,44] and African ancestry [10], and that most of this change results from loss of trabecular bone density. One method by which to increase our understanding of the mechanisms influencing early bone loss is to determine whether such longitudinal loss is heritable and, if so, to identify the specific genes involved.

In this study we show substantial variation in the individual rates of early areal BMD change at the femoral neck, total hip, ultradistal radius, and 33% radius, and estimate that genetic factors are responsible for up to 35% of this variation. In contrast, we found that early bone loss at the lumbar spine was not heritable, though the reason for this is unclear. Both baseline and follow-up spine BMD are heritable in young Mexican Americans ($h^2_r = 0.78$ and 0.71 , respectively), and the rate of change in BMC and area does not differ qualitatively from the other skeletal sites. Although our results include both men and women, they are consistent with those of Hui et al. [35] who reported heritability of change in femoral neck BMD in premenopausal women only. Our estimates of heritability for early BMD change are also similar, with a few exceptions, to those for bone loss among older individuals at several skeletal sites [36,37]. Together, this body of evidence suggests that rates of BMD loss are heritable for many skeletal sites, in both men and women, young and old.

Because BMD is calculated from the mineral content (g) over the 2-dimensional projected area (cm^2) of a bone, change in BMD may be reflective of either change in bone mineral content (BMC), change in bone area, or some combination of the two. In this sample, both BMC and area were changing, on average, in the same direction ($r = 0.53$ to 0.78), though the rates of percent change differed between BMC and area. For example, femoral neck BMC (mean change = -0.7% per year) was decreasing faster than area (-0.5% per year, $p = 0.001$). Percent change in BMC was also less than percent change in area for the ultradistal radius and lumbar spine ($p < 0.001$ for both). In contrast, percent change in BMC was greater than percent change in area for the total hip (1.2% versus 0.7% per year) and 33% radius ($p < 0.001$ for both). These observations suggest that genes affecting the interplay between BMC and bone area, rather than one or the other, are responsible for the observed heritability of BMD change; the specific genetic relationships among these traits within skeletal sites remain to be resolved.

Our linkage scan yielded a putative QTL influencing bone loss in young adults. The QTL for rate of change in femoral neck BMD on chromosome 1q23 (LOD = 3.6) has been reported to influence variation in peak spine BMD in several [17,21,27,30], but not all [51], previous cross-sectional studies; specifically, this QTL was not observed in our Mexican American sample for either peak femoral neck BMD [19] or femoral neck BMD loss in older individuals [37]. These possibly contradictory results could be explained by a gene or genes influencing BMD change, rather than peak BMD, in an age-specific manner. We know that that trabecular and cortical bone loss varies by age and skeletal site [10,43]. Thus, linkage studies for peak BMD [17,21] and BMD change in young individuals are well-suited to detect a genetic regulator of early BMD change, whereas linkage [19] and exclusion [51] analysis for BMD in mixed-aged samples, and linkage analysis for BMD change in older samples [37], are not.

The one-LOD unit support interval for the femoral neck QTL on chromosome 1q23 ranges from 146 to 155 cM (approximately 133 Mbp to 157 Mbp) in a gene-rich region containing 429 known or hypothetical genes. In particular, this region includes BGLAP/osteocalcin, which has well known effects on bone formation [52,53], and IL6R, for which previous studies have shown genetic association to BMD [54], and linkage to BMD and osteopenia [55]. Additionally, a zinc/iron transporter (SLC39A1; mouse homolog expressed in osteoblasts of developing bone) [56], and 17 calcium binding proteins (S100A1-A7, -A7.1, -A7.2-A14, -A16) are contained in the support interval. One or more of these genes, or other linked loci, may contribute to our observed linkage signal.

We also obtained suggestive evidence for a QTL influencing early BMD change at the 33% ulna on chromosome 11p14-15 (LOD = 2.5). Due to its position, this QTL is likely distinct from those detected in previous studies for peak BMD of the spine at 11q12-13, the *LRP5* locus [18,22], or 11q23 [26]. The one-LOD unit support interval for our suggestive QTL ranges from 22 to 42 cM (approximately 14 to 25 Mbp), covering 88 genes. Notable genes in this region are calcitonin alpha (*CALCA*), beta (*CALCB*), and pseudogene (*CALCP*). Calcitonin is a thyroid hormone regulating serum calcium levels, and has known inhibitory effects on the resorptive activity of osteoclasts [57]. Calcitonin could also be involved in early BMD change and is a reasonable positional candidate gene for BMD change.

We speculate that because the femoral neck is comprised of both cortical and trabecular bone, and because growing evidence suggests that early bone loss occurs primarily in the latter [10,43], the observed linkage to chromosome 1q23 may be due to one or more (possibly site-specific) genes affecting femoral neck trabecular bone loss. Though direct measurements of trabecular and cortical content are not available in this study, this hypothesis reflects recent findings in other populations [10,43], and is consistent with the largely uncorrelated linkage scans for different skeletal sites.

A major strength of the current study is the combined longitudinal and family-based study design. Our assessment of 5-year BMD change is better suited to elucidate the genetic effects of early bone loss than cross-sectional linkage scans, and the inclusion of many types of relatives pairs within our extended families allows us to better model truly genetic effects on BMD change (as opposed to familial non-genetic effects, such as household effects), than do studies using twins or sibships. These strengths notwithstanding, several limitations of the present study also need to be addressed. First, while statistical power to detect heritability of rate of BMD change was adequate, power to detect linkage at genome-wide significance was poor, and our sample size precluded testing for gene $\times$ sex or gene $\times$ environment interactions. Also, densitometry equipment was upgraded during the follow-up interim which may have introduced measurement bias; however, due to the independent standardization of baseline and follow-up measurements and excellent agreement observed in our cross-calibration experiment, any unknown effects of our equipment upgrade are probably minimal. Finally, DXA technology is inherently limited in its ability to detect BMD change because areal projection may not adequately account for the size (depth) of bone or possible change in bone size over time, and cannot measure trabecular and cortical content separately.

In conclusion, this study reports that rate of early change in BMD at several skeletal sites is heritable and similar in magnitude to the heritability of late bone loss [35,37]. The QTL for rate of early BMD change that we detected on chromosome 1q23 is among the first loci implicated in longitudinally-assessed BMD change, and may act as candidate regions for additional investigations into the genetic determinants of early bone loss. Identification of specific genes or pathways influencing early BMD change may lead to better understanding of bone health and possible early therapeutic interventions for individuals at greater risk of developing osteoporosis.

Acknowledgements

We are deeply grateful to the participants of the San Antonio Family Osteoporosis Study. We would also like to acknowledge two anonymous reviewers for their thoughtful suggestions. This work was funded by grants M01-RR-01346, P01-HL-45522, R01-AR-43351, and R37-MH-59490, awarded by the National Institutes of Health.

References

1. Hui SL, Slemenda CW, Johnston CC Jr. The contribution of bone loss to postmenopausal osteoporosis. *Osteoporos Int* 1990;1:30–34. [PubMed: 2133638]
2. Hannan MT, Felson DT, Dawson-Hughes B, Tucker KL, Cupples LA, Wilson PW, Kiel DP. Risk factors for longitudinal bone loss in elderly men and women: the Framingham Osteoporosis Study. *J Bone Miner Res* 2000;15:710–720. [PubMed: 10780863]
3. Jordan KM, Cooper C. Epidemiology of osteoporosis. *Best Pract Res Clin Rheumatol* 2002;16:795–806. [PubMed: 12473274]
4. Wilkins CH, Birge SJ. Prevention of osteoporotic fractures in the elderly. *Am J Med* 2005;118:1190–1195. [PubMed: 16271899]
5. Dirschl DR, Henderson RC, Oakley WC. Accelerated bone mineral loss following a hip fracture: a prospective longitudinal study. *Bone* 1997;21:79–82. [PubMed: 9213011]
6. Engelke K, Kemmler W, Lauber D, Beeskov C, Pintag R, Kalender WA. Exercise maintains bone density at spine and hip EFOPS: a 3-year longitudinal study in early postmenopausal women. *Osteoporos Int* 2006;17:133–142. [PubMed: 16096715]
7. Melton LJ 3rd, Khosla S, Atkinson EJ, O'Connor MK, Ofallon WM, Riggs BL. Cross-sectional versus longitudinal evaluation of bone loss in men and women. *Osteoporos Int* 2000;11:592–599. [PubMed: 11069193]
8. Warming L, Hassager C, Christiansen C. Changes in bone mineral density with age in men and women: a longitudinal study. *Osteoporos Int* 2002;13:105–112. [PubMed: 11905520]

9. Christian JC, Yu PL, Slemenda CW, Johnston CC Jr. Heritability of bone mass: a longitudinal study in aging male twins. *Am J Hum Genet* 1989;44:429–433. [PubMed: 2916585]
10. Wang X, Kammerer CM, Wheeler VW, Patrick AL, Bunker CH, Zmuda JM. Genetic and environmental determinants of volumetric and areal BMD in multi-generational families of African ancestry: the Tobago Family Health Study. *J Bone Miner Res* 2007;22:527–536. [PubMed: 17227221]
11. Mitchell BD, Kammerer CM, Schneider JL, Perez R, Bauer RL. Genetic and environmental determinants of bone mineral density in Mexican Americans: results from the San Antonio Family Osteoporosis Study. *Bone* 2003;33:839–846. [PubMed: 14623060]
12. Brown LB, Streeten EA, Shapiro JR, McBride D, Shuldiner AR, Peyser PA, Mitchell BD. Genetic and environmental influences on bone mineral density in pre- and post-menopausal women. *Osteoporos Int* 2005;16:1849–1856. [PubMed: 15997421]
13. Brown LB, Streeten EA, Shuldiner AR, Almasy LA, Peyser PA, Mitchell BD. Assessment of sex-specific genetic and environmental effects on bone mineral density. *Genet Epidemiol* 2004;27:153–161. [PubMed: 15305331]
14. Deng HW, Chen WM, Conway T, Zhou Y, Davies KM, Stegman MR, Deng H, Recker RR. Determination of bone mineral density of the hip and spine in human pedigrees by genetic and life-style factors. *Genet Epidemiol* 2000;19:160–177. [PubMed: 10962476]
15. Deng HW, Xu FH, Huang QY, Shen H, Deng H, Conway T, Liu YJ, Liu YZ, Li JL, Zhang HT, Davies KM, Recker RR. A whole-genome linkage scan suggests several genomic regions potentially containing quantitative trait loci for osteoporosis. *J Clin Endocrinol Metab* 2002;87:5151–5159. [PubMed: 12414886]
16. Devoto M, Specchia C, Li HH, Caminis J, Tenenhouse A, Rodriguez H, Spotila LD. Variance component linkage analysis indicates a QTL for femoral neck bone mineral density on chromosome 1p36. *Hum Mol Genet* 2001;10:2447–2452. [PubMed: 11689491]
17. Econs MJ, Koller DL, Hui SL, Fishburn T, Conneally PM, Johnston CC Jr, Peacock M, Foroud TM. Confirmation of linkage to chromosome 1q for peak vertebral bone mineral density in premenopausal white women. *Am J Hum Genet* 2004;74:223–228. [PubMed: 14730478]
18. Johnson ML, Gong G, Kimberling W, Recker SM, Kimmel DB, Recker RB. Linkage of a gene causing high bone mass to human chromosome 11 (11q12-13). *Am J Hum Genet* 1997;60:1326–1332. [PubMed: 9199553]
19. Kammerer CM, Schneider JL, Cole SA, Hixson JE, Samollow PB, O'Connell JR, Perez R, Dyer TD, Almasy L, Blangero J, Bauer RL, Mitchell BD. Quantitative trait loci on chromosomes 2p, 4p, and 13q influence bone mineral density of the forearm and hip in Mexican Americans. *J Bone Miner Res* 2003;18:2245–2252. [PubMed: 14672361]
20. Karasik D, Myers RH, Cupples LA, Hannan MT, Gagnon DR, Herbert A, Kiel DP. Genome screen for quantitative trait loci contributing to normal variation in bone mineral density: the Framingham Study. *J Bone Miner Res* 2002;17:1718–1727. [PubMed: 12211443]
21. Koller DL, Econs MJ, Morin PA, Christian JC, Hui SL, Parry P, Curran ME, Rodriguez LA, Conneally PM, Joslyn G, Peacock M, Johnston CC, Foroud T. Genome screen for QTLs contributing to normal variation in bone mineral density and osteoporosis. *J Clin Endocrinol Metab* 2000;85:3116–3120. [PubMed: 10999795]
22. Koller DL, Rodriguez LA, Christian JC, Slemenda CW, Econs MJ, Hui SL, Morin P, Conneally PM, Joslyn G, Curran ME, Peacock M, Johnston CC, Foroud T. Linkage of a QTL contributing to normal variation in bone mineral density to chromosome 11q12-13. *J Bone Miner Res* 1998;13:1903–1908. [PubMed: 9844108]
23. Peacock M, Koller DL, Fishburn T, Krishnan S, Lai D, Hui S, Johnston CC, Foroud T, Econs MJ. Sex-specific and non-sex-specific quantitative trait loci contribute to normal variation in bone mineral density in men. *J Clin Endocrinol Metab* 2005;90:3060–3066. [PubMed: 15741260]
24. Peacock M, Koller DL, Hui S, Johnston CC, Foroud T, Econs MJ. Peak bone mineral density at the hip is linked to chromosomes 14q and 15q. *Osteoporos Int* 2004;15:489–496. [PubMed: 15205721]
25. Ralston SH, Galwey N, MacKay I, Albagha OM, Cardon L, Compston JE, Cooper C, Duncan E, Keen R, Langdahl B, McLellan A, O'Riordan J, Pols HA, Reid DM, Uitterlinden AG, Wass J, Bennett

- ST. Loci for regulation of bone mineral density in men and women identified by genome wide linkage scan: the FAMOS study. *Hum Mol Genet* 2005;14:943–951. [PubMed: 15746152]
26. Shen H, Zhang YY, Long JR, Xu FH, Liu YZ, Xiao P, Zhao LJ, Xiong DH, Liu YJ, Dvornyk V, Rocha-Sanchez S, Liu PY, Li JL, Conway T, Davies KM, Recker RR, Deng HW. A genome-wide linkage scan for bone mineral density in an extended sample: evidence for linkage on 11q23 and Xq27. *J Med Genet* 2004;41:743–751. [PubMed: 15466007]
 27. Streeten EA, McBride DJ, Pollin TI, Ryan K, Shapiro J, Ott S, Mitchell BD, Shuldiner AR, O'Connell JR. Quantitative trait loci for BMD identified by autosome-wide linkage scan to chromosomes 7q and 21q in men from the Amish Family Osteoporosis Study. *J Bone Miner Res* 2006;21:1433–1442. [PubMed: 16939402]
 28. Styrkarsdottir U, Cazier JB, Kong A, Rolfsson O, Larsen H, Bjarnadottir E, Johannsdottir VD, Sigurdardottir MS, Bagger Y, Christiansen C, Reynisdottir I, Grant SF, Jonasson K, Frigge ML, Gulcher JR, Sigurdsson G, Stefansson K. Linkage of osteoporosis to chromosome 20p12 and association to BMP2. *PLoS Biol* 2003;1:E69. [PubMed: 14691541]
 29. Hsu YH, Xu X, Terwedow HA, Niu T, Hong X, Wu D, Wang L, Brain JD, Bouxsein ML, Cummings SR, Rosen CJ, Xu X. Large-scale genome-wide linkage analysis for loci linked to BMD at different skeletal sites in extreme selected sibships. *J Bone Miner Res* 2007;22:184–194. [PubMed: 17228995]
 30. Ioannidis JP, Ng MY, Sham PC, Zintzaras E, Lewis CM, Deng HW, Econs MJ, Karasik D, Devoto M, Kammerer CM, Spector T, Andrew T, Cupples LA, Foroud T, Kiel DP, Koller B, Langdahl B, Mitchell BD, Peacock M, Recker R, Shen H, Sol-Church K, Spotila LD, Uitterlinden AG, Wilson SG, Kung AW, Ralston SH. Meta-Analysis of Genome Wide Scans Provides Evidence for Gender and Site Specific Regulation of Bone Mass. *J Bone Miner Res* 2007;22:173–183. [PubMed: 17228994]
 31. Karasik D, Cupples LA, Hannan MT, Kiel DP. Age, gender, and body mass effects on quantitative trait loci for bone mineral density: the Framingham Study. *Bone* 2003;33:308–316. [PubMed: 13678771]
 32. Lee YH, Rho YH, Choi SJ, Ji JD, Song GG. Meta-analysis of genome-wide linkage studies for bone mineral density. *J Hum Genet* 2006;51:480–486. [PubMed: 16534542]
 33. Wilson SG, Reed PW, Bansal A, Chiano M, Lindersson M, Langdown M, Prince RL, Thompson D, Thompson E, Bailey M, Kleyn PW, Sambrook P, Shi MM, Spector TD. Comparison of genome screens for two independent cohorts provides replication of suggestive linkage of bone mineral density to 3p21 and 1p36. *Am J Hum Genet* 2003;72:144–155. [PubMed: 12478480]
 34. Xiao P, Shen H, Guo YF, Xiong DH, Liu YZ, Liu YJ, Zhao LJ, Long JR, Guo Y, Recker RR, Deng HW. Genomic regions identified for BMD in a large sample including epistatic interactions and gender-specific effects. *J Bone Miner Res* 2006;21:1536–1544. [PubMed: 16995807]
 35. Hui SL, L KD, Foroud TM, Econs MJ, Johnston CC, Peacock M. Heritability of changes in bone size and bone mass with age in premenopausal white sisters. *J Bone Miner Res* 2006;21:1121–1125. [PubMed: 16813533]
 36. Makovey J, Nguyen TV, Naganathan V, Wark JD, Sambrook PN. Genetic effects on bone loss in peri- and postmenopausal women: a longitudinal twin study. *J Bone Miner Res* 2007;22:1773–1780. [PubMed: 17620052]
 37. Shaffer JR, Kammerer CM, Bruder JM, Cole SA, Dyer TD, Almasy L, MacCluer JW, Blangero J, Bauer RL, Mitchell BD. Genetic influences on bone loss in The San Antonio Family Osteoporosis Study. *Osteoporos Int*. 2008(in press)
 38. Kelly PJ, Nguyen T, Hopper J, Pocock N, Sambrook P, Eisman J. Changes in axial bone density with age: a twin study. *J Bone Miner Res* 1993;8:11–17. [PubMed: 8427043]
 39. Seeman E. Pathogenesis of bone fragility in women and men. *Lancet* 2002;359:1841–1850. [PubMed: 12044392]
 40. Slemenda C, Longcope C, Peacock M, Hui S, Johnston CC. Sex steroids, bone mass, and bone loss. A prospective study of pre-, peri-, and postmenopausal women. *J Clin Invest* 1996;97:14–21. [PubMed: 8550826]
 41. Hui SL, Zhou L, Evans R, Slemenda CW, Peacock M, Weaver CM, McClintock C, Johnston CC Jr. Rates of growth and loss of bone mineral in the spine and femoral neck in white females. *Osteoporos Int* 1999;9:200–205. [PubMed: 10450407]

42. Uusi-Rasi K, Sievanen H, Pasanen M, Oja P, Vuori I. Association of physical activity and calcium intake with the maintenance of bone mass in premenopausal women. *Osteoporos Int* 2002;13:211–217. [PubMed: 11991440]
43. Riggs BL, Melton LJ, Robb RA, Camp JJ, Atkinson EJ, McDaniel L, Amin S, Rouleau PA, Khosla S. A population-based assessment of rates of bone loss at multiple skeletal sites: evidence for substantial trabecular bone loss in young adult women and men. *J Bone Miner Res* 2008;23:205–214. [PubMed: 17937534]
44. Nordstrom P, Neovius M, Nordstrom A. Early and rapid bone mineral density loss of the proximal femur in men. *J Clin Endocrinol Metab* 2007;92:1902–1908. [PubMed: 17311855]
45. Green, P.; Falls, K.; Crooks, S. Documentation for CRI-MAP, Version 2.4. St. Louis, MO, USA: Washington University School of Medicine; 1990. In:
46. Heath SC. Markov chain Monte Carlo segregation and linkage analysis for oligogenic models. *Am J Hum Genet* 1997;61:748–760. [PubMed: 9326339]
47. Heath SC, Snow GL, Thompson EA, Tseng C, Wijsman EM. MCMC segregation and linkage analysis. *Genet Epidemiol* 1997;14:1011–1016. [PubMed: 9433616]
48. Blangero J, Williams JT, Almasy L. Robust LOD scores for variance component-based linkage analysis. *Genet Epidemiol* 2000;19:S8–S14. [PubMed: 11055364]
49. Atwood LD, Heard-Costa NL, Fox CS, Jaquish CE, Cupples LA. Sex and age specific effects of chromosomal regions linked to body mass index in the Framingham Study. *BMC Genet* 2006;7:7. [PubMed: 16438729]
50. Almasy L, Blangero J. Multipoint quantitative-trait linkage analysis in general pedigrees. *Am J Hum Genet* 1998;62:1198–1211. [PubMed: 9545414]
51. Chen XD, Shen H, Lei SF, Li MX, Yang YJ, Deng HW. Exclusion mapping of chromosomes 1, 4, 6 and 14 with bone mineral density in 79 Caucasian pedigrees. *Bone* 2006;38:450–455. [PubMed: 16249131]
52. Aubin, JE.; Lian, JB.; Stein, GS. Chapter 4. Bone formation: maturation and function activities of osteoblast lineage cells. In: Lian, JB.; Goldring, SR., editors. *Primer on the metabolic bone diseases and disorders of mineral metabolism*. The American Society of Bone and Mineral Research; 2006. p. 20-29.
53. Rosen, CJ.; Kiel, DP. Chapter 12. Age-related osteoporosis. In: Shane, E.; Kleerekoper, M., editors. *Primer on the metabolic bone diseases and disorders of mineral metabolism*. The American Society for Bone and Mineral Research; 2006. p. 71-74.
54. Bustamante M, Nogues X, Mellibovsky L, Agueda L, Jurado S, Caceres E, Blanch J, Carreras R, Diez-Perez A, Grinberg D, Balcells S. Polymorphisms in the interleukin-6 receptor gene are associated with bone mineral density and body mass index in Spanish postmenopausal women. *Eur J Endocrinol* 2007;157:677–684. [PubMed: 17984249]
55. Ota N, Hunt SC, Nakajima T, Suzuki T, Hosoi T, Orimo H, Shirai Y, Emi M. Linkage of interleukin 6 locus to human osteopenia by sibling pair analysis. *Hum Genet* 1999;105:253–257. [PubMed: 10987653]
56. Lioumi M, Ferguson CA, Sharpe PT, Freeman T, Marenholz I, Mischke D, Heizmann C, Ragoussis J. Isolation and characterization of human and mouse ZIRT1, a member of the IRT1 family of transporters, mapping within the epidermal differentiation complex. *Genomics* 1999;62:272–280. [PubMed: 10610721]
57. Deftos, LJ. Chapter 18. Calcitonin. In: Christakos, S.; Holick, MF., editors. *Primer on the metabolic diseases and disorders of mineral metabolism*. The American Society for Bone and Mineral Research; 2006. p. 115-117.

Figure 1. Distributions of change in BMD (%SD/year) for (A) femoral neck, (B) total hip, (C) ultradistal radius, (D) 33% radius, and (E) total spine. Grey bars represent the total sample (women + men); black bars represent women.

Figure 2.
Autosome-wide multipoint linkage scans

Figure 3. Multipoint LOD score profile for femoral neck BMD change on chromosome 1 in total sample (solid line) and subset of men and premenopausal women (dashed line).

Table 1

Mean (SD) population characteristics

variable	all	(SD)	women	(SD)	men	(SD)
sample size, n	325		210		115	
follow-up (years)	5.6	(0.7)	5.5	(0.7)	5.6	(0.7)
age (years)	34.4	(5.9)	34.6	(5.8)	34.2	(6.1)
anthropometrics						
height (cm)	162.2	(8.8)	157.4	(6.1)	170.8	(6.2)
weight (kg)	80.4	(20.7)	75.9	(19.2)	88.6	(20.8)
BMI (kg/m ²)	30.5	(7.0)	30.6	(7.4)	30.3	(6.3)
annual weight gain (kg/year)	0.69	(1.39)	0.82	(1.40)	0.45	(1.35)
annual change in BMI (kg/m ² /year)	0.27	(0.52)	0.33	(0.55)	0.15	(0.45)
medical						
diabetes (%)	10.7	-	10.4	-	11.2	-
pre-menopausal (%)	-	-	95.3	-	-	-
oral contraceptives (%)	-	-	20.9	-	-	-
interim entrance into menopause (%)	-	-	9.2	-	-	-
lifestyle						
alcohol consumption (%)	50.8	-	44.5	-	62.1	-
smoking history (%)	20.8	-	19.4	-	23.3	-
baseline BMD (g/cm ²)						
femoral neck	0.89	(0.13)	0.87	(0.12)	0.92	(0.13)
total hip	0.99	(0.15)	0.96	(0.13)	1.05	(0.15)
radius (ultradistal)	0.50	(0.07)	0.47	(0.05)	0.55	(0.06)
radius (33%)	0.72	(0.07)	0.68	(0.04)	0.79	(0.05)
total spine	1.05	(0.12)	1.05	(0.12)	1.04	(0.13)
follow-up BMD (g/cm ²)						
femoral neck	0.87	(0.12)	0.86	(0.12)	0.89	(0.13)
total hip	1.01	(0.14)	0.99	(0.13)	1.06	(0.14)
radius (ultradistal)	0.49	(0.06)	0.47	(0.05)	0.53	(0.07)
radius (33%)	0.74	(0.07)	0.71	(0.05)	0.80	(0.06)
total spine	1.03	(0.12)	1.04	(0.11)	1.01	(0.12)

Table 2
Relationship between rate of early BMD change and covariates: beta-coefficients (p-values)

	femoral neck		total hip		radius (ultradistal)		radius (33%)		total spine	
N	300		300		313		313		295	
R ²	0.24		0.16		0.14		0.06		0.14	
sex	1.9 ^a	(0.023)	2.2	(<0.001)	5.1	(<0.001)	2.5	(0.001)	3.8	(<0.001)
age	-	(0.738)	-	(0.284)	-0.2	(0.002)	-1.6	(0.012)	-	(0.489)
meno ^c	-4.4	(0.008)	-3.3	(0.009)	-	(0.699)	-	(0.336)	-5.5	(0.001)
BMI	5.1	(<0.001)	0.2	(<0.001)	-0.1	(0.097)	-	(0.343)	-0.1	(0.045)
ΔBMI	0.6	(<0.001)	3.0	(<0.001)	1.4	(0.064)	-1.6	(0.03)	-	(0.642)
BMD	-15.9	(<0.001)	-	(0.663)	14.9	(0.045)	-	(0.164)	-10.6	(0.002)

R² = proportion of variance due to covariates
beta-coefficients are interpreted as %SD/year per unit of covariate
^a effect of female sex with respect to male sex
^b only covariates with significant effects at $\alpha = 0.1$ are included in genetic models
^c meno = entrance into menopause during interim between baseline and follow-up;
ΔBMI = yearly change in BMI during interim (kg/m²/year);
BMD = site-specific baseline BMD (g/cm²)

Table 3

Residual heritability of early BMD change

Trait	n	h^2_r	SE	p-value
femoral neck	300	0.22	0.16	0.06
total hip	300	0.34	0.17	0.01
radius (ultradistal)	313	0.34	0.16	0.004
radius (33%)	313	0.27	0.16	0.03
total spine	295	0.00	-	0.50

h^2_r = residual heritability, while simultaneously accounting for variation due to covariates enumerated in Table 2. Only covariates with significant effects at $\alpha = 0.1$ are included in genetic models.