Published in final edited form as:

J Am Chem Soc. 2009 April 29; 131(16): 5763–5765. doi:10.1021/ja901283q.

A Method for the Preparation of Differentiated *trans*-1,2-Diol Derivatives with Enantio- and Diastereocontrol

Sang Min Lim, Nicholas Hill, and Andrew G. Myers

Department of Chemistry and Chemical Biology, Harvard University, Cambridge, Massachusetts 02138

Abstract

We describe a synthetic sequence that allows for the preparation of optically active *trans*-1,2-diol monosilyl ether derivatives from ketones, providing a new means for retrosynthetic simplification of differentiated diol and polyol targets. The sequence involves silyl enol ether formation, Shi asymmetric epoxidation, then regio- and stereospecific addition of hydride, methide, or higher alkylide. The tactical combination presented has not been integrated in synthetic problem solving, so far as we are aware, but has promise for broad application, we believe.

We describe a synthetic sequence that allows for the preparation of optically active *trans*-1,2-diol monosilyl ether derivatives from ketones, providing a new means for retrosynthetic simplification of differentiated diol and polyol targets. The sequence is exemplified in equation 1 and proceeds by silyl enol ether formation, Shi asymmetric epoxidation,¹ then regio- and stereospecific addition of hydride, methide, or higher alkylide. The tactical combination presented has not been integrated in synthetic problem solving, so far as we are aware, but has promise for broad application, we believe.

(1)

It has been shown that α -hydroxy ketones can be prepared enantioselectively by application of the Shi asymmetric epoxidation protocol to *tert*-butyldimethylsilyl and trimethylsilyl enol ether substrates followed by hydrolysis.² Shi and coworkers noted that certain substrates were problematic in the transformation, especially silyl enol ethers derived from cyclic ketones (racemization and dimerization were specifically mentioned as complicating factors). This led them to direct their studies toward enol ester substrates, where the primary epoxidation products were sufficiently stable to be isolated and manipulated, providing an essential enablement of their general method for the enantiocontrolled synthesis of α -acyloxy ketones.³

While prior research has shown that trimethylsilyloxy and *tert*-butyldimethylsilyloxy epoxides can be prepared as discrete intermediates, especially by epoxidation with non-acidic oxygen atom-transfer agents such as arylsulfonyloxaziridines⁴ and dimethyldioxirane,⁵ it is also evident from this prior work that the reactivity of the products complicate their isolation and handling, absent special stabilizing structural features. With these precedents in mind, and aware of the particular stability of *tert*-butyldiphenylsilyl ethers toward acid-catalyzed hydrolysis,⁶ we undertook to reinvestigate the Shi asymmetric epoxidation of silyl enol ether derivatives with a focus on variation of the trialkylsilyl group. Here we show that *tert*-butyldiphenylsilyl enol ether derivatives of both cyclic and acyclic ketones undergo enantioselective epoxidation according to the Shi paradigm,¹ that the *tert*-butyldiphenylsilyloxy epoxide products can be isolated by simple extraction, and, most significantly, that these products undergo highly regio- and stereospecific addition reactions with certain hydride and alkylidene donors (*vide infra*).

tert-Butyl(cyclohexenyloxy)diphenylsilyl ether (**1**), prepared from cyclohexanone in >95% yield under soft enolization conditions (*tert*-butyldiphenylsilyl triflate,⁷ triethylamine, dichloromethane, 0 °C, 1 h)⁸ or by deprotonation with potassium hexamethyldisilazide followed by trapping with the silyl chloride, is, as expected, substantially more stable than the corresponding triisopropylsilyl and *tert*-butyldimethylsilyl enol ethers^{6,9} (in that order) under conditions such as exposure to silica gel (see Supporting Information). All three substrates were transformed into the corresponding silyloxy epoxides under typical Shi epoxidation conditions (0.30 equiv of Shi catalyst,¹⁰ borate buffer, slow addition of Oxone and potassium carbonate solutions over 90 min at 0 °C) with similar, high enantioselectivities (91-93% ee, *vide infra*).² The products were efficiently extracted into pentane without decomposition.⁴ Two-dimensional thin-layer chromatographic analysis of the product solutions revealed the following stability order: *tert*-butyldiphenylsilyl (compound **2**) >> triisopropylsilyl > *tert*-butyldimethylsilyl (see Supporting Information). Solutions of the epoxide **2** could be concentrated and the crude product stored neat for several weeks at -20 °C without evident decomposition. ¹H NMR spectral data were consistent with expectations for the proposed structure (**2**) and revealed that the product had been formed cleanly, with the only significant

contaminant being a small amount of Shi catalyst that had partitioned into the pentane layer during the extraction. While future applications may benefit from or require that purified silyloxy epoxides be used, all of the transformations described herein were conducted using the pentane extracts of the Shi epoxidation reaction mixtures directly, without purification, or even concentration in many cases.

Among the new and useful reactions of silyloxy epoxides we have investigated is a simple regio- and stereoselective reduction with borane-THF, which gives rise to *trans*-diol monosilyl ethers.¹¹ In a specific illustration (eq 1 and entry 1, Table 1), addition of a 1.0 M solution of borane-THF (1.2 equiv) to an ice-cold solution of the pentane extracts containing the Shi epoxidation product **2**, stirring at 0 °C for 1 h, then careful quenching with an aqueous solution of tris(hydroxy-methyl)aminomethane, extractive isolation, and chromatographic purification afforded pure (1*R*,2*R*)-2-(*tert*-butyldiphenylsilyloxy)cyclohexanol **3** in 91% yield for the two-step (epoxidation-reduction) sequence. The product was shown to be of 93% ee by ¹H NMR analysis of the corresponding Mosher ester.¹² Similarly, the triisopropylsilyl and *tert*-butyldimethylsilyl enol ethers (entries 2 and 3, respectively, Table 1) were transformed directly into the corresponding *trans*-1,2-diol monosilyl ethers, in yields that reflected the stabilities of the silyloxy epoxide intermediates.

As the examples of Table 1 reveal, the two-step Shi epoxidation-reduction sequence appears to hold promise as a general method for the enantio- and diastereocontrolled synthesis of differentiated cyclic *trans*-1,2-diols, and as entry 9 suggests, may also allow for simplification of certain polyol targets by simultaneous multiple application.^{13,14} Because epimerization is unlikely to have occurred at any point during the transformations summarized in Table 1, we believe the ee values presented there provide an accurate assessment of the enantioselectivities of the epoxidation step, which are typically above 90%, in keeping with Shi's prior observations.¹

The stereochemistry of hydride addition supports a mechanism involving prior coordination of borane to the epoxide oxygen atom followed by epoxide opening and internal hydride transfer, as has been proposed for reductions of glycol epoxide-like substrates with borane-THF,¹⁵ although the trajectories for hydride addition are presumably quite different.¹⁶

In further evaluating the scope of the epoxidation-reduction sequence we explored acyclic *tert*-butyldiphenylsilyl enol ethers as substrates and in this context gained mechanistic insight into the reduction process. Asymmetric epoxidation-reduction of *trans*-1-*tert*-butyldiphenylsilyloxypropene¹⁷ afforded (*R*)-1-(*tert*-butyldimethylsilyloxy)propan-2-ol in 90% yield and 82% ee (eq 2). Employing BD₃-THF in lieu of BH₃-THF we found that the reduction proceeded with >95% stereospecificity (eq 3),¹⁸ as observed in the cyclic substrate series (Table 1). Epoxidation-reduction (also with BD₃-THF) of *cis*-1-*tert*-butyldiphenylsilyloxypropene,¹⁹ however, proceeded with substantially diminished stereospecificity (~33%, eq 4), which we rationalize in Figure 1.²⁰

(2)

(3)

(4)

Two additional examples in the acyclic series, employing trisubstituted *Z*-*tert*-butyldiphenylsilyl enol ethers as substrates (each $\geq 14:1$, *Z*:*E*, prepared by enolate formation with potassium hexamethyldisilazide followed by trapping with *tert*-butyldiphenylsilyl chloride), suggest that the present method may have general value for the preparation of differentiated *anti*-1,2-diols (eqs 5 and 6).

(5)

(6)

Lastly, we have observed that *tert*-butyldiphenylsilyloxy epoxides react stereospecifically with trimethyl- and triethylaluminum to form differentiated *trans*-1,2-diol products (eqs 7-9). It is noteworthy that the tertiary hydroxyl groups of the product diols emerge bearing the *tert*-butyldiphenylsilyl protecting group. Thus far, we have not seen evidence of silyl group transfer. Here, too, substantial literature precedent exists for stereospecific additions of trialkylaluminum reagents to glycol epoxides²¹ although, as discussed above in the context of

hydride addition,¹⁶ the stereoelectronic features of the present transformations are presumably very different and seemingly less favorable.

(7)

(8)

(9)

We imagine that the asymmetric epoxidation-reduction and epoxidation-alkylide addition sequences presented herein will be useful for the preparation of a number of complex diol and polyol targets.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgment

We thank the NSF (CHE-0749566), the NIH/NCI (CHE-0749566), the NIH/NIGMS (GM007598-30) (NH), the Kwanjeong Educational Foundation Fellowship (SML), the Eli Lilly Organic Chemistry Fellowship (SML), the

Harvard College Research Program, Pfizer Inc., Amgen, and Merck & Co., Inc. for financial support of this research. We thank Dr. Richard Staples and Dr. Douglas Ho for X-ray crystallographic analyses.

References

- (1). (a) Tu Y, Wang Z-X, Shi Y. J. Am. Chem. Soc 1996;118:9806–9807. (b) Wang Z-X, Tu Y, Frohn M, Zhang J-R, Shi Y. J. Am. Chem. Soc 1997;119:11224–11235. (c) Frohn M, Shi Y. Synthesis 2000:1979–2000. (d) Shi Y. Acc. Chem. Res 2004;37:488–496. [PubMed: 15311947]
- (2). (a) Adam W, Fell RT, Saha-Möller CR, Zhao C-G. Tetrahedron: Asymmetry 1998;9:397–401. (b) Zhu Y, Tu Y, Yu H, Shi Y. Tetrahedron Lett 1998;39:7819–7822. (c) Solladié-Cavallo A, Lupattelli P, Jerry L, Bovicelli P, Angeli F, Antonioletti R, Klein A. Tetrahedron Lett 2003;44:6523–6526.
- (3). Zhu Y, Manske KJ, Shi Y. J. Am. Chem. Soc 1999;121:4080–4081.
- (4). Davis FA, Sheppard AC. J. Org. Chem 1987;52:954–955.
- (5). (a) Chenault HK, Danishefsky SJ. J. Org. Chem 1989;54:4249–4250. (b) Adam W, Hadjiarapoglou L, Wang X. Tetrahedron Lett 1989;30:6497–6500. (c) Schaumann E, Tries F. Synthesis 2002:191–194.
- (6). Hanessian S, Lavallee P. Can. J. Chem 1975;53:2975–2977.
- (7). Dimopoulos P, George J, Tocher DA, Manaviazar S, Hale KJ. Org. Lett 2005;7:5377–5380. [PubMed: 16288510]
- (8). For a different method of preparing compound **1**, see: Martel A, Leconte S, Dujardin G, Brown E, Maisonneuve V, Retoux R. Eur. J. Org. Chem 2002:514–525.
- (9). Cunico RF, Bedell L. J. Org. Chem 1980;45:4797–4798.
- (10). The catalyst was prepared in two steps from d-Fructose as described in reference 1b.
- (11). The transformation is functionally equivalent to an asymmetric hydroboration-oxidation of a silyl enol ether substrate. In the one example of such a process of which we are aware cyclohexanone trimethylsilyl enol ether was transformed into *trans*-1,2-cyclohexanediol monotrimethylsilyl ether with diisopinocampheylborane-alkaline hydrogen peroxide in 31% yield and 28% ee: Peterson PE, Stepanian M. J. Org. Chem 1988;53:1903–1907.
- (12). Dale JA, Dull DL, Mosher HS. J. Org. Chem 1969;34:2543–2549. The absolute stereochemistry of product **3** was confirmed by comparison (optical rotation) with an authentic sample prepared by an independent route.
- (13). The absolute stereochemistry of the product (entry 9, Table 1) was established by X-ray crystallographic analysis of the corresponding bis-*p*-bromophenyl ester derivative (see Supporting Information). Interestingly, we observed that when a pentane-ethyl acetate solution of the bis-epoxide intermediate of entry 9 was shaken briefly with a 1 N aqueous hydrochloric acid solution, the hemiketal monosilyl ether depicted was obtained as a stable substance. The structure was verified by X-ray crystallography (see Supporting Information).
- (14). (a) Xiong Z, Corey EJ. J. Am. Chem. Soc 2000;122:9328–9329. (b) Lorenz JC, Frohn M, Zhou X, Zhang J-R, Tang Y, Burke C, Shi Y. J. Org. Chem 2005;70:2904–2911. [PubMed: 15822948]
- (15). (a) Bazin HG, Kerns RJ, Linhardt RJ. Tetrahedron Lett 1997;38:923–926. (b) Koyama Y, Yamaguchi R, Suzuki K. Angew. Chem. Int. Ed 2008;47:1084–1087.
- (16). We speculate that hydride transfer in the two processes may involve different conformations of the respective six-membered rings: (ref 15a)
- (17). Prepared by iridium-catalyzed isomerization of allyloxy(*tert*-butyl)diphenylsilane: Ohmura T, Yamamoto Y, Miyaura N. Organometallics 1999;18:413–416.
- (18). Using a commercial source of BD₃-THF with 11% hydrogen content (determined by ¹H NMR analysis of the corresponding BD₃-dimethylphenylphosphine complex), we found the product of eq 5 was enriched in hydrogen (17% hydrogen content at the site of transfer), consistent with a kinetic isotope effect of approximately 1.6.
- (19). Guha SK, Shibayama A, Abe D, Sakaguchi M, Ukaji Y, Inomata K. Bull. Chem. Soc. Jpn 2004;77:2147–2157.
- (20). The products of this two-step transformation were also formed with diminished enantioselectivity (~40% ee), in keeping with prior examples of Shi epoxidations of *cis*-1,2-disubstituted alkenes (see reference 1b).

- (21). a) Bailey JM, Craig D, Gallagher PT. Synlett 1999:132–134. (b) Rainier JD, Cox JM. Org. Lett 2000;2:2707–2709. [PubMed: 10990433]

Figure 1.

Reduction of the *cis*-*tert*-butyldiphenylsilyloxy epoxide **4** with $\text{BD}_3\text{-THF}$ provides evidence for a short-lived carbocationic intermediate. The data suggest that the rate of deuteride transfer is slightly more rapid than rotation about the internal C-C bond.

Table 1

Cyclic *trans*-1,2-Diol Monosilyl Ether Derivatives Formed by Shi Asymmetric Epoxidation^a of Silyl Enol Ethers followed by Stereospecific Reduction with Borane-THF

Entry	Substrate
-------	-----------

TBDPSO

1

Entry

Substrate

2

EntrySubstrate

TBDPSO

3

Entry

Substrate

4

5

Entry	Substrate
6

Entry

Substrate

TBDPSO

N
Boc

7

Entry

Substrate

8^d

Entry

Substrate

^a Shi epoxidation conditions: 0.3 equiv Shi catalyst, 1.38 equiv Oxone, 5.8 equiv K₂CO₃, CH₃CN-CH₃OCH₂OCH₃-borate buffer, 0 °C, 2 h.

^b Isolated yields over two steps. ¹H NMR analysis revealed that in all cases diastereoselectivities were > 20:1, favoring the *trans*-1,2-diol derivatives.

^c Enantiomeric excesses were determined by ¹H NMR analysis of the corresponding Mosher esters, except for entries 7 and 8, where ee's were determined by HPLC using a chiral column (see Supporting Information for details).

^d These entries involved slight procedural modifications; see Supporting Information for details.