Regulation of p53 by TopBP1: a Potential Mechanism for p53 Inactivation in Cancer^{∇†}

Kang Liu,¹ Naresh Bellam,¹ Hui-Yi Lin,^{2‡} Bing Wang,¹ Cecil R. Stockard,³
William E. Grizzle,³ and Weei-Chin Lin^{1,4*}

Division of Hematology and Oncology, Department of Medicine,¹ Medical Statistics Section, Department of Medicine,²
Department of Pathology,³ and Department of Cell Biology,⁴ University of Alabama at Birmingham,
Birmingham, Alabama 35294-3300

Received 18 July 2008/Returned for modification 29 August 2008/Accepted 3 March 2009

Proper control of the G₁/S checkpoint is essential for normal proliferation. The activity of p53 must be kept at a very low level under unstressed conditions to allow growth. Here we provide evidence supporting a crucial role for TopBP1 in actively repressing p53. Depletion of TopBP1 upregulates p53 target genes involved in cell cycle arrest and apoptosis and enhances DNA damage-induced apoptosis. The regulation is mediated by an interaction between the seventh and eighth BRCT domains of TopBP1 and the DNA-binding domain of p53, leading to inhibition of p53 promoter binding activity. Importantly, TopBP1 overexpression is found in 46 of 79 primary breast cancer tissues and is associated with high tumor grade and shorter patient survival time. Overexpression of TopBP1 to a level comparable to that seen in breast tumors leads to inhibition of p53 target gene expression and DNA damage-induced apoptosis and G₁ arrest. Thus, a physiological level of TopBP1 is essential for normal G₁/S transition, but a pathological level of TopBP1 in cancer may perturb p53 function and contribute to an aggressive tumor behavior.

The tumor suppressor protein p53 is mutated or inactivated in most human cancers. Mutations of p53 are found in about half of these tumors. Even in tumors without p53 mutations, p53 can be inactivated indirectly through other mechanisms such as binding to viral proteins or alterations in p53 regulators (47). The tumor suppressor function of p53 is ascribed mainly to its pivotal role in causing cell cycle arrest or apoptosis in response to genomic, hypoxic, or oncogenic stresses (15). In response to DNA damage, p53 activates p21^{Cip1} and arrests the cells in G₁ phase—this constitutes the major G₁/S checkpoint. Induction of p53 can also lead to apoptosis by activating proapoptotic proteins such as BAX, PUMA, and NOXA. With the activities in growth arrest and apoptosis, p53 must be tightly controlled during normal growing conditions. The regulation of p53 occurs at multiple levels. Through phosphorylation by ATM and Chk2 kinases, p53 is stabilized and activated in response to DNA damage. Other modifications such as acetylation, methylation, and ubiquitination also regulate p53 function (36). In addition, many proteins have been identified to interact with p53 and modulate its activity (1, 2). It is believed that through these complex but delicate regulations, different activities of p53 can be modulated properly in response to various environmental stresses.

TopBP1 (topoisomerase IIβ binding protein) contains eight BRCA1 carboxyl-terminal (BRCT) motifs, which are found in proteins involved in DNA repair and cell cycle checkpoints.

TopBP1 also contains a conserved ATR-activating domain (23) and is recruited by the Rad9-Hus1-Rad1 (9-1-1) clamp to stalled replication forks for Chk1 activation (6). Xmus101, the *Xenopus laevis* homolog of TopBP1, is required for the loading of Cdc45 and DNA polymerases α and ε to replication origins (11, 45). Human TopBP1 interacts with DNA polymerase ε as well (29) and may either monitor the replication (22) or actively recruit the replication preinitiation complex to chromatin (18).

In addition to a role in replication, TopBP1 also possesses activity in transcriptional regulation. TopBP1 binds to a specific member of E2F transcription factors, E2F1 (26), and represses its proapoptotic activity by recruiting Brg1/Brm chromatin-remodeling complex (27). The repression of E2F1 activity by TopBP1 requires phosphatidylinositol 3-kinase/Akt. Akt phosphorylates TopBP1 at Ser¹¹⁵⁹ and induces its oligomerization. The oligomerized TopBP1 then binds and represses E2F1 (28). TopBP1 also binds and represses Miz1 to inhibit p21^{Cip1} expression (14). Like E2F1, the binding and functional repression of Miz1 require Akt-dependent oligomerization of TopBP1 (28). Phosphorylation by Akt at Ser¹¹⁵⁹ is required for TopBP1 to regulate other transcription factors such as human papillomavirus type 16 E2 (28) and an ePHD protein, SPBP (38), as well. TopBP1 also contains several transcriptional regulatory domains (51). Thus, TopBP1 may have a more general role in transcriptional regulation.

Consistent with a role in promoting growth and survival, the expression of TopBP1 rises at G₁/S transition and S phase. In fact, TopBP1 is an E2F target (27, 52) and feed-back-regulates E2F1 by blocking E2F1-mediated apoptosis during S-phase entry. This is evidenced by induction of E2F1-dependent apoptosis upon depletion of TopBP1 (27). However, the apoptosis induced by TopBP1 small interfering RNA (siRNA) is not completely abrogated in *E2F1*^{-/-}

* Corresponding author. Mailing address: Shelby 815, 1530 3rd Ave. S, Birmingham, AL 35294-3300. Phone: (205) 934-3979. Fax: (205) 975-6911. E-mail: wclin@uab.edu.

† Supplemental material for this article may be found at <http://mcb.asm.org/>.

‡ Present address: Department of Biostatistics, Moffitt Cancer Center and Research Institute, Tampa, FL 33612.

[∇] Published ahead of print on 16 March 2009.

FIG. 1.

MEFs (mouse embryonic fibroblasts), suggesting that TopBP1 also regulates other proapoptotic molecules or a proapoptotic pathway. Besides apoptosis, TopBP1 plays a role in the regulation of p21^{Cip1} and p27 expression for activation of cyclin E/cdk2 (18). Therefore, TopBP1 appears

to function as a modulator between G₁ and S phases to ensure proper transition into DNA replication.

In this report, we show that TopBP1 is required to keep p53 activities in check during normal growth. This regulation is mediated through an interaction between the seventh

FIG. 1—Continued.

and eighth BRCT domains (BRCT78) of TopBP1 and the p53 DNA-binding domain (DBD), leading to inhibition of the promoter-binding activity of p53. Deregulation of this control can have pathological consequences. We show that overexpression of TopBP1 is frequently seen in breast cancer and is associated with high-grade tumors and poor survival. Therefore, abnormally high levels of TopBP1 may inactivate p53 potentially through deregulation of the Rb pathway and contribute to an aggressive behavior of breast tumors. Our results suggest that targeting TopBP1 can enhance the response of p53 and potentiate the apoptotic response during chemotherapy and radiation.

MATERIALS AND METHODS

Cell culture and transfection. HEK293, C33A, NIH 3T3, MCF7, and Saos-2 cells and primary human foreskin fibroblasts were maintained in Dulbecco's modified Eagle's medium supplemented with 10% fetal bovine serum (FBS). HCT116 ($p53^{+/+}$ and $p53^{-/-}$) and U2OS cells were grown in McCoy's medium supplemented with 10% FBS. H358 cells were grown in RPMI 1640 supplemented with 10% FBS. A standard calcium phosphate method was used for transfection of C33A cells. HEK293 cells were transfected with the calcium phosphate method or the Gene Pulser Xcell electroporation system (Bio-Rad) according to the manufacturer's instruction for 293 cells. Saos-2 cells were transfected using Lipofectamine 2000 (Invitrogen).

Plasmid construction. All the TopBP1-related constructs were described previously (26–28). Full-length human p53, p53 DBD, p53 C terminus, and p53 Δ C deletion mutants were amplified by PCR and then inserted into expression vector pGEX6P1, encoding glutathione S-transferase (GST). GST-p53N (containing amino acids [aa] 1 to 70) has been described previously (25). The sequences were verified by sequencing. Primer sequences for PCR are 5'-CTCGGATCCATGG AGGAGCCGAGCAG-3' and 5'-CTCGAATTCTCAGTCTGAGTCAGGCC C-3' for full length, 5'-CTCGGATCCATGTCATCTTCTGTCCTTCCAG-3' and 5'-CTCGAATTCTCATTCTTTCGGGAGATTCTC-3' for DBD, and 5'-CTCGGATCCATGCCAGGGAGCACTAAGCGAGC-3' and 5'-CTCGAATTC TCAGTCTGAGTCAGGCC-3' for C terminus.

EYFP1-p53 and EYFP2-p53 were obtained by cloning the EGFP1-p53 BspEI/ApaI fragment into pcDNA3.1EYFP(1 and 2)-zipper vector (35). EYFP1-TopBP1 and YFP2-TopBP1 were obtained by cloning the HcRed1-TopBP1 BspEI/ApaI fragment into pcDNA3.1EYFP(1 and 2)-zipper vector. Ad-siGFP and Ad-siTopBP1 were established with the AdEasy system (12). The NotI/SalI fragments of pSUPER constructs containing H1 RNA gene promoter and RNA interference for green fluorescent protein (28) and TopBP1 (27) were cloned to pShuttle. Viruses were purified by CsCl banding.

Immunoprecipitation and Western blot analysis. Cells were harvested 24 to 48 h after transfection in TNN buffer (50 mM Tris [pH 7.5], 0.2 M NaCl, 5 mM EDTA [pH 8.0], 0.5% NP-40), and immunoprecipitation was carried out as described previously (26). The specific signals were detected with appropriate antibodies. The antibodies specific to p53 (FL-393), GST (B-14), hemagglutinin (HA) (Y-11), Myc (A-14), p21 (C-19), and p63 (4A4) were purchased from Santa Cruz. The cyclin A antibody was purchased from Upstate Biotechnology (06-138). The β -actin antibody was purchased from Sigma. p53 (Ab-1) was purchased from Calbiochem. Glyceraldehyde-3-phosphate dehydrogenase (GAPDH) antibody was purchased from Alexis. p73 (5B429) was purchased from Imgenex. The anti-green fluorescent protein antibody was purchased from Clontech. The poly-(ADP-ribose) polymerase antibody and the monoclonal and polyclonal TopBP1 antibodies were purchased from BD Transduction Laboratories and Bethyl Laboratories, respectively. A goat polyclonal TopBP1 antibody (L20) for immunoprecipitation was purchased from Santa Cruz.

GST pull-down assay. The GST fusion proteins were induced by 0.1 mM IPTG (isopropyl- β -D-thiogalactopyranoside) in *Escherichia coli* strain BL21 and purified. The GST portion on GST-TopBP1 and its mutants were excised by Pre-Scission protease (Pharmacia). One microgram of purified GST-p53, its different deletion mutants, or GST was incubated in NETN-A buffer (50 mM NaCl, 1 mM EDTA, 20 mM Tris, 0.5% NP-40) with 2 μ g purified TopBP1 or its deletion mutants and rotated at 4°C for 3 h. GST-p53 and its mutants were pulled down with glutathione Sepharose, and the beads were washed six times with NETN-B buffer (100 mM NaCl, 1 mM EDTA, 0.2 mM phenylmethylsulfonyl fluoride) and then subjected to sodium dodecyl sulfate-polyacrylamide gel electrophoresis (SDS-PAGE) and analyzed by Western blotting with anti-TopBP1 or anti-GST antibody.

ChIP assay. A chromatin immunoprecipitation (ChIP) assay was performed following the protocol as described previously (42). Briefly, HCT116 cells grown in 15-cm² dishes were left untreated or treated with adriamycin (5 μ M) for 5 h and then cross-linked with formaldehyde. Cells were collected, and chromatin was extracted and sonicated. One-tenth of supernatants were used as input control for PCR, and the other chromatin was precleared with protein G plus/protein A-agarose beads and then immunoprecipitated with 4 μ g of each antibody (p53 and Ab-1 from Calbiochem; TopBP1 from BD Transduction Laboratories). The antibody-bound complexes were recovered on protein A/G beads. Immunoprecipitates were washed under stringent conditions, and cross-links of chromatin were reversed by incubating samples at 65°C overnight. The resulting DNA was purified and analyzed by PCR. The PCR primers for E2F1 and β -actin promoters have been described previously (27). The PCR primers derived from p21^{Cip1} promoter are 5'-GTGGCTCTGATTGGCTTTCTG-3' and 5'-CTGAA AACAGGCAGCCCAAG-3'. The PCR primers derived from HDM2 promoter are 5'-TGGGCAGGTTGACTCAGCTTTTCTC-3' and 5'-TGGCGTGCCTC CGTCCCCAC-3'. The PCR primers derived from NOXA promoter are 5'-TG CCCATCTCTTCAAGTTCG-3' and 5'-AGGGTATTGCGGCAGACGC-3'. The PCR primers derived from the IGFBP3 promoter are 5'-TGCTGAGGTG CCTGGAGT-3' and 5'-TCCAGGCAGGAAGCGCTGATC-3'.

Real-time and semiquantitative RT-PCR. NIH 3T3 and HCT116 ($p53^{+/+}$ or $p53^{-/-}$) cells were infected with Ad-siGFP or Ad-siTopBP1, AdCMV, or Ad-TopBP1 and then either left untreated or treated with adriamycin (1 μ M) for 24 h. RNA was extracted using TRIzol reagent (Invitrogen). Reverse transcription-PCR (RT-PCR) was performed using the following primer pairs: p21, 5'-GAGTCAGGCGCAGATCCACAG-3' and 5'-GGGCACTTCAGGGTTTTC-3', 521 bp; mouse MDM2, 5'-GGATCCTTTGCAAGCGCCAC-3' and 5'-TCA AAGGACAGGGACCTGCG-3', 162 bp; human HDM2, 5'-ATCTTGGCCAG TATATTATG-3' and 5'-GTTCTGTAGATCATGGTAT-3', 152 bp; mouse NOXA, 5'-CGTCGGAACGCGCCAGTGAACCC-3' and 5'-TCCTTCTGAGG AGGTCCCTTCTTGC-3', 336 bp; IGFBP3, 5'-GACAGCCAGCGCTACAAA GTTGAC-3' and 5'-ACTTATCCACACACCAGCAGAAGC-3', 259 bp; CAB1, 5'-AAGCTCGGCCAGATGCTGAGCATC-3' and 5'-GGGAACAGG

FIG. 1. Interaction between TopBP1 and p53 in vivo. (A) Interaction between endogenous TopBP1 and p53 in primary human foreskin fibroblasts (HFF), HEK293, and MCF7 cells. HFF, HEK293, or MCF7 cells were left untreated or treated with the radiomimetic agent NCS at 300 ng/ml or adriamycin at 5 μ M for 5 h. Lysates were immunoprecipitated (IP) with an anti-TopBP1 antibody, an anti-p53 antibody, or a control mouse immunoglobulin G (IgG) antibody as indicated, followed by immunoblotting. One-tenth of cell lysates before immunoprecipitation were analyzed by Western blotting and probed with p53 and TopBP1. The films with the same exposure times as those for immunoprecipitation are shown. (B) HEK293 cells were transfected with TopBP1-FLAG and then left untreated or treated with NCS at 300 ng/ml for 3 h. FLAG-tagged TopBP1 was immunoprecipitated from cell lysates, and the coimmunoprecipitated p53 was detected by immunoblotting. The relative intensities of the signals were quantified by densitometry and are shown below each panel. (C) $p53^{+/+}$ HCT116 cells were suspended in hypotonic buffer (10 mM HEPES-KOH, pH 7.5, 10 mM KCl, 3 mM MgCl₂, 0.05% NP-40, 1 mM EDTA) for 30 min. Cytosolic and nuclear fractions were lysed in TNN buffer and subject to p53 immunoprecipitation followed by immunoblotting. Immunoblotting of a cytoplasmic protein, vinculin, was performed to ensure the effectiveness of subcellular fractionation. Arrows indicate the position of p53. The asterisk indicates a cytosolic protein which cross-reacts with rabbit anti-p53 antibody. (D) Growing HCT116 and MCF7 cells were harvested in TNN buffer. The cell lysates were either left untreated or pretreated with 1 μ g/ml DNase for 20 min at 37°C or 10 μ g/ml ethidium bromide (EtBr) on ice for 30 min. The lysates were then immunoprecipitated with an anti-p53 antibody or a control mouse immunoglobulin G antibody followed by immunoblotting as indicated. Due to the short exposure time, p53 signal in the input lane was not detected, but it was detected after longer exposure (lower panels). One-tenth of cell lysates before immunoprecipitation were analyzed by Western blotting (lower panels). (E and F) Bimolecular fluorescence complementation assay. HEK293 or NIH 3T3 cells were transiently transfected with split yellow fluorescent protein expression plasmids as indicated. The split yellow fluorescent protein was tagged at the N terminus of p53 and TopBP1. Nuclei were stained with Hoechst 33258. The reconstituted yellow fluorescent protein signals due to protein-protein interaction between YFP1 and YFP2 fusion proteins were captured on a Zeiss fluorescence microscope. Because cells are limited by the transfection efficiency, not every cell shows yellow fluorescent protein signal. Magnification, $\times 100$ (E). (G) Cellular lysates of HEK293 cells in panel E were analyzed for expression of YFP1- or YFP2-TopBP1 or p53 using either TopBP1 or p53 antibody for immunoblotting. Endogenous TopBP1 or p53 was also detected. YFP1 peptide added 17.9 kDa to the fused protein, and YFP2 peptide added 9.3 kDa.

FIG. 2. Mapping the interaction regions of TopBP1 and p53 by GST pull-down assay. (A) A direct interaction between TopBP1 and p53 in vitro. TopBP1 and GST-p53 proteins were produced and purified from *E. coli*. Purified TopBP1 or TopBP1(S1159A) protein was first phosphorylated with Akt kinase and then incubated with purified GST or GST-p53. GST pull-down with glutathione-Sepharose was performed, and GST was probed with the indicated antibodies. Bottom panel: Coomassie blue staining of input TopBP1 (wild-type [W]) or TopBP1(S1159A) (A) proteins. (B and C) HEK293 cells were transfected with FLAG-TopBP1 expression plasmid or FLAG-tagged truncated mutants. The lysates were then incubated with GST or GST-p53, and a GST pull-down assay was performed as described for panel A. One-tenth of input lysates were loaded along with GST pull-down in panel B. One-fifth of input lysates were immunoprecipitated with FLAG beads and run along with GST pull-down in panel C. "IgG" (immunoglobulin G) denotes the position of anti-FLAG heavy chain. (D and E) GST pull-down assay was performed to test the interaction between purified TopBP1, TopBP1-BRCT78, or TopBP1-BRCT678 and purified GST-p53 or its domains: p53 DBD, p53CT (C terminus), p53ΔCT (lacking C terminus), and p53N (N terminus). All the proteins were produced in *E. coli* and purified as described in Materials and Methods. The proteins in panel D were resolved by 12% SDS-PAGE. Coomassie blue staining of the GST fusion proteins is shown in the bottom panel of panel E. We repeatedly observed the presence of a higher-molecular-weight product of GST-p53CT (**), which can be recognized by both anti-GST and anti-p53 antibodies and might represent a dimeric form of GST-p53CT, since the C terminus is known to mediate tetramerization of p53. * denotes the monomeric form.

CCTTCTGGAAGCATG-3', 347 bp; HIC1, 5'-AAGAGCAGCGAGGAGACC GGT-3' and 5'-TGGCCGGTCTTGTAGCTCTTGT-3', 373 bp; mouse GADD45, 5'-GCGGTTTCAGAAAGATGCAGGC-3' and 5'-GGTTGTGCCCAA TGTCCTCG-3', 303 bp; human GADD45, 5'-CGGCGCCTGTGAGTGAGTG

C-3' and 5'-GATGTTGATGTCGTTCTCGC-3', 370 bp; human PUMA, 5'-G ACCTCAACGCACAGTA-3' and 5'-CTAATTGGGCTCCATCT-3', 145 bp; mouse and human TopBP1, 5'-AAGAGTTTCCTTGTGTTTGGG-3' and 5'-CA TGCCTTCTTTCGATTGG-3', 402 bp; 14-3-3 σ , 5'-GGAGAGAGCCAGTCT

FIG. 3. A physiological role of TopBP1 in the control of endogenous p53 target genes. (A) NIH 3T3 cells were transfected with a control siRNA vector (pSUPER-siGFP) or pSUPER-siTopBP1. Twenty-four hours later, the cells were infected with adenoviruses either harboring an empty vector or expressing human TopBP1 at a multiplicity of infection of 400. Two days later, RNA was extracted and RT-PCR analysis was performed using primers specific for the selected p53 target genes or GAPDH as indicated. Left panels: mRNA expression levels of p53 target genes were measured using semiquantitative RT-PCR. The "no RNA" lane represents a control RT-PCR without input RNA. Right panels: the changes of p53 target gene expression were confirmed using quantitative real-time PCR. Results were normalized to GAPDH levels and are expressed relative to the expression of the gene in the siGFP control. Except for 14-3-3 σ , the *P* values for the differences of gene expression between siGFP and siTopBP1 and between siTopBP1 and rescue groups are all less than 0.02 (paired two-tailed *t* test). (B) Saos-2 cells were transfected with the expression vectors for p53 and TopBP1 or the TopBP1 domain-deletional mutants, along with a p53 activity reporter plasmid (a p21 promoter-luciferase plasmid) and pCMV- β gal. Luciferase activity of transfected p53 was determined as induction relative to that of empty-vector control. Each sample was assayed in triplicate. A portion of the cellular lysates was immunoblotted with antibody for p53 or immunoprecipitated with anti-FLAG agarose followed by anti-FLAG immunoblotting (upper panels). The asterisks indicate the corresponding TopBP1 or mutant products. (C) A schematic diagram of the TopBP1 and p53 mutant proteins used in Fig. 2 and this figure and a summary of their binding properties and functional activities. p53 aa 1 to 42, transactivation domain; 63 to 97, proline-rich domain; 324 to 355, tetramerization domain; 363 to 393, regulatory domain. (D) HEK293 cells were transfected with expression plasmids as indicated. Myc-tagged p63 α , HA-tagged p73 α , or FLAG-tagged TopBP1 was immunoprecipitated from cell lysates and immunoblotted by antibodies specific to TopBP1, Myc, or HA. (E) The transcriptional activity of p53, p63 α , or p73 α was assayed in Saos-2 cells, which were transfected with the expression vectors for p53, p63 α , or p73 α and TopBP1, along with a p21 promoter-luciferase plasmid and pCMV- β gal as described for panel B. The expression levels of p63 α , p73 α , and TopBP1 were determined by immunoblotting (upper panels).

FIG. 3—Continued.

GATC-3' and 5'-TCTCCTTCTGCTGATGTCC-3', 482 bp; Bax, 5'-AGGGTTCATCCAGGATCGAGC-3' and 5'-ACAGGGCCTTGAGCACCAGTTTC-3', 294 bp; GAPDH, 5'-TGAAGGTCCGAGTCAACGGATTGGT-3' and 5'-AAATGAGCCCCAGCCTTCTCCATG-3', 325 bp. PCR was performed for 22 to 28 cycles at 95°C for 30 s for the denaturation step, 55°C for 1 min for annealing, and 72°C for 2 min. We ensured linear amplification in all cases. Quantitative PCR was performed in triplicate on an MX3000p thermal cycler (Stratagene) using the SYBR green dye method to track the progress of the reactions with ROX dye added as reference. GAPDH was run in parallel with test genes. Results were analyzed with MxPro 4.0 quantitative PCR software (Stratagene).

Luciferase assay. The expression constructs (5 μ g for pCMV-p53; 5 μ g for Myc-p63; 5 μ g for HA-p73; 10 μ g for FLAG-WT-TopBP1, FLAG- Δ 123TopBP1, FLAG- Δ 45TopBP1, FLAG- Δ 678TopBP1, FLAG-678TopBP1, FLAG- Δ 78TopBP1, and FLAG-78TopBP1; 10 or 15 μ g for pcDNA3-TopBP1; 10 or 15 μ g for pcDNA3-TopBP1-S1159A; 10 μ g for Brg1 and Brm), the promoter plasmid (1 μ g for pGL2-p21 promoter-Luc; gift of Xinbin Chen), and 1 μ g of β -galactosidase plasmid were transfected into Saos-2 cells or C33A cells. Cells were harvested 2 days later. An aliquot of cells was lysed in SDS lysis buffer for Western blot analysis in some experiments; the other cell extracts were lysed in reporter lysis buffer (Promega) for luciferase activity and β -galactosidase activity as described previously (26). The luciferase activity was normalized against the β -galactosidase activity. All transient expression steps in this assay were carried out in triplicate.

p53 EMSA. Nuclear extracts from adenovirus-infected $p53^{-/-}$ HCT116 cells were prepared as described previously (24). The p53 electrophoretic mobility shift assay (EMSA) was carried out as described previously (5) except that the LightShift chemiluminescent EMSA kit (Pierce, Rockford, IL) was used for detection. The biotin-labeled DNA probe is derived from p21 promoter (5'-biotin-AGCTTGAACATGTCCCAACATGTTGAGCTC-3'). The wild-type and mutant p53 binding-site-containing oligonucleotides for competition are as follows: wild type, 5'-AGCTTGAACATGTCCCAACATGTTGAGCTC-3', and mutant, 5'-AGCTTGAATGCATCCCAACATGTTGA-3' (boldface sequence indicates swapping of nucleotides to mutate the p53 binding site). The complexes were resolved in a 6% polyacrylamide gel.

Immunohistochemical staining. Sections of paraffin-embedded tissue (5 μ m) were mounted on Bond-Rite slides from Richard-Allan Scientific if the antigen retrieval was to be in pH 6 buffer (10 mM citrate). If the section was destined for pH 9 antigen retrieval (10 mM Tris, 1 mM EDTA), the slides were precoated with chrome alum gelatin for better tissue adhesion. The slides were then heated to 60°C for 2 h. Paraffin was removed from the sections by three changes of xylene and rehydrated through graded alcohols from absolute to 70% for 5 minutes each. For immunohistochemistry, the slides were rinsed with deionized water and high-temperature antigen retrieval was performed in pH 9 or pH 6 buffer in a pressure cooker for 10 minutes. The slides were then allowed to cool for 20 min and transferred to Tris buffer (Trizma base [0.05 M], NaCl [0.15 M], Triton X-100 [0.1%], pH adjusted to 7.6 with HCl). Endogenous peroxidases were quenched with an aqueous solution of H₂O₂ for 5 minutes. Biotin was blocked by treating the tissues with streptavidin (Jackson Immuno Research, West Grove PA) at 10 μ g/ml in phosphate-buffered saline for 15 min, rinsing them with Tris buffer, and applying biotin (Sigma-Aldrich, St. Louis, MO) at 200

μ g/ml in PBE buffer (phosphate-buffered saline with 1% bovine serum albumin, 1 mM EDTA, and 0.15 mM Na₂S₂O₃). Nonspecific binding was blocked with 3% goat serum in PBE buffer on the sections for 20 min. The primary antibodies were diluted in PBE buffer and applied to the sections for 1 hour at room temperature. TopBP1 antibody (clone 33; BD Transduction Laboratories, San Jose, CA) was used at a dilution of 1:250 with pH 9 antigen retrieval or a dilution of 1:50 with pH 6 antigen retrieval. Negative controls were incubated with 3% goat serum. Slides were rinsed with Tris buffer after each step. Immunodetection was performed with a biotinylated anti-mouse secondary antibody and streptavidin-horseradish peroxidase (Signet Pathology Systems, Dedham, MA) for 20 min each. The chromogen used was 3,3'-diaminobenzidine (Scy Tek, Logan, UT). After 7 minutes the slides were rinsed with water and lightly counterstained with Mayer's hematoxylin. The sections were dehydrated through graded alcohols, 70% to absolute, and three xylene baths for 5 minutes each. The coverslips were mounted with Permount.

Flow cytometry. To assay apoptosis, cells were stained with annexin V-phycoerythrin (PE) (Pharmingen) and 7-amino-actinomycin (7-AAD) (Pharmingen). The annexin-7-AAD profile was analyzed by flow cytometry. The DNA content profiles of the HCT116 ($p53^{+/+}$ or $p53^{-/-}$) cells were analyzed by propidium iodide staining followed by flow cytometry.

Frozen breast cancer tissue collection. Frozen breast tumor or normal tissues were collected with informed consent through the University of Alabama at Birmingham (UAB) Tissue Procurement Core Facility under an institutional review board-approved protocol. Samples 1-20, 1T, and 7T were collected between 1990 and 2003. The rest of the samples were collected between 2004 and 2007. All patients underwent surgery at the UAB and were diagnosed at UAB. The majority of patients were followed at UAB for further treatment, but seven patients underwent follow-up with oncologists at different medical centers. The clinical and pathological information was obtained from a retrospective review of the medical record at UAB. The frozen tissues were minced and lysed in 1% SDS-60 mM Tris-HCl and boiled for 5 min. The lysates were sonicated briefly and clarified by centrifugation to remove insoluble tissues. To analyze TopBP1 protein levels, 100 μ g of lysates from tumor samples and normal breast control tissues was fractionated by SDS-PAGE and electrophoretically transferred to an Immobilon-P membrane (Millipore). TopBP1 was detected by either a monoclonal antibody (BD Transduction Laboratories) or a polyclonal antibody (Bethyl Laboratories).

Statistical analysis. Descriptive statistics were used to summarize patients' demographic and clinical characteristics by TopBP1 status (positive/negative). Means, standard deviations, and ranges were applied for continuous variables, and frequency and percentage were calculated for categorical variables. The differences of these characteristics between TopBP1-positive and -negative groups were compared using the *t* test, chi-square test, or Fisher's exact test. Survival time is defined as the number of months between date of initial diagnosis and date of death or last follow-up date. Time to disease progression is defined as the number of months between date of initial diagnosis and date of first disease relapse/progression or last follow-up date. We analyzed survival data using the Kaplan-Meier method. The difference of survival time by TopBP1 status was examined using the log-rank test. *P* values less than 0.05 were considered statistically significant. Statistical analyses were performed using SAS version 9.0 software (SAS Institute).

FIG. 4. A role of TopBP1 in the control of p53-induced apoptosis and G₁ arrest during DNA damage. (A) $p53^{+/+}$ or $p53^{-/-}$ HCT116 cells were infected with Ad-siGFP or Ad-siTopBP1 at a multiplicity of infection of 200. Twenty-four hours later, cells were either left untreated or treated with adriamycin at 5 μ M for 5 h. RNA was then extracted, and RT-PCR analysis was performed using primers specific for the selected p53 target genes or GAPDH. Left panels: semiquantitative RT-PCR. The "no RNA" lane represents a control RT-PCR without input RNA. An aliquot of the cell lysates was analyzed by Western blotting (left lower panels). Right panels: the changes of p53 target gene expression in $p53^{+/+}$ HCT116 cells were confirmed using quantitative real-time PCR. Results were normalized to GAPDH levels and are expressed relative to the expression of the gene in the siGFP control. The *P* values for the differences of gene expression between siGFP and siTopBP1 and between siGFP and siGFP-plus-adriamycin groups are all less than 0.01 (paired two-tailed *t* test). (B) $p53^{+/+}$ or $p53^{-/-}$ HCT116 cells were infected with Ad-siGFP or Ad-siTopBP1 at a multiplicity of infection of 200 as described for panel A. Twenty-four hours later, cells were labeled with BrdU for 1 hour and the incorporated BrdU was detected with fluorescein isothiocyanate-conjugated anti-BrdU antibody following the manufacturer's instructions (Becton Dickinson). The experiment was performed in triplicate. The *P* values are based on a paired two-tailed *t* test. (C) TopBP1 inhibits p53-induced apoptosis. A p53-null lung cancer cell line, H358, was infected with recombinant adenoviruses expressing empty vector (AdCMV), TopBP1, or/and p53 (multiplicity of infection, 400 for AdTopBP1 and 100 for Adp53). Two days later, cells were harvested for annexin-PE staining followed by flow cytometry or Western blot analysis. The data shown are the means $\pm$ standard errors of three independent experiments. The paired two-tailed *t* test *P* value is shown. (D) TopBP1 depletion promotes p53-dependent 5-FU-induced apoptosis. $p53^{+/+}$ HCT116 and $p53^{-/-}$ HCT116 cells were infected with adenovirus Ad-siGFP or Ad-siTopBP1 at a multiplicity of infection of 200. Twenty-four hours later, cells were either left untreated or treated with 5-FU (375 μ M) for 48 h. The cells were then harvested, stained with annexin V-PE and 7-AAD, and analyzed by flow cytometry. An aliquot of cell lysates was analyzed by Western blotting with TopBP1 and β -actin antibodies (upper panels).

FIG. 5. TopBP1 prevents p53 from binding to its target gene promoters. (A) Repression of p53 transcriptional activity by TopBP1 does not require Brg1/Brm or Akt phosphorylation. p53 transcriptional activity was measured by a p21 promoter-luciferase construct in a Brg1/Brm-deficient C33A cell line with or without reconstitution of Brg1 and Brm. p53 was cotransfected with either wild-type TopBP1 or TopBP1(S1159A) with or without Brg1 and Brm as indicated to C33A cells. Luciferase activity of transfected p53 was determined as induction relative to that of the empty-vector control. Each sample was assayed in triplicate. The experiments were repeated multiple times with consistent results. One-tenth of each sample was lysed with SDS lysis buffer for Western blot analysis (upper panels). (B and C) Nuclear extracts were prepared from $p53^{-/-}$ HCT116 cells infected with AdCMV (control extracts), Adp53, AdTopBP1, or Adp53 plus AdTopBP1 as described for Fig. 4C. p53 DNA-binding activity in extracts (6 μ g) was measured by EMSA. The p53-DNA complex (solid arrow) was specifically competed by untagged (cold) oligonucleotides containing wild-type p53-binding sites and supershifted by incubating the extracts with 0.2 μ g anti-p53 (PAb421). The supershifted PAb421-p53-DNA complex (open arrow) migrated further into the gel upon a longer run and could be competed by untagged oligonucleotides containing wild-type p53-binding sites (lower panel of panel B). Western blot analysis of the nuclear extracts is shown (lower panels of panel C). (D) Nuclear extracts (6 μ g) from Adp53-infected $p53^{-/-}$ HCT116 cells were incubated with 50 ng bovine serum albumin (BSA) or purified TopBP1 protein (produced from *E. coli*) for 3 h in the presence of 50 mM NaCl and then subjected to EMSA. (E and F) TopBP1 overexpression inhibits p53 promoter occupancy. $p53^{+/+}$ HCT116 cells were either left uninfected or infected with AdTopBP1 at a multiplicity of infection of 200 for 24 h. Some uninfected cells were treated with adriamycin (5 μ M) for 5 h. A ChIP assay was performed using antibodies against p53, TopBP1, and a control normal mouse immunoglobulin G (IgG) control, respectively, as indicated. Mock immunoprecipitations correspond to control reaction mixtures lacking antibodies. The precipitated DNA was amplified with primers derived from p21^{Cip1}, HDM2, NOXA, IGFBP3, E2F1, or β -actin promoter. The input represented 0.5% of the total amount of chromatin added to each immunoprecipitation reaction mixture. An aliquot of cell lysates was analyzed by Western blotting with p53 and TopBP1 antibodies (right panels). (F) The ChIP assay results for each promoter were quantified by real-time PCR.

FIG. 5—Continued.

RESULTS

TopBP1 interacts with p53 in vitro and in vivo. In a pilot microarray study (see Table S1 and Fig. S1 in the supplemental material), we observed induction of several known p53 target genes upon depletion of TopBP1. The observation prompted us to examine whether TopBP1 regulates p53. Indeed, endogenous TopBP1 interacted with p53 in the presence or absence of a radiomimetic agent, neocarzinostatin (NCS), in primary human fibroblasts and HEK293 cells (Fig. 1A). The interaction was also detected in the breast cancer cell line MCF7 by reciprocal coimmunoprecipitation (Fig. 1A, right panels). The interaction between overexpressed TopBP1 and endogenous

p53 was also seen in HEK293 cells (Fig. 1B). The increased coimmunoprecipitation during adriamycin or NCS treatment is likely due to induction of p53 and TopBP1 levels. A subcellular fractionation experiment in the colon cancer cell line HCT116 further showed an interaction in the nucleus but not in the cytosol (Fig. 1C). Since both p53 and TopBP1 can interact with DNA, the formation of their complex could indirectly result from DNA-protein interactions. To exclude this possibility, we pretreated cell extracts of MCF7 and HCT116 cells with ethidium bromide or DNase and then performed immunoprecipitation. We detected the interaction between endogenous TopBP1 and p53 in both cells even after treatment with ethidium bromide or DNase (Fig.

1D). These data indicate that the interaction is not mediated by DNA and suggest a direct physical interaction. We further verified their interaction in living cells by a bimolecular fluorescence complementation assay (21). YFP1-p53 (containing enhanced yellow fluorescent protein aa 1 to 158 fused with p53) interacted with YFP2-TopBP1 (enhanced yellow fluorescent protein aa 159 to 239 fused with TopBP1) within the nucleus but not with a control zipper domain fusion protein (Fig. 1E to G). The YFP1-zipper and YFP2-zipper proteins are localized to the nucleus (Fig. 1F) and thus can serve as proper negative controls. We also observed interaction between YFP1-p53 and YFP2-p53 or between YFP1-TopBP1 and YFP2-TopBP1, consistent with oligomerization of p53 (9) and TopBP1 (28). Lastly, we purified both p53 and TopBP1 proteins produced from *E. coli* and tested their interaction in vitro. Indeed, the interaction between p53 and TopBP1 is direct, since purified p53 and TopBP1 interacted with each other in a GST pull-down assay (Fig. 2). This interaction does not require Akt phosphorylation (Fig. 2A), suggesting that oligomerization of TopBP1 is not needed for p53 binding. We also performed domain-mapping analysis and mapped the interaction domains to BRCT78 of TopBP1 and the DBD of p53 (Fig. 2B to E).

A physiological role for TopBP1 in the control of endogenous p53 target gene expression and p53 activity. To test a physiological role for TopBP1 in the control of p53, we examined p53 target gene expression in TopBP1-depleted NIH 3T3 cells. Depletion of TopBP1 activated several classic p53 target genes including NOXA, BAX, GADD45, and p21 (Fig. 3A). Several other target genes that were identified in our microarray experiment, including MDM2, HIC1, CABCl, and IGFBP3, were also induced, confirming our microarray data from assays performed on U2OS cells (see Table S1 in the supplemental material). The expression of TopBP1 was then reconstituted to its physiological level by a recombinant adenovirus that expresses human TopBP1 transcript resistant to the siRNA against murine TopBP1 due to codon degeneracy. Reconstitution of TopBP1 levels did resuppress the expression of these genes (Fig. 3A). The changes of gene expression by TopBP1 siRNA or rescue were confirmed by quantitative real-time PCR (Fig. 3A, right panels). Interestingly, TopBP1 siRNA did not alter the level of 14-3-3 σ , a known p53 target gene (13). These results suggest a new physiological role for TopBP1 in the control of specific target genes of p53.

Binding of TopBP1 to the DBD of p53 and activation of p53 target gene expression by TopBP1 depletion suggested to us that TopBP1 might repress the transcriptional activity of p53. To test this hypothesis, we cotransfected TopBP1 and p53 expression vectors with a p53 activity reporter p21 promoter-luciferase plasmid in a p53-null cell line, Saos-2. Expression of TopBP1 significantly inhibited p53 transcriptional activity (Fig. 3B), without affecting the protein level of p53. Importantly, experiments using a series of TopBP1 deletion mutants demonstrated that the p53-binding domain BRCT78 was necessary and sufficient to repress p53 transcriptional activity (Fig. 3B). We performed the experiments using another p53 reporter assay (20) and obtained similar results (see Fig. S2 in the supplemental material). Taken together, the results of the binding and functional regulation analyses of these TopBP1 truncated mutants as summarized in Fig. 3C indicate that BRCT78 of TopBP1 is responsible for p53 binding and regu-

lation. The requirement of the interaction domain for p53 regulation also strongly suggests that TopBP1 regulates p53 through a direct physical interaction.

We also tested whether the regulation is specific to p53 or is general to other p53 family members, p63 and p73, since they share 60% amino acid identity in the DBD with p53. Despite multiple attempts, no interaction between TopBP1 and p73 α or p63 α (full-length form) could be detected when they were overexpressed (Fig. 3D). TopBP1 also failed to repress the transcriptional activities of p73 α and p63 α in a p21 promoter luciferase reporter assay (Fig. 3E). These results suggest that the regulation is specific to p53 but not p73 α or p63 α . It is worth noting that p53, p63, and p73 each have multiple isoforms (33). We have not tested the interaction of TopBP1 with other isoforms of p53, p63, and p73 and therefore cannot rule out their interactions.

A role for TopBP1 in the control of p53-induced apoptosis and G₁/S arrest during DNA damage. To further establish the p53 dependency of the gene regulation by TopBP1, we knocked down TopBP1 in a pair of HCT116 cell lines with either wild-type p53 (p53^{+/+}) or null p53 (p53^{-/-}) status (3). These cells were infected with Ad-siTopBP1 or a control virus, Ad-siGFP. Twenty-four hours later, some cells were subjected to a 5-h treatment with adriamycin before harvesting. As shown in Fig. 4A, TopBP1 knockdown did not alter p53 protein levels or induce p53 Ser15 phosphorylation but upregulated the expression of its target genes including p21, HDM2, IGFBP3, BAX, CABCl, PUMA, and GADD45 and enhanced their induction by adriamycin. This effect was dependent on p53 since it was not seen in p53^{-/-} HCT116 cells. The induction of p21 protein was also confirmed by Western blot analysis (Fig. 4A, lower panels). Consistent with the induction of p21, depletion of TopBP1 at the time of RNA harvest also coincided with the inhibition of bromodeoxyuridine (BrdU) incorporation in p53^{+/+} but not in p53^{-/-} HCT116 cells (Fig. 4B).

We considered the possibility that TopBP1 depletion might indirectly activate p53 through loss of chromosome integrity (22). In experiments where analysis was carried out within 24 to 48 h after introduction of TopBP1 siRNA into cells, TopBP1 siRNA did not induce p53 induction or phosphorylation, E2F1 induction, or Chk2 phosphorylation (27) (Fig. 4; see also Fig. S3 in the supplemental material). These results suggest that no significant DNA damage was associated with TopBP1 depletion in our transient assays that could be responsible for induction of p53 target genes. This assertion is further supported by the lack of induction of p53 protein and phosphorylation (Fig. 4A) and 14-3-3 σ transcript (Fig. 3A and 4A), all known to be induced upon DNA damage (13, 47), in the TopBP1-depleted cells where induction of p53 target genes was seen. These data indicate that induction of p53 target genes in these TopBP1-depletion experiments is not a consequence of general DNA damage response. To further investigate this, we analyzed the time course of p21 induction following TopBP1 depletion in Ad-siTopBP1-infected HCT116 cells. As shown in Fig. S4 in the supplemental material, induction of p21 immediately followed the decrease of TopBP1, which occurred around 14 h postinfection, and no evidence of induction of H2AX phosphorylation (as a marker of DNA damage) could be seen at that time. Phosphorylation of H2AX could be detected only much later (around 48 to 72 h after infection). In

the experiments carried out by Kim et al. (22), phosphorylation of H2AX was also observed at least 3 days after transfection of TopBP1 siRNA. Therefore, we conclude that the induction of p53 target genes by TopBP1 depletion reflects a direct regulation of p53 transcriptional activity by TopBP1. Taken together, we conclude that TopBP1 is required to control p53 transcriptional activity during normal growth and upon DNA damage.

The repression of p53 proapoptotic genes such as NOXA, PUMA, and BAX by TopBP1 suggested that TopBP1 could inhibit p53-mediated apoptosis. Indeed, expression of TopBP1 in a p53-null lung cancer cell line, H358, inhibited the apoptosis induced by infection of a recombinant adenovirus expressing p53 (Adp53) (Fig. 4C; see also Fig. S5 in the supplemental material). To examine the role of TopBP1 in the control of p53-induced apoptosis during chemotherapy, we again utilized *p53*^{+/+} and *p53*^{-/-} HCT116 cells for the experiments. We depleted TopBP1 in these cells and measured the apoptosis induced by 5-fluorouracil (5-FU), which has been shown to induce p53-dependent apoptosis in HCT116 cells (4). Knockdown of TopBP1 did enhance p53-dependent 5-FU-induced apoptosis (Fig. 4D). Thus, TopBP1 represses p53-mediated apoptosis during DNA damage.

TopBP1 binds to the DBD of p53 and inhibits p53 binding to promoters. TopBP1 regulates p53 probably through a mechanism distinct from that which regulates E2F1, since it involves only its seventh and eighth BRCT domains (as opposed to the sixth to eighth BRCT domains for E2F1 regulation) and does not require Akt phosphorylation (Fig. 2A and 3B). To determine the mechanism, we tested whether Brg1/Brm is required for TopBP1 to repress p53 transcriptional activity. We utilized a Brg1/Brm mutant cell line, C33A, and C33A cells in which Brg1 or Brm was reconstituted (27) to perform a p53 activity reporter assay. As shown in Fig. 5A, the regulation of p53 transcriptional activity by TopBP1, unlike the regulation of E2F1, was not affected by the status of Brg1/Brm. This result indicates that Brg1/Brm is not required for TopBP1 to repress p53. Consistent with the notion that Akt phosphorylation is also not involved, an Akt phosphorylation site mutant, TopBP1(S1159A), can repress p53 transcriptional activity as well as does wild-type TopBP1 (Fig. 5A).

TopBP1-BRCT78 directly interacts with the p53 DBD and is necessary and sufficient to repress p53 transcriptional activity on the p21 promoter (Fig. 2 and 3), suggesting that TopBP1 regulates p53 through a direct interaction. The interaction at the p53 DBD also suggests a potential mechanism for repressing p53 activity. 53BP1 also interacts with p53 through the BRCT repeat of 53BP1 and the DBD of p53 (7, 19). The 53BP1 BRCT repeats and the linker between the repeats bind p53 on a region that overlaps with the DNA-binding surface of p53 (7, 19); therefore, p53 cannot simultaneously bind to 53BP1 and a DNA fragment carrying a consensus p53 binding site (17). To test whether binding of TopBP1 to p53 directly inhibits its specific DNA-binding activity, we performed p53 nonisotopic EMSA with double-stranded oligonucleotides containing p53-binding sites. Nuclear extracts prepared from *p53*^{-/-} HCT116 cells infected with AdCMV do not contain specific DNA-binding activity. Upon infection with Adp53, the nuclear extracts exhibited specific binding activity toward a p21 promoter fragment containing p53-binding sites. The p53-

TABLE 1. Characteristics of patients and tumor samples with (positive) or without (negative) overexpression of TopBP1

Characteristic	Mean $\pm$ SD or n (%) for patients with TopBP1 status:		P value ^a
	Negative (n = 33)	Positive (n = 46)	
Age at diagnosis (yrs; range, 28–89)	52.7 $\pm$ 13.1	52.7 $\pm$ 12.1	0.989
Race			0.181
Black	6 (18.8)	17 (38.6)	
White	24 (75.0)	24 (54.5)	
Other	2 (6.3)	3 (6.8)	
Unknown	1	2	
Survival			0.043
Alive	29 (93.5)	31 (75.6)	
Dead	2 (6.5)	10 (24.4)	
Unknown	2	5	
Progression			0.009
No	29 (93.5)	28 (68.3)	
Yes	2 (6.5)	13 (31.7)	
Unknown	2	5	
Tumor grade ^b			0.007
0	1 (3.2)	0	
1	4 (12.9)	0	
2	15 (48.4)	15 (33.3)	
3	11 (35.5)	30 (66.7)	
Unknown	2	1	
ER status			0.032
Negative	5 (17.2)	17 (41.5)	
Positive	24 (82.8)	24 (58.5)	
Unknown	4	5	
PR status			0.900
Negative	13 (44.8)	19 (46.3)	
Positive	16 (55.2)	22 (53.7)	
Unknown	4	5	
Lymph node involvement			0.369
No	13 (41.9)	14 (31.8)	
Yes	18 (58.1)	30 (68.2)	
Unknown	2	2	
Stage ^c			0.237
0	1 (3.1)	0	
1	4 (12.5)	2 (4.4)	
2	15 (46.9)	20 (44.4)	
3	11 (34.4)	23 (51.1)	
4	1 (3.1)	0	
Unknown	1	1	
Her2/Neu status			0.826
Negative	21 (77.8)	32 (80.0)	
Positive	6 (22.2)	8 (20.0)	
Unknown	6	6	

^a P value of t test, chi-square test, or Fisher's exact test without including unknowns.

^b Test based on the difference of tumor grades 0 to 2 versus grade 3.

^c Test based on the difference of stages 0 to 2 versus stages 3 and 4.

DNA complex could be "supershifted" by a p53 antibody and competed by untagged competitive oligonucleotides containing wild-type p53-binding sites but not by oligonucleotides containing mutated p53-binding sites (Fig. 5B). Importantly,

TopBP1 overexpression — negative
- - - positive

A

Overall survival curves

TopBP1 overexpression

negative: n=31 (2 deaths)

positive: n=41 (10 deaths)

Log-rank p-value=0.003

B

Progression-free Survival Curves

TopBP1 overexpression

negative: n=31 (2 progression)

positive: n=41 (13 progression)

Log-rank p-value=0.001

C

Survival Curves for Stages 3-4

TopBP1 overexpression

negative: n=12 (2 deaths)

positive: n=20 (8 deaths)

Log-rank p-value=0.003

D

Survival Curves for lymph node involvement

TopBP1 overexpression

negative: n=18 (2 deaths)

positive: n=26 (8 deaths)

Log-rank p-value=0.002

E

Survival Curves for ER+ or PR+

TopBP1 overexpression

negative: n=24 (1 deaths)

positive: n=27 (6 deaths)

Log-rank p-value=0.012

FIG. 6.

FIG. 6—Continued.

coinfection of AdTopBP1 inhibited the specific DNA-binding activity of p53 (Fig. 5C). Moreover, incubation of the p53-containing extracts with purified TopBP1 protein prepared from *E. coli* also blocked its DNA-binding activity (Fig. 5D).

To further test whether binding of TopBP1 inhibits p53 binding to endogenous promoters, we performed a ChIP assay to examine the occupancy of TopBP1 and p53 on the promoters of p53-responsive genes. As shown in Fig. 5E and F, overexpression of TopBP1 inhibited the occupancy of p53 on HDM2, NOXA, p21, and IGFBP3 promoters. No significant occupancy of TopBP1 on these promoters could be detected. This is not because of inefficiency of the TopBP1 ChIP assay, since E2F1 promoter sequences were detected on the same TopBP1 ChIP samples. Together with the data from EMSA and domain-mapping studies, these results indicate that TopBP1 directly binds the DBD of p53 and inhibits its promoter-binding activity.

TopBP1 is frequently overexpressed in breast cancer, and its overexpression is associated with poor survival. TopBP1 is induced by E2F (27, 52), the activity of which is often upregulated in cancer and might lead to upregulation of TopBP1. We examined TopBP1 protein levels in a cohort of frozen breast cancer samples collected from 79 patients. A majority of these patients have either stage 2 (35 patients) or stage 3 (34 patients) disease (see Table S2 in the supplemental material). Among them, we also obtained matched normal breast tissues from 47 patients. The expression of TopBP1 in normal breast tissues was below the sensitivity of our Western blot analysis, such that we could detect only a low level of TopBP1 in 2 out of 47 normal breast tissues. In contrast, 46 out of 79 (58.2%) breast cancer samples contained detectable and higher levels of TopBP1 compared with normal breast tissues (examples shown in Fig. S6 in the supplemental material). Although this is a retrospective study, the patient characteristics (Table 1) between the TopBP1-negative (not overexpressing) and TopBP1-positive (overexpressing) patients are similar in the age at diagnosis, race, stage, lymph node involvement, progesterone receptor (PR) status, and Her2/Neu status. However, patients with overexpression of TopBP1 tend to have higher grades of breast cancer ($P = 0.007$) and negative estrogen receptor (ER) status ($P = 0.032$) compared with those without overexpression of TopBP1. In addition, patients with overexpression of TopBP1 in the tumors have significantly shorter overall survival time (log-rank, $P = 0.003$) and shorter progression-free survival time ($P = 0.001$) (Fig. 6A and B) than those without overexpression of TopBP1. The median survival time for patients without overexpression of TopBP1 is 165 months (95% confidence interval [CI] = 138, 192) and for those with overexpression of TopBP1 is 40 months (95% CI = 33, not achievable [N/A]). The median progression-free sur-

vival time for patients without overexpression of TopBP1 is 107 months (95% CI = N/A, N/A) and for those with overexpression of TopBP1 is 37 months (95% CI = 25, N/A). The survival difference between TopBP1-positive and -negative tumors remains statistically significant for patients with disease stage 3 to 4 ($P = 0.003$), lymph node involvement ($P = 0.002$), and ER-positive or PR-positive status ($P = 0.012$) (Fig. 6C and E). These data strongly suggest that TopBP1 may be an important prognostic marker for aggressive subgroups of breast cancers. Given the activity of TopBP1 in repressing E2F1 and p53 function as presented earlier, TopBP1 levels might affect patients' clinical outcomes.

Going and colleagues reported that TopBP1 is aberrantly localized to cytosol in some areas of certain breast cancer cases (out of 61 cases examined, 37 had predominantly nuclear staining, two had undetectable staining, 13 had a nuclear and nuclear-plus-cytoplasmic pattern, and nine had a nuclear-plus-cytoplasmic and purely cytoplasmic pattern [10]). We also performed immunohistochemical staining on 20 formalin-fixed paraffin-embedded breast tissue sections chosen randomly from the TopBP1-overexpressing group. All these sections showed a predominantly nuclear overexpression pattern of TopBP1 (Fig. 6G to K) compared with normal mammary epithelium (Fig. 6L). Although some cells have cytoplasmic staining in addition to intense nuclear staining, we have not observed exclusively cytoplasmic distribution in these sections. To ensure that a difference in antigen retrieval techniques did not result in a different apparent distribution of TopBP1, tissues from random cases were stained separately with antigen retrieval using pH 6.0 citrate (used in the study by Going et al. [10]) or pH 9.0 EDTA. The staining was less with citrate than with EDTA, but both methods demonstrated primarily nuclear patterns with little cytoplasmic staining (Fig. 6M).

TopBP1 overexpression represses p53 target gene expression and inhibits p53-dependent apoptosis and G₁/S arrest. Since TopBP1 overexpression is frequently seen in breast cancer, we tested whether TopBP1 has the capacity to repress p53 activities when it is expressed at a level comparable to that seen in breast cancer. We overexpressed TopBP1 in p53^{+/+} and p53^{-/-} HCT116 cells and then examined the p53 target genes and the response to adriamycin. The expression of TopBP1 was achieved by recombinant adenoviral infection to achieve a uniform gene transduction. As shown in Fig. 7A, TopBP1 overexpression inhibited the expression of p21, HDM2, IGFBP3, BAX, CABCl, PUMA, and GADD45 but not 14-3-3 σ during normal growth and upon adriamycin treatment. The level of TopBP1 overexpression in this experiment is close to that seen in primary breast cancer tissues (Fig. 7A, right lower panels). Inhibition of BAX and PUMA expression by TopBP1 suggests that TopBP1 might mitigate p53-dependent

FIG. 6. (A to E) Kaplan-Meier curves of patients' overall survival or progression-free survival. (F) Negative control (3% goat serum without TopBP1 antibody) for immunohistochemical staining of breast cancer section (sample identifier 9 [see Table S2 in the supplemental material]) shown at a magnification of $\times 400$. The same tissue was stained with TopBP1 antibody and is shown in panel G. (G to K) Representative pictures of the TopBP1 staining in breast cancer at a magnification of $\times 600$ (G to I and K) or $\times 250$ (J). The corresponding sample identifier for panels G and H is 9, that for panel I is 80T, that for panel J is 67T, and that for panel K is 25T. Open arrows indicate infiltrating carcinoma. Solid arrows indicate adjacent normal mammary epithelium. (L) TopBP1 staining in normal breast tissue at a magnification of $\times 400$ from a reduction mammoplasty. (M) Representative pictures of the TopBP1 staining at a magnification of $\times 600$ in breast cancer (sample identifier #80T) with antigen retrieval in sodium citrate pH 6.0 (the same condition as that used in the study by Going et al. [10]).

apoptosis induced by chemotherapeutic agents. Indeed, TopBP1 expression inhibited 5-FU-induced apoptosis in a p53-dependent manner (Fig. 7B; see also Fig. S7 in the supplemental material).

We also investigated whether TopBP1 overexpression might interfere with p53-dependent G₁/S arrest during DNA damage. It has been shown that a low dose of adriamycin can induce p53-dependent G₁/S arrest and p53-independent G₂/M arrest in *p53*^{+/+} HCT116 cells and lead to accumulation of 4N cells in *p53*^{-/-} HCT116 cells (4). It is worth noting that deletion of p53 or p21 in HCT116 cells leads to a decrease of G₁ fraction and an increase of G₂/M fraction after adriamycin treatment (4, 48). Utilizing a pair of isogenic *p53*^{+/+} and *p53*^{-/-} HCT116 cells, we tested whether TopBP1 could inhibit the p53-dependent G₁/S arrest. Indeed, TopBP1 overexpression decreased the degree of adriamycin-induced G₁/S arrest (Fig. 7C and D). In AdCMV-infected control cells, adriamycin treatment induced G₁ arrest in 66.4% of the cells. Upon TopBP1 overexpression, the adriamycin-induced G₁ arrest declined to 26.3%. In adriamycin-treated *p53*^{-/-} HCT116 cells, there was still about 10 to 15% of the 2N fraction. Similar to *p53*^{-/-} HCT116 cells, TopBP1 overexpression also led to an accumulation of the 4N fraction (from 27.1% to 63.4%) in adriamycin-treated cells. The degree of G₁/S arrest after adriamycin treatment was also measured by BrdU incorporation (Fig. 7E), which also shows abrogation of the adriamycin effect by TopBP1 overexpression.

Collectively, these data strongly support a role for TopBP1 in the regulation of p53-dependent cell cycle arrest and apoptosis upon DNA damage. We also demonstrate that TopBP1 overexpression in cancer cells may hamper normal function of p53.

DISCUSSION

We provide compelling evidence supporting a physiological role for TopBP1 in the control of p53 during normal growth and upon DNA damage. An interaction between endogenous TopBP1 and p53 has been shown in multiple cell types including primary human fibroblasts, HEK293, MCF7, and HCT116 cells. A role for TopBP1 in repressing the expression of endogenous p53 target genes is also demonstrated in U2OS, NIH 3T3, and HCT116 cells. We further show that the mechanism of this regulation is through an interaction between BRCT78 of TopBP1 and the p53 DBD, leading to inhibition of the promoter-binding activity of p53. These findings may provide a link between two major pathways in the control of cellular proliferation, Rb/E2F and p53 (Fig. 8A). TopBP1 is controlled by Rb/E2F and is induced when cells enter into S phase (27, 52) to participate in DNA replication and ATR activation. TopBP1 in turn regulates the transcriptional functions of p53 (this study), E2F1 (26–28), and Miz1 (14). Through the regulation of p53 and Miz1, TopBP1 represses p21 expression. TopBP1 also inhibits apoptosis by controlling p53 and E2F1. Thus, the transcriptional function of TopBP1 appears to play a critical role during G₁/S transition to ensure smooth entry to S phase.

The TopBP1-mediated control may have pathological implications in cancer. Dereglulation of Rb leads to activation of E2F activity, which in turn may cause overexpression of

TopBP1. This is supported by the observation that some TopBP1 overexpression is correlated with higher levels of E2F1 protein or transcript (see Fig. S6A and C in the supplemental material). Other mechanisms could also cause TopBP1 overexpression. For example, TopBP1 is regulated by Egr1 as well (43). As a result, the G₁/S checkpoint mediated by p53 and Miz1 and the apoptosis checkpoint mediated by p53 and E2F1 may be inhibited (Fig. 8B). Thus, deregulation of the Rb pathway may inhibit p53 functions through upregulation of TopBP1. This mechanism may be partly responsible for the high-grade phenotype and poor patient survival of TopBP1-overexpressing breast tumors.

Among the p53-interacting partners, ASPP1 and -2 are activated upon DNA damage. They then interact with the DBD of p53 and enhance p53's apoptotic capability by increasing selective binding of p53 to the promoters of proapoptotic genes, such as BAX, but not the promoters of cell cycle arrest genes such as p21 or regulatory genes such as MDM2 (37). Another p53 DBD-binding protein, Hzf, induces preferentially the expression of p53 target genes that block the cell cycle such as p21 and 14-3-3 σ (5). In contrast, TopBP1 binds to the DBD of p53 and represses the expression of many p53 target genes including BAX, PUMA, NOXA, GADD45, HIC1, CABCl, IGFBP3, p21, and MDM2 but not 14-3-3 σ . The lack of regulation of 14-3-3 σ by TopBP1 could be because 14-3-3 σ is also regulated by p63 and p73 (30, 50) and TopBP1 does not control p63 or p73 activities. The relative nonselectivity in target gene regulation is consistent with a model in which TopBP1 binds the p53 DBD and inhibits its DNA-binding activity. Also in agreement with a potential tumor suppressor role for ASPP and a potential oncogenic role for TopBP1 is the fact ASPP1 and -2 are frequently downregulated in cancer (41), whereas TopBP1 is overexpressed in more than 50% of human breast cancers.

The prevalence of TopBP1 overexpression in breast cancer and its association with poor clinical outcome underscore the importance of the pathway delineated in our studies. Analysis of three breast cancer microarray databases (8, 16, 46) with a total of 537 breast tumor samples via a data mining tool, ONCOMINE, shows a very significant and consistent association between high levels of TopBP1 RNA and high-grade tumors ($P = 2.2\text{E}-7$, $1.4\text{E}-11$, and $8.6\text{E}-5$, respectively). Analysis of a microarray database from 226 ER-positive and 69 ER-negative breast tumors (44) also shows a strong association between high TopBP1 expression and negative ER status ($P = 6.1\text{E}-8$). Moreover, when breast tumors are classified as either low genomic risk or high genomic risk based on their 76-gene prognostic signature (8), TopBP1 is expressed at higher levels in high-risk breast tumors ($n = 143$) than in low-risk tumors ($n = 55$) ($P = 8.3\text{E}-6$). Thus, several large sets of microarray data are in line with our cohort study by protein analysis. The association between TopBP1 overexpression and high-grade tumors was also observed in feline and canine mammary neoplasia by immunohistochemical staining (31, 32). Most TopBP1 staining is nuclear, but as the degree of malignancy increases, nuclear and cytoplasmic staining is observed in human, feline, and canine mammary tumors (10, 31, 32). Although purely cytoplasmic staining of TopBP1 was observed in some areas of certain breast cancer cases in the study by Goings et al. (10), nuclear plus cytoplasmic staining was also observed

FIG. 7. TopBP1 overexpression leads to inhibition of p53 function in cancer. (A) *p53*^{+/+} or *p53*^{-/-} HCT116 cells were infected with AdCMV or AdTopBP1 at a multiplicity of infection of 200. Twenty-four hours later, cells were either left untreated or treated with adriamycin at 5 μ M for 5 h. RNA was then extracted, and RT-PCR analysis was performed using primers specific for the selected p53 target genes or GAPDH. The "no RNA" lane represents a control RT-PCR mixture without input RNA. Right panels: the changes of p53 target gene expression in *p53*^{+/+} HCT116 cells were confirmed using quantitative real-time PCR. Results were normalized to GAPDH levels and are expressed relative to the expression of the gene in AdCMV control. The *P* values for the differences of gene expression between AdCMV and AdTopBP1 and between AdCMV and AdCMV-plus-adriamycin groups are all less than 0.05 (paired two-tailed *t* test). Lower panels: an aliquot of the cell lysates was analyzed by Western blotting, and the level of TopBP1 in HCT116 cells was compared with that in breast tumor samples. (B) TopBP1 inhibits p53-dependent 5-FU-induced apoptosis. *p53*^{+/+} HCT116 and *p53*^{-/-} HCT116 cells were infected with adenovirus AdCMV or AdTopBP1 at a multiplicity of infection of 200. Twenty-four hours later, cells were either left untreated or treated with 5-FU (375 μ M) for 48 h. The cells were then harvested, stained with annexin V-PE and 7-AAD, and analyzed by flow cytometry. An aliquot of cell lysates was analyzed by Western blotting with TopBP1 and β -actin antibodies (right panels). (C and D) TopBP1 reduces adriamycin-induced G₁ arrest. *p53*^{+/+} and *p53*^{-/-} HCT116 cells were infected with adenovirus AdCMV or AdTopBP1 at a multiplicity of infection of 200. Cells were then either left untreated or treated with adriamycin (Adr) (0.34 μ M) for 24 h. The cells were harvested, and the cell cycle profile was analyzed by propidium iodide flow cytometry. Shown are the means $\pm$ standard deviations of the fractions of each phase of the cell cycle from three independent infections. Western blot analysis of the lysates from these samples is shown in panel D. (E) *p53*^{+/+} and *p53*^{-/-} HCT116 cells were infected with adenovirus AdCMV or AdTopBP1 and then treated with adriamycin as described for panel C. The cells were then labeled with BrdU for 1 hour, and the incorporated BrdU was detected with fluorescein isothiocyanate-conjugated anti-BrdU antibody. The experiment was performed in triplicate. The *P* values are based on a paired two-tailed *t* test.

FIG. 8. An integrated model for the function of TopBP1 in the normal cell cycle and in cancer cells. (A) TopBP1 is induced by E2F during G₁/S transition. It participates in DNA replication and ATR activation. It also restricts p53 activity and E2F1-dependent apoptosis to ensure smooth progression of S phase. (B) TopBP1 is upregulated through deregulation of the pRb pathway or other mechanisms. Overexpression of TopBP1 in some cancer cells then may inhibit both p53 and E2F1 activities in checkpoint or apoptosis. In this manner, up-regulation of TopBP1 links pRb and p53 pathways in cancer.

in other areas of the same samples. Among our 20 cases, all have a predominantly nuclear pattern, and no purely cytoplasmic staining was observed. There was also no exclusively cytoplasmic pattern in feline and canine mammary tumors (31, 32). Since the majority of overexpressed TopBP1s in breast cancers are nuclear, TopBP1 could therefore affect the functions of transcription factors such as E2F1 and p53.

Interestingly, 11 out of 12 triple-negative (ER⁻ PR⁻ Her2/Neu⁻) tumors overexpress TopBP1 in the current series (compared with 29/54 in samples in which there is at least one of these receptors; $P = 0.0205$). This suggests that TopBP1 may be a valuable therapeutic target for triple-negative tumors for which there is no existing targeted therapy. The overexpressed TopBP1 in these breast cancers might lead to impairment of p53 function as demonstrated in HCT116 cells (Fig. 7). In fact, we analyzed the expression levels of TopBP1 and p53 target genes in primary breast tumors from several published microarray databases (34, 39, 40, 49, 53). TopBP1 expression levels inversely correlate with the expression levels of p53 target genes, such as p21, HDM2, HIC1, CABCl, PUMA, etc. For example, based on breast cancer microarray data obtained from invasive ductal carcinoma and lobular carcinoma (53), the Pearson correlation coefficient between TopBP1 and HIC1 is -0.54 ($P < 0.0001$, $n = 60$), that between TopBP1 and HDM2 is -0.34 ($P = 0.0109$, $n = 55$), that between TopBP1 and p21 is -0.317 ($P = 0.02$, $n = 53$), and that between TopBP1 and GADD45A is -0.327 ($P = 0.01$, $n = 60$). The Pearson correlation coefficient between TopBP1 and p21 from another breast cancer microarray database (34) is -0.67 ($P < 0.0001$, $n = 70$). Our study provides a potential mechanism for the inverse correlation between the levels of TopBP1 and p53 target genes in breast cancer.

The association between TopBP1 overexpression and aggressive breast cancer also has therapeutic implications. TopBP1 overexpression might contribute to treatment resistance in breast cancer therapy by way of repressing E2F1- and

p53-mediated apoptosis, both being required for adequate treatment response. This hypothesis will need to be tested in prospective clinical trials. Whether TopBP1 is an independent prognostic marker for aggressive breast cancer also deserves further investigation in other larger clinical trials. Our studies outline a strategy for a novel mechanism-based therapy where we can use specific TopBP1 inhibitors in patients whose breast cancer has high expression of TopBP1, with the goal to reactivate both E2F1- and p53-dependent apoptosis and augment therapeutic efficacy.

ACKNOWLEDGMENTS

We gratefully acknowledge the UAB Tissue Procurement Facility for providing breast tumor specimens, Xinbin Chen for Myc-p63 α and HA-p73 α expression constructs, Stephen W. Michnick for YFP1 and YFP2 vectors, and Bert Vogelstein for p53^{+/+} and p53^{-/-} HCT116 cells. We also thank Fannie Lin for valuable discussions and reagents and Piotr Kulesza for help in real-time quantitative PCR. We acknowledge critical reading of the manuscript by Jason Paik and Fannie Lin.

The work was supported by grants from the NIH (CA100857), the Department of Defense Breast Cancer Research Program (W81XWH-04-1-0442), and the Susan G. Komen Foundation (BCTR0600484) and a pilot grant from the UAB Cancer Center Structural Biology Program. W.-C.L. is a Leukemia and Lymphoma Society Scholar.

All authors disclose that they have no financial interests that will pose a conflict of interest regarding the article.

REFERENCES

- Aylon, Y., and M. Oren. 2007. Living with p53, dying of p53. *Cell* 130:597–600.
- Braithwaite, A. W., G. Del Sal, and X. Lu. 2006. Some p53-binding proteins that can function as arbiters of life and death. *Cell Death Differ.* 13:984–993.
- Bunz, F., A. Dutriaux, C. Lengauer, T. Waldman, S. Zhou, J. P. Brown, J. M. Sedivy, K. W. Kinzler, and B. Vogelstein. 1998. Requirement for p53 and p21 to sustain G2 arrest after DNA damage. *Science* 282:1497–1501.
- Bunz, F., P. M. Hwang, C. Torrance, T. Waldman, Y. Zhang, L. Dillehay, J. Williams, C. Lengauer, K. W. Kinzler, and B. Vogelstein. 1999. Disruption of p53 in human cancer cells alters the responses to therapeutic agents. *J. Clin. Invest.* 104:263–269.
- Das, S., L. Raj, B. Zhao, Y. Kimura, A. Bernstein, S. A. Aaronson, and S. W. Lee. 2007. Hsf1 determines cell survival upon genotoxic stress by modulating p53 transactivation. *Cell* 130:624–637.
- Delacroix, S., J. M. Wagner, M. Kobayashi, K. Yamamoto, and L. M. Karnitz. 2007. The Rad9-Hus1-Rad1 (9-1-1) clamp activates checkpoint signaling via TopBP1. *Genes Dev.* 21:1472–1477.
- Derbyshire, D. J., B. P. Basu, L. C. Serpell, W. S. Joo, T. Date, K. Iwabuchi, and A. J. Doherty. 2002. Crystal structure of human 53BP1 BRCT domains bound to p53 tumour suppressor. *EMBO J.* 21:3863–3872.
- Desmedt, C., F. Piette, S. Loi, Y. Wang, F. Lallemand, B. Haibe-Kains, G. Viale, M. Delorenzi, Y. Zhang, M. S. d'Assignies, J. Bergh, R. Lidereau, P. Ellis, A. L. Harris, J. G. Kljij, J. A. Foekens, F. Cardoso, M. J. Piccart, M. Buyse, and C. Sotiriou. 2007. Strong time dependence of the 76-gene prognostic signature for node-negative breast cancer patients in the TRANSBIG multicenter independent validation series. *Clin. Cancer Res.* 13:3207–3214.
- Friedman, P. N., X. Chen, J. Bargonetti, and C. Privet. 1993. The p53 protein is an unusually shaped tetramer that binds directly to DNA. *Proc. Natl. Acad. Sci. USA* 90:3319–3323.
- Going, J. J., C. Nixon, E. S. Dornan, W. Boner, M. M. Donaldson, and I. M. Morgan. 2007. Aberrant expression of TopBP1 in breast cancer. *Histopathology* 50:418–424.
- Hashimoto, Y., and H. Takisawa. 2003. Xenopus Cut5 is essential for a CDK-dependent process in the initiation of DNA replication. *EMBO J.* 22:2526–2535.
- He, T. C., S. Zhou, L. T. da Costa, J. Yu, K. W. Kinzler, and B. Vogelstein. 1998. A simplified system for generating recombinant adenoviruses. *Proc. Natl. Acad. Sci. USA* 95:2509–2514.
- Hermeking, H., C. Lengauer, K. Polyak, T. C. He, L. Zhang, S. Thiagalingam, K. W. Kinzler, and B. Vogelstein. 1997. 14-3-3 sigma is a p53-regulated inhibitor of G2/M progression. *Mol. Cell* 1:3–11.
- Herold, S., M. Wanzel, V. Beuger, C. Frohme, D. Beul, T. Hillukkala, J. Syvaaja, H. P. Saluz, F. Haenel, and M. Eilers. 2002. Negative regulation of the mammalian UV response by Myc through association with Miz-1. *Mol. Cell* 10:509–521.
- Horn, H. F., and K. H. Vousden. 2007. Coping with stress: multiple ways to activate p53. *Oncogene* 26:1306–1316.

16. Ivshina, A. V., J. George, O. Senko, B. Mow, T. C. Putti, J. Smeds, T. Lindahl, Y. Pawitan, P. Hall, H. Nordgren, J. E. Wong, E. T. Liu, J. Bergh, V. A. Kuznetsov, and L. D. Miller. 2006. Genetic reclassification of histologic grade delineates new clinical subtypes of breast cancer. *Cancer Res.* **66**: 10292–10301.
17. Iwabuchi, K., P. L. Bartel, B. Li, R. Marraccino, and S. Fields. 1994. Two cellular proteins that bind to wild-type but not mutant p53. *Proc. Natl. Acad. Sci. USA* **91**:6098–6102.
18. Jeon, Y., K. Y. Lee, M. J. Ko, Y. S. Lee, S. Kang, and D. S. Hwang. 2007. Human TopBP1 participates in cyclin E/CDK2 activation and preinitiation complex assembly during G1/S transition. *J. Biol. Chem.* **282**:14882–14890.
19. Joo, W. S., P. D. Jeffrey, S. B. Cantor, M. S. Finnin, D. M. Livingston, and N. P. Pavletich. 2002. Structure of the 53BP1 BRCT region bound to p53 and its comparison to the Brca1 BRCT structure. *Genes Dev.* **16**:583–593.
20. Kern, S. E., J. A. Pietenpol, S. Thiagalingam, A. Seymour, K. W. Kinzler, and B. Vogelstein. 1992. Oncogenic forms of p53 inhibit p53-regulated gene expression. *Science* **256**:827–830.
21. Kerppola, T. K. 2006. Visualization of molecular interactions by fluorescence complementation. *Nat. Rev. Mol. Cell Biol.* **7**:449–456.
22. Kim, J. E., S. A. McAvoy, D. I. Smith, and J. Chen. 2005. Human TopBP1 ensures genome integrity during normal S phase. *Mol. Cell. Biol.* **25**:10907–10915.
23. Kumagai, A., J. Lee, H. Y. Yoo, and W. G. Dunphy. 2006. TopBP1 activates the ATR-ATRIP complex. *Cell* **124**:943–955.
24. Leone, G., J. DeGregori, Z. Yan, L. Jakoi, S. Ishida, R. S. Williams, and J. R. Nevins. 1998. E2F3 activity is regulated during the cell cycle and is required for the induction of S phase. *Genes Dev.* **12**:2120–2130.
25. Lin, W. C., F. T. Lin, and J. R. Nevins. 2001. Selective induction of E2F1 in response to DNA damage, mediated by ATM-dependent phosphorylation. *Genes Dev.* **15**:1833–1844.
26. Liu, K., F. T. Lin, J. M. Ruppert, and W. C. Lin. 2003. Regulation of E2F1 by BRCT-domain containing protein TopBP1. *Mol. Cell. Biol.* **23**:3287–3304.
27. Liu, K., Y. Luo, F. T. Lin, and W. C. Lin. 2004. TopBP1 recruits Brg1/Brm to repress E2F1-induced apoptosis, a novel pRb-independent and E2F1-specific control for cell survival. *Genes Dev.* **18**:673–686.
28. Liu, K., J. C. Paik, B. Wang, F. T. Lin, and W. C. Lin. 2006. Regulation of TopBP1 oligomerization by Akt/PKB for cell survival. *EMBO J.* **25**:4795–4807.
29. Makinemi, M., T. Hillukkala, J. Tuusa, K. Reini, M. Vaara, D. Huang, H. Pospiech, I. Majuri, T. Westerling, T. P. Makela, and J. E. Syvaaja. 2001. BRCT domain-containing protein TopBP1 functions in DNA replication and damage response. *J. Biol. Chem.* **276**:30399–30406.
30. Melino, G., X. Lu, M. Gasco, T. Crook, and R. A. Knight. 2003. Functional regulation of p73 and p63: development and cancer. *Trends Biochem. Sci.* **28**:663–670.
31. Morris, J. S., C. Nixon, A. Bruck, L. Nasir, I. M. Morgan, and A. W. Philbey. 2008. Immunohistochemical expression of TopBP1 in feline mammary neoplasia in relation to histological grade, Ki67, ERalpha and p53. *Vet. J.* **175**:218–226.
32. Morris, J. S., C. Nixon, O. J. King, I. M. Morgan, and A. W. Philbey. 2009. Expression of TopBP1 in canine mammary neoplasia in relation to histological type, Ki67, ERalpha and p53. *Vet. J.* **179**:422–429.
33. Murray-Zmijewski, F., D. P. Lane, and J. C. Bourdon. 2006. p53/p63/p73 isoforms: an orchestra of isoforms to harmonise cell differentiation and response to stress. *Cell Death Differ.* **13**:962–972.
34. Perou, C. M., T. Sorlie, M. B. Eisen, M. van de Rijn, S. S. Jeffrey, C. A. Rees, J. R. Pollack, D. T. Ross, H. Johnsen, L. A. Akslen, O. Fluge, A. Pergamenschikov, C. Williams, S. X. Zhu, P. E. Lonning, A. L. Borresen-Dale, P. O. Brown, and D. Botstein. 2000. Molecular portraits of human breast tumours. *Nature* **406**:747–752.
35. Remy, I., A. Montmarquette, and S. W. Michnick. 2004. PKB/Akt modulates TGF-beta signalling through a direct interaction with Smad3. *Nat. Cell Biol.* **6**:358–365.
36. Rozan, L. M., and W. S. El-Deiry. 2007. p53 downstream target genes and tumor suppression: a classical view in evolution. *Cell Death Differ.* **14**:3–9.
37. Samuels-Lev, Y., D. J. O'Connor, D. Bergamaschi, G. Trigiante, J. K. Hsieh, S. Zhong, I. Campargue, L. Naumovski, T. Crook, and X. Lu. 2001. ASPP proteins specifically stimulate the apoptotic function of p53. *Mol. Cell* **8**:781–794.
38. Sjøttem, E., C. Rekdal, G. Svineng, S. S. Johnsen, H. Klenow, R. D. Uglehus, and T. Johansen. 2007. The ePHD protein SPBP interacts with TopBP1 and together they co-operate to stimulate Ets1-mediated transcription. *Nucleic Acids Res.* **35**:6648–6662.
39. Sorlie, T., C. M. Perou, R. Tibshirani, T. Aas, S. Geisler, H. Johnsen, T. Hastie, M. B. Eisen, M. van de Rijn, S. S. Jeffrey, T. Thorsen, H. Quist, J. C. Matese, P. O. Brown, D. Botstein, P. Eystein Lonning, and A. L. Borresen-Dale. 2001. Gene expression patterns of breast carcinomas distinguish tumor subclasses with clinical implications. *Proc. Natl. Acad. Sci. USA* **98**:10869–10874.
40. Sorlie, T., R. Tibshirani, J. Parker, T. Hastie, J. S. Marron, A. Nobel, S. Deng, H. Johnsen, R. Pesich, S. Geisler, J. Demeter, C. M. Perou, P. E. Lonning, P. O. Brown, A. L. Borresen-Dale, and D. Botstein. 2003. Repeated observation of breast tumor subtypes in independent gene expression data sets. *Proc. Natl. Acad. Sci. USA* **100**:8418–8423.
41. Sullivan, A., and X. Lu. 2007. ASPP: a new family of oncogenes and tumour suppressor genes. *Br. J. Cancer* **96**:196–200.
42. Takahashi, Y., J. B. Rayman, and B. D. Dynlacht. 2000. Analysis of promoter binding by the E2F and pRB families in vivo: distinct E2F proteins mediate activation and repression. *Genes Dev.* **14**:804–816.
43. Usskilat, C., C. Skerka, H. P. Saluz, and F. Hanel. 2006. The transcription factor Egr-1 is a regulator of the human TopBP1 gene. *Gene* **380**:144–150.
44. van de Vijver, M. J., Y. D. He, L. J. van't Veer, H. Dai, A. A. Hart, D. W. Voskuil, G. J. Schreiber, J. L. Peterse, C. Roberts, M. J. Marton, M. Parrish, D. Atsma, A. Witteveen, A. Glas, L. Delahaye, T. van der Velde, H. Bartelink, S. Rodenhuis, E. T. Rutgers, S. H. Friend, and R. Bernards. 2002. A gene-expression signature as a predictor of survival in breast cancer. *N. Engl. J. Med.* **347**:1999–2009.
45. Van Hatten, R. A., A. V. Tutter, A. H. Holway, A. M. Khederian, J. C. Walter, and W. M. Michael. 2002. The Xenopus Xmus101 protein is required for the recruitment of Cdc45 to origins of DNA replication. *J. Cell Biol.* **159**:541–547.
46. van 't Veer, L. J., H. Dai, M. J. van de Vijver, Y. D. He, A. A. Hart, M. Mao, H. L. Peterse, K. van der Kooy, M. J. Marton, A. T. Witteveen, G. J. Schreiber, R. M. Kerkhoven, C. Roberts, P. S. Linsley, R. Bernards, and S. H. Friend. 2002. Gene expression profiling predicts clinical outcome of breast cancer. *Nature* **415**:530–536.
47. Vogelstein, B., D. Lane, and A. J. Levine. 2000. Surfing the p53 network. *Nature* **408**:307–310.
48. Waldman, T., K. W. Kinzler, and B. Vogelstein. 1995. p21 is necessary for the p53-mediated G1 arrest in human cancer cells. *Cancer Res.* **55**:5187–5190.
49. West, R. B., D. S. Nuyten, S. Subramanian, T. O. Nielsen, C. L. Corless, B. P. Rubin, K. Montgomery, S. Zhu, R. Patel, T. Hernandez-Boussard, J. R. Goldblum, P. O. Brown, M. van de Vijver, and M. van de Rijn. 2005. Determination of stromal signatures in breast carcinoma. *PLoS Biol.* **3**:e187.
50. Westfall, M. D., D. J. Mays, J. C. Snizek, and J. A. Pietenpol. 2003. The ΔNp63α phosphoprotein binds the p21 and 14-3-3σ promoters in vivo and has transcriptional repressor activity that is reduced by Hay-Wells syndrome-derived mutations. *Mol. Cell. Biol.* **23**:2264–2276.
51. Wright, R. H., E. S. Dornan, M. M. Donaldson, and I. M. Morgan. 2006. TopBP1 contains a transcriptional activation domain suppressed by two adjacent BRCT domains. *Biochem. J.* **400**:573–582.
52. Yoshida, K., and I. Inoue. 2004. Expression of MCM10 and TopBP1 is regulated by cell proliferation and UV irradiation via the E2F transcription factor. *Oncogene* **23**:6250–6260.
53. Zhao, H., A. Langerod, Y. Ji, K. W. Nowels, J. M. Nesland, R. Tibshirani, I. K. Bukholm, R. Karesen, D. Botstein, A. L. Borresen-Dale, and S. S. Jeffrey. 2004. Different gene expression patterns in invasive lobular and ductal carcinomas of the breast. *Mol. Biol. Cell* **15**:2523–2536.