Published in final edited form as:

Tetrahedron Lett. 2009 March 18; 50(11): 1161–1163. doi:10.1016/j.tetlet.2008.12.096.

Total Synthesis of Tovophyllin B

Valer Jeso and K.C. Nicolaou

Department of Chemistry and The Skaggs Institute for Chemical Biology, The Scripps Research Institute, 10550 North Torrey Pines Road, La Jolla, California 92037, and Department of Chemistry and Biochemistry, University of California, San Diego, 9500 Gilman Drive, La Jolla, California 92037, USA

Abstract

The first total synthesis of tovophyllin B (**2**), an antimicrobial xanthone derived from mangosteen, has been accomplished through a convergent strategy from building blocks **6** and **7** involving lithium-mediated coupling, dehydrative cyclization, and 6π electrocyclization as key steps.

Tovophyllin B, a prenylated pentacyclic xanthone, was first isolated in 1972 from the wood of *Tovomita macrophylla* by de Oliveira et al.¹ The originally proposed structure (**1**; Figure 1) was corrected in 1975 on the basis of chemical and spectroscopic data to the currently accepted structure (**2**; Figure 1) in which one of the phenolic groups is H-bonded to the carbonyl moiety of the molecule.² Recent investigations of the antituberculosis potential of the fruit hulls of mangosteen (*Garcinia mangostana*), traditionally used in Thai folk medicine for alleviation of a number of maladies, led to re-isolation of this natural product and revealed that tovophyllin B (**2**) possesses a significant inhibitory activity against *Mycobacterium tuberculosis* (MIC = 25 μ g/mL).³ Therefore, as a lead compound for drug discovery, the total synthesis of tovophyllin B and its analogs is deemed important. In this letter we report a short and efficient total synthesis of this molecule that may also serve to prepare designed analogs.

Scheme 1 shows the retrosynthetic analysis of the molecule of tovophyllin B (**2**) in which a 6π electrocyclization (a) plays the role of casting the final heterocyclic ring in the synthetic direction (**3**→**2**). An aldol/dehydration sequence (b) then traces **3**, expected to be a transient intermediate, back to xanthone **4** and prenal (**4'**). Disconnection of the indicated carbon-oxygen bond within xanthone **4** through a retro dehydrative cyclization leads to bisaryl ketone **5**, whose disassembly through a retro lithium-mediated coupling as shown reveals benzaldehyde **6** and benzopyran derivative **7** as the required building blocks for the projected synthesis.

Scheme 2 summarizes the synthesis of advanced intermediate bisaryl ketone **5**. Thus, *O*-propargylation⁴ of the readily available phenol **8** (prepared from 2,4-dihydroxybenzaldehyde in three steps)⁵ with methyl 2-methyl-3-yn-2-yl carbonate (**9**) in the presence of DBU and catalytic amounts of CuCl₂ proceeded smoothly to afford 1,1-dimethylpropargyl ether **10**, which, upon heating in toluene at 140 °C, underwent Claisen rearrangement, leading to 2,2-dimethylchromene **11**, in 74% overall yield for the two steps.⁶

Regioselective lithiation of **11** facilitated by the two OMOM groups (*n*BuLi, THF, 25 °C), followed by quenching of the resulting lithiated species with prenyl bromide furnished

Correspondence to: K.C. Nicolaou.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

prenylated chromene **7** in 86% yield. Pleasantly, the generation of the lithiated species **12** from **7** proceeded smoothly on exposure of the later to *n*BuLi in THF at 25 °C and was essentially complete within 15 minutes (ca. 95% deuterium incorporation upon quenching with D₂O). Addition of bis-benzyl-MOM protected benzaldehyde derivative **6** (prepared from phloroglucinol carboxaldehyde in two steps)⁷ to lithiated species **12** at –78 °C furnished coupling product **13** (58% yield),⁸ whose hydroxy group was oxidized with Dess-Martin periodinane to afford ketone **14**⁹ in 80% yield. Initial attempts to selectively remove the benzyl groups from the latter compound with hydrogen and a variety of catalysts failed primarily due to the interference of the olefinic bonds residing within the molecule. This problem was, however, solved through the employment of a palladium-catalyzed transfer hydrogenation protocol utilizing a triethylammonium formate/formic acid buffer system.¹⁰ Under these conditions the two benzyl groups were cleanly cleaved from **14** to form the corresponding diphenol (**15**, 89% yield), from which the MOM groups were removed by exposure to CSA in MeOH at ambient temperature, leading to the desired tricyclic precursor **5**.

The final steps of the synthesis of tovophyllin B (**2**) are shown in Scheme 3. Thus, cycloetherification of **5** under the influence of silica gel (impregnation of the substrate on a silica gel plate, followed by elution with EtOAc)¹¹ furnished, through dehydration, xanthone **4**⁹ in 65% yield. The missing 2,2-dimethylchromene ring of the growing molecule was finally forged through a cascade sequence involving CaO-induced aldol-type reaction¹² of xanthone **4** with prenal (**4'**) that proceeded via intermediates **16** and **3** to afford the target molecule (**2**) in 55% overall yield as shown in Scheme 3. Synthetic tovophyllin B (**2**) exhibited identical physical properties to those reported for the naturally derived material.^{1,2}

The described chemistry demonstrates the power of cascade reactions in total synthesis and provides a practical route to tovophyllin B (**2**) and related compounds, including designed analogs for biological investigations.

Acknowledgments

We thank Drs. Dee H. Huang and Gary Siuzdak for NMR spectroscopic and mass spectrometric assistance, respectively. Financial support for this work was provided by the National Institutes of Health (USA) (AI 055475-08) and the Skaggs Institute for Chemical Biology.

References and notes

- de Oliveira WG, Gottlieb OR, Mesquita AAL. *Phytochemistry* 1972;11:3323–3325.
- Mesquita AAL, de Oliveira WG, Neiva RMT. *Phytochemistry* 1975;14:803–806.
- Suksamrarn S, Suwannapoch N, Phakhodee W, Thanuhiranlert J, Ratananukul P, Chimnoi N, Suksamrarn A. *Chem Pharm Bull* 2003;51:857–859. [PubMed: 12843596]
- Godfrey JD, Mueller RH, Sedergran TC, Soundararajan N, Colandrea VJ. *Tet Lett* 1994;35:6405–6408.
- Miyake M, Hanaoka Y, Fujimoto Y, Sato Y, Taketomo N, Yokota I, Yoshiyama Y. *Heterocycles* 1996;43:665–674.
- a) Nicolaou KC, Sasmal PK, Xu H, Namoto K, Ritzen A. *Angew Chem Int Ed* 2003;42:4225–4229. b) Quillinan AJ, Scheinmann F. *J Chem Soc Perkin Trans 1* 1972:1382–1387.
- Anderson JC, Headley C, Stapleton PD, Taylor PW. *Tetrahedron* 2005;61:7703–7711. [PubMed: 19330042]
- a) Nicolaou KC, Xu H, Wartmann M. *Angew Chem Int Ed* 2005;44:756–761. b) Iikubo K, Ishikawa Y, Ando N, Umezawa K, Nishiyama S. *Tet Lett* 2002;43:291–293. c) Nicolaou KC, Li J. *Angew Chem Int Ed* 2001;40:4264–4268.
- Physical properties of selected compounds. **14**: IR (film): $\nu_{\max}$ 3065, 2974, 2928, 1650, 1600, 1582, 1430, 1322, 1206, 1152, 1112, 1067, 926, 843, 731, 696 cm⁻¹; ¹H NMR (400 MHz, CDCl₃) δ 7.40–7.18 (m, 10 H), 6.43 (d, *J* = 9.9 Hz, 1 H), 6.42 (s, 1 H), 6.28 (d, *J* = 1.8 Hz, 1 H), 5.53 (d, *J* = 9.9 Hz,

1 H), 5.18 (t, $J = 6.6$ Hz, 1 H), 5.12 (s, 2 H), 5.04 (s, 2 H), 4.96 (s, 2 H), 4.94 (s, 2 H), 4.69 (s, 2 H), 3.56 (s, 3 H), 3.41 (d, $J = 6.3$ Hz, 2 H), 3.31 (s, 3 H), 3.29 (s, 3 H), 1.71 (s, 3 H), 1.63 (s, 3 H), 1.41 (s, 6 H); ^{13}C NMR (100 MHz, CDCl_3) δ 192.99, 162.03, 159.07, 158.08, 146.98, 144.88, 141.80, 136.12, 136.09, 131.03, 130.36, 129.98, 129.45, 128.54, 128.34, 128.14, 127.61, 127.58, 126.84, 120.30, 119.09, 115.39, 101.27, 98.59, 94.72, 94.63, 94.28, 75.41, 70.39, 70.06, 57.37, 57.23, 55.99, 27.50, 25.58, 24.01, 17.89; HRMS(ESI-TOF) m/z calcd for $\text{C}_{43}\text{H}_{49}\text{O}_{10}$ ($\text{M} + \text{H}^+$) 725.3320, found 725.3320. Compound **4**: IR (film): ν_{max} 3356, 2973, 2925, 2876, 1650, 1612, 1570, 1431, 1321, 1293, 1253, 1167, 1138, 1084, 1031, 966, 893, 830, 813, 768 cm^{-1} ; ^1H NMR (400 MHz, $\text{MeOH}-d_4$) δ 7.92 (d, $J = 10$ Hz, 6.22 (s, 1 H), 6.09 (s, 1 H), 5.76 (d, $J = 10$ Hz, 1 H), 5.23 (t, $J = 7.6$ Hz, 1 H), 3.51 (d, $J = 7.2$ Hz, 2 H), 1.87 (s, 3 H), 1.67 (s, 3 H), 1.45 (s, 6 H); ^{13}C NMR (100 MHz, $\text{MeOH}-d_4$) δ 182.54, 164.73, 163.37, 163.36, 157.33, 151.05, 150.22, 137.61, 131.59, 131.40, 131.38, 121.53, 120.84, 117.50, 177.49, 115.60, 107.35, 102.77, 97.49, 92.82, 75.75, 26.05, 24.74, 22.21, 16.96; HRMS(ESI-TOF) m/z calcd for $\text{C}_{23}\text{H}_{23}\text{O}_6$ ($\text{M} + \text{H}^+$) 395.1489, found 395.1491.

10. Tani K, Ono N, Okamoto S, Sato F. *J Chem Soc Chem Commun* 1993:386–387.
11. Hamada M, Iikubo K, Ishikawa Y, Ikeda A, Umezawa K, Nishiyama S. *Bioorg Med Chem Lett* 2003;13:3151–3153. [PubMed: 12951083]
12. a) Nicolaou KC, Sasmal PK, Xu H. *J Am Chem Soc* 2004;126:5493–5501. [PubMed: 15113221] b) Tisdale EJ, Slobodov I, Theodorakis EA. *Proc Natl Acad Sci U S A* 2004;101:12030–12035. [PubMed: 15210986] c) Saimoto H, Yoshida K, Murakami T, Morimoto M, Sashiwa H, Shigemasa Y. *J Org Chem* 1996;61:6768–6769. [PubMed: 11667564]

1: tovophyllin B
[original structure]

2: tovophyllin B
[revised structure]

Figure 1.
Originally proposed (**1**) and revised (**2**) structures of tovophyllin B.

Scheme 1.

Retrosynthetic analysis of tophoyllin B (**2**): a) 6π electrocyclization; b) aldol/dehydration.

Scheme 2.

Synthesis of bisaryl ketone **5**. Reagents and conditions: a) **9** (3.0 equiv), DBU (3.0 equiv), CuCl₂ (0.01 equiv), MeCN, 0 °C, 12 h; b) toluene, 140 °C, 48 h, 74% (over two steps); c) *n*BuLi (1.3 equiv), THF, 25 °C, 15 min; then prenyl bromide (1.5 equiv), THF, 25 °C, 2 h, 86%; d) *n*BuLi (1.5 equiv), THF, 25 °C, 15 min; e) **6** (1.5 equiv), THF, −78 °C, 15 h, 58%; f) DMP (1.4 equiv), NaHCO₃ (5.0 equiv), CH₂Cl₂, 25 °C, 2 h, 80%; g) 10% Pd/C (0.34 equiv), Et₃N (1.0 equiv), HCO₂H (32 equiv), acetone, 25 °C, 1.5 h, 89%; h) CSA (5.0 equiv), MeOH, 25 °C, 8 h.

Scheme 3.

Completion of the total synthesis of tophophyllin B (2). Reagents and conditions: a) impregnation on a silica gel plate (1.5 h); then elution with EtOAc, 65%, b) CaO (7.5 equiv), prenal (4') (15 equiv), MeOH, 25 °C, 16 h, 55%.