Published in final edited form as:

Nat Prod Rep. 2009 May ; 26(5): 661–680. doi:10.1039/b811227b.

Synthetic Efforts toward Cyathane Diterpenoid Natural Products

John A. Enquist Jr. and Brian M. Stoltz^{*}

The Arnold and Mabel Beckman Laboratories of Chemical Synthesis Division of Chemistry and Chemical Engineering, California Institute of Technology 1200 East California Boulevard, MC 164-30, Pasadena, California, 91125, USA

1. Introduction

This review is intended to present an overview of synthetic efforts published toward the cyathane diterpenoid natural products. The work summarized here represents an update to the previous review on this topic which was reported in 2000, and as such will only cover cyathane core syntheses described in the literature past this date.¹ For the purpose of comprehensive review and comparison, however, all known completed total syntheses will be detailed herein. The focus of this review is centered upon the ring-forming and stereogenic strategies employed within the published strategies toward cyathane natural products.

2. Overview of the Cyathane Diterpenoids

The cyathane diterpene natural products are isolated from a diverse variety of fungi, sponges, and fruiting plants. However, despite their myriad natural sources, are all unified by the presence of a characteristic 5-6-7 tricyclic fused core structure (**1**, Figure 1). Within this class of natural products can be found the cyathins, the allocyathins, the erinacines, the sarcodonins, the scabronines, the striatals, the cyanthiwiggins, and the cyaftrins, all of which display the conserved carbon scaffold of the cyathane skeleton (**1**). Of the twenty carbons that comprise this cyathane framework, the C(6) and C(9) carbons present all-carbon quaternary stereocenters, which bear angular methyl groups at the points of ring fusion. Almost all of the compounds within the cyathane class display these methyl groups with an anti relative stereochemical relationship, though the cyanthiwigin natural products possess a syn arrangement. The cyathane core structure is additionally characterized by the presence of an isopropyl side chain at C(3) and an exocyclic carbon atom connected to C(12)

2.1 Isolation

In 1970, Ayer and Brodie published a report in which an extract from the bird's nest fungus *Cyathus helenae* was scrutinized to better understand its antimicrobial activity. Though no full structural assignment was made for any compound within the extract, the active components of the mixture were separated via chromatography. After isolation and elemental analysis, the compounds responsible for the observed antimicrobial activity were named (without structural elucidation) as cyathin A₃, A₄, B₃, B₄, C₅, and allocyathin A₄.² The first fully characterized cyathane diterpene natural products were subsequently reported by Ayer and coworkers in 1972, when the substance previously identified as cyathin A₃ was found to be a mixture of isomeric compounds, which they named cyathin A₃ (**2**) and allocyathin B₃ (**3**).³ Numerous other cyathin and allocyathin natural products were discovered subsequent to this report, including allocyathin B₂ (**4**),⁴ a cyathane diterpenoid which has since become the focus of numerous synthetic studies.

Following the identification of the primary cyathin diterpenoids, several structurally related compounds were isolated from additional natural sources. It was discovered that the fruiting bodies of *Hericium erinaceum* contained a number of glycosylated allocyathin B₂ analogues, which were eventually named the erinacines. Among the compounds obtained from *Hericium erinaceum* were erinacine A (**5**), erinacine B (**6**), and the more structurally complex erinacine E (**7**).⁵

In 1989 Nakayama and coworkers published the isolation and structural assignment of eight new cyathane molecules from the fungus *Sarcodon scabrosus* which they named the sarcodonins A–H.⁶ These compounds possess the conserved 5-6-7 tricyclic core found in all other cyathane natural products, but are distinct in that they display additional oxidation at the C(19) position, such as is observed with sarcodonin G (**8**). An additional class of cyathane diterpene compounds was later identified from the same *Sarcodon scabrosus* fungus by Oshima in 1998, when isolation and characterization of scabronine A–F were disclosed. Six years later, Liu *et al.* disclosed the isolation of two additional scabronine molecules from the same source, including scabronine G (**9**).⁷ Structural elucidation of these natural products revealed the presence of a carboxyl group at C(17), a feature which marks the scabronines as distinct from the remainder of the cyathanes.

In 1992, Kashman and coworkers published a report detailing the isolation and characterization of the first cyanthiwigin molecules. Initially isolated from the marine sponge *Epipolasis reiswigi*, cyanthiwigins A–D were fully characterized and assigned absolute configuration via NMR, X-ray, and Mosher ester analysis.⁸ A decade later the laboratories of Hamann isolated the same four cyanthiwigins, plus an additional 23 compounds of this class, from the Jamaican sea sponge *Myrmekioderma styx*. This isolation included the products cyanthiwigin F (**10**) and cyanthiwigin U (**11**). In years subsequent to their initial report, Hamann and coworkers have detailed the isolation and characterization of the additional cyanthiwigin molecules AB–AG, including the structurally unique spirocyclic cyanthiwigin AC (**12**).⁹

2.2 Bioactivity

The biological activities of the cyathane natural products vary widely among the different molecules within the class. Many of these diterpene compounds possess antibiotic or antimicrobial activity, such as the previously discussed cyathin A₃ (**2**) and allocyathin B₃ (**3**). Indeed, almost all of the subcategories of cyathane molecules display mild activity in this regard.^{2,4,9} In addition to serving an antibiotic function, some members of the cyanthiwigin compounds have displayed limited cytotoxic activity against human primary tumor cells, as well as P388 murine leukemia cells and A549 lung tumor cells.^{9,10}

However, the most significant biological activity reported among these diterpenoid molecules is their powerful ability to encourage the synthesis of Nerve Growth Factor (NGF). Both the erinacine and the scabronine natural products have displayed considerable potency in the stimulation of NGF synthesis,^{5,7b} a capacity that implicates their potential as therapeutic agents to treat neurodegenerative ailments such as Alzheimer's or Parkinson's disease.¹¹

2.3 Biosynthesis

The details of cyathane diterpenoid biosynthesis have been covered in Wright's review, and as such will only be briefly summarized here.¹ Subsequent to his isolation of the parent cyathins, Ayer conducted an in-depth study to scrutinize the biosynthetic origin of the cyathane diterpenoid core. By growing *Cyathus earlei* in the presence of ¹³C-labelled sodium acetate, Ayer was able to isolate and examine the cyathin molecules produced by these fungal bodies. Analysis of these compounds via ¹³C NMR allowed Ayer to conclude that the biosynthetic

pathway toward the cyathane core tricycle involves cascade cyclization and subsequent rearrangement of geranylgeranyl phosphate.¹²

3. Strategy Summary

Since the isolation of the first cyathane natural products in the early 1970s, numerous research groups have endeavoured to synthesize these compounds. A multitude of differing strategies have been documented in the literature, and many of these approaches are presented here in schematic form. The synthetic efforts reported to date can be roughly categorized by their key transformations, which are further divided as either transition metal-mediated reactions (Scheme 2), or alkylation / aldol reactions (Scheme 3).

A majority of the cyathane syntheses published to date leverage various metal-catalyzed transformations to accomplish difficult or complicated ring-forming reactions. For example, the strategy employed by Trost *et al.* relies upon Ru-catalyzed cycloisomerization to close the central B-ring of the cyathane core, an approach which leads back to allylic alcohol **17**. Similarly, Desmaële's route relies upon a Pd-catalyzed intramolecular Heck cyclization to construct the core B-ring, invoking diene **21** as a critical retrosynthetic precursor. Danishefsky's approach to the tricycle implements an Fe-catalyzed Nazarov cyclization for construction of the five-membered A-ring, allowing for strategic disconnection back to dienone **23**. Beyond the A- and B-rings, metal-catalyzed methodology has also been used to target the seven-membered C-ring of the cyathane core. For example, Snider's method toward these diterpene molecules employs an Al-catalyzed carbonyl-ene reaction for C-ring construction from bicyclic aldehyde **18**. The power of transition-metal catalysis has also enabled routes that employ a strategy of simultaneous multicyclic construction. For example, Phillips employs an efficient Ru-catalyzed ringopening ring-closing metathesis strategy to build both the A- and C-rings of the cyathane core in tandem, thus invoking bridged bicycle **22** as a retrosynthetic precursor. Wender's [5 + 2] cycloaddition approach toward the cyathane skeleton allows for cascade construction of both the B- and C-rings simultaneously, thus retrosynthetically disconnecting the core backward to ynone **20**. The strategy employed by Stoltz relies upon Ru-catalyzed C-ring construction, but more imperatively, implements Pd-catalyzed alkylation for quaternary stereocenter formation. This invokes bis-b-ketoester **19** as a critical retrosynthetic precursor.

Another unifying approach by which the cyathane tricycle has been targeted is that of an alkylation strategy, often specifically in the form of an aldol reaction. In order to construct the C-ring of the tricyclic diterpene core, Ward and his group employ an ozonolysis and aldol sequence. This ring-expanding strategy invokes tricycle **28** as an important synthetic precursor (Scheme 3a). Tori implements a similar technique to C-ring formation in his synthesis, in which bisaldehyde **26** is invoked via a disconnection of the cyathane core by means of an intramolecular aldol reaction. The cyathane C-ring has also been targeted via a unique [3 + 4] annulation reaction developed by Takeda, which employs **25** as a bicyclic precursor to the larger tricyclic skeleton. A convergent strategy targeting the B-ring of the cyathane core is carried out in Nakada's approach toward these natural products. By disconnecting the central ring of the tricycle, Nakada envisions tethered system **27** as the critical substrate for an intramolecular aldol reaction. Non-aldol alkylation procedures have also played a role in cyathane synthetic design. Reddy disconnects the smaller six-membered C-ring of cyanthiwigin AC via an enolate spiro-alkylation strategy to invoke bicycle **29** as a precursor (Scheme 3b). An anionic alkylation approach has also been employed with a focus upon construction of the five-membered cyathane A-ring. Piers' retrosynthetic analysis opens the A-ring to vinyl iodide **24**, a structure, which after lithium-halogen exchange and ketone trapping, is closed to give a tricyclic system.

4. Cyathane Core Syntheses

The section detailed below is intended as an update to the review published in 2000. As such, only cyathane core syntheses published after this date will be summarized here.

Several groups have devised efficient strategies toward the cyathane tricyclic core, with possible extensions toward multiple completed cyathane natural products. These efforts have focused upon preparation of the tricyclic framework found in the typical cyathane molecules.

4.1 Wender's Cyathane Core Synthesis

A general route toward the construction of the 5-6-7 tricyclic diterpene core was reported by the laboratories of Wender in 2001 by implementation of a [5 + 2] Rh-catalyzed cycloaddition reaction.¹³ Beginning with (–)-limonene (**30**), hydrogenation, oxidative olefin cleavage, and intramolecular aldol condensation afforded enal **31** (Scheme 4). After reduction and vinyl ether formation, a thermal Claisen rearrangement yielded aldehyde **32** as a 10:1 mixture of inseparable diastereomers. Cyclopentane **32** was thereafter advanced along four steps, including ozonolysis of the exocyclic methylene and addition of lithium cyclopropylacetylide, to furnish cyclopentanol **33** as a mixture of diastereomers. Stereoselective reduction of cyclopropyl alkyne **33** allowed access to diol **34**, which was oxidized, then exposed to 1-propynylmagnesium bromide to generate propargyl alcohol **35**. Attempts to execute a [5 + 2] cycloaddition reaction using alcohol **35** as the starting material were unfortunately unsuccessful, and yielded only a complicated mixture of products. For this purpose, enyne **35** was oxidized to conjugated ketone **20** before executing the [5 + 2] cycloaddition reaction.

Upon exposure of vinyl cyclopropane **20** to 5 mol% of [Rh(CO)₂Cl]₂, the desired cycloaddition reaction proceeded in high yield to provide tricycle **38** as a single diastereomer (Scheme 5).¹⁴ The critical [5 + 2] cycloaddition reaction initiates with complexation of the rhodium catalyst to both the alkyne moiety and the vinyl cyclopropane group. This is then followed by oxidative cyclometallation to form an intermediate metallocyclopentane (**36**), which in turn undergoes strain-driven cyclopropane ring-opening and ringexpansion to generate a transient metallocyclooctadiene species (**37**). Reductive elimination of rhodium from **37** thereafter yields tricyclic structure **38**, representing a completed cyathane core. The structure of tricyclic enone **38** was verified by single-crystal X-ray crystallography. Overall, this strategy allows access to a completed 5-6-7 tricyclic structure in 14 steps and 13% overall yield.

4.2 Desmaële's Cyathane Core Synthesis

A generalized synthetic route toward the tricyclic cyathane core structure was developed by Desmaële and coworkers in 2002, and an updated version of this strategy was later published in 2005.¹⁵ When addressing the challenges present in constructing the tricyclic cyathane framework, Desmaële states that establishing the anti relationship between the methyl groups of C(6) and C(9) represents the most significant obstacle. In order to solve this stereochemical issue, the strategy envisioned employs late-stage construction of the central B-ring of the tricycle via intramolecular Heck reaction between tethered A- and C-ring fragments.

The synthesis was initiated from known, enantioenriched keto-ester **39** (Scheme 6).¹⁶ Mukaiyama aldol reaction with acetaldehyde, followed by dehydration and isomerization, yielded enone **40**. After cuprate addition to and saponification of ester **40**, the intermediate keto-acid obtained was then subjected to the Kochi modification of the Hunsdiecker reaction to afford primary iodide **41**. Displacement of iodide **41** with the lithium enolate of methyl ester **42** then provided tethered intermediate **21**, which was subjected to a four step sequence to access the critical Heck cyclization precursor **43** (Scheme 7). Notably, prior attempts at an intramolecular Heck cyclization involving a sevenmembered ring dieneone proved quite

difficult to advance along this synthetic path, and for this reason Desmaële opted to employ a six-membered ring to serve as a surrogate for the cyathane C-ring.

Several attempts to cyclize precursor **43** via intramolecular Heck reaction under standard conditions were met with difficulty, with most attempts yielding either undesired acetate addition products or incorrect relative stereochemistry.^{15a} Eventually, further optimization of this reaction lead Desmaële to discover that exposure of triflate **43** to 20 mol% of Pd(OAc)₂ in the presence of PPh₃ and *n*-Bu₄NBr could execute the desired Heck reaction to yield tricycle **44** in 73% yield and a 19:1 diastereomeric ratio based on the newly formed stereocenter at C (6) (Scheme 8). This transformation constructed the central B-ring of the cyathane core while simultaneously establishing the necessary all-carbon quaternary stereocenter at C(6) via desymmetrization of the pendent C-ring precursor.^{15b,c}

Elaboration of tricyclic dienone **44** toward the cyathane core structure proceeded via hydrogenation of the disubstituted olefin using Wilkinson's catalyst to give enone **45**. This reduction was then followed by treatment of tricycle **45** with trimethyl aluminum and trimethylsilyl diazomethane to effect an organoaluminium-promoted ring expansion.¹⁷ This reaction afforded the completed 5-6-7 tricyclic framework as a 6:1 mixture of ketone (**46**) and enone (**47**) isomers. By obtaining these tricyclic structures, Desmaële accomplished the construction of the cyathane core tricycle over 15 steps and an overall combined yield of 1.4% for both isomers obtained.

4.3 Takeda's Cyathane Core Synthesis

A synthesis of the cyathane core leveraging a [4 + 3] annulation strategy was disclosed by Takeda and coworkers in 2000.¹⁸ Starting from known racemic enone **48**, addition of ethynyl Grignard was followed by a Rupe rearrangement to give extended conjugated enone system **25** (Scheme 9). Formation of the lithium enolate of **25** was followed by addition to acyl silane **49**. Initial nucleophilic addition of the enolate of **25** to **49** produces intermediate alkoxide **50**, which undergoes Brook rearrangement and subsequent nucleophilic addition to form cyclopropane **51**. Divinyl cyclopropane species **51** subsequently undergoes spontaneous [3,3] sigmatropic rearrangement, a mechanism which is further accelerated by the presence of an alkoxide. After rearrangement occurs, the completed cyathane core structure **52** is afforded as a single diastereomer.¹⁹

Continued functionalization of tricycle **52** was accomplished by diastereoselective DIBAL reduction, a process which provided the isomerically pure silyl enol ether **53** (Scheme 10). Notably, the stereochemistry imparted by this reduction afforded the alcohol epimer analogous to allocyathin B₂ at C(14).¹⁹ Takeda and coworkers thereafter concluded their efforts with oxidation of the enol silane present in **53** and subsequent cleavage of the C-bound trimethylsilyl group. This provided **54** as the final product of the synthetic sequence.

Beginning from known enone **48**, the des-methyl cyathane core was established in three steps and 19% yield, while the more elaborated cyathane analog **54** was produced in 11% yield over five steps.

5. Cyathane Total Syntheses

The following section is intended as a comprehensive review of all disclosed total syntheses of cyathane diterpenoid natural products published to date.

5.1 Snider's (±)-Allocyathin B₂ and (–)-Erinacine A

The first total synthesis of any cyathane diterpenoid natural product was accomplished by Snider and coworkers in 1996 with their preparation of allocyathin B₂.²⁰ Snider's synthetic

plan invoked the use of a carbonyl-ene reaction to target the cyathane core, and this strategy was later extended beyond allocyathin B₂ in order to achieve the synthesis of (+)-erinacine E via glycosylation.

Beginning with known racemic enone **48**,²¹ triflate formation, palladium-catalyzed carbonylation, and oxidation state manipulation allowed access to enal **55** (Scheme 11). Conjugate addition of a cuprate species generated from Grignard reagent **56** to the β -position of extended unsaturated system **55** provided aldehyde **57**, which was subsequently methylated at the α -position to afford bicycle **18**. At this point in the synthesis, Snider's route called for construction of the C-ring via intramolecular carbonyl-ene reaction of aldehyde **18**. In the event, treatment of **18** with Me₂AlCl initiated rearrangement to give a single isomer of alcohol **58** in excellent yield, thus completing the final ring of the tricyclic natural product.²²

The synthesis was completed over ten additional transformations, which involved protecting group manipulation, oxidation state modification, and palladiumcatalyzed carbonylation starting from tricycle **58** (Scheme 12). The completed natural product ($\pm$)-allocyathin B₂ (**4**) was thus attained from precursor **48** in 17 steps and 6.4% overall yield. Because allocyathin B₂ represents an aglycone substrate for the erinacine natural products, Snider and coworkers were well equipped at this point to address the total synthesis of the erinacine compounds. As such, glycosylation of allocyathin B₂ with 2,3,4-tri-*O*-acetyl- α -D-xylopyranosyl (**59**) bromide and successive global deprotection generated the natural product (–)-erinacine A (**5**) and isomeric structure **60** as a 1:1 mixture of diastereomers.

Erinacine A was prepared in 19 steps and 1.0% yield. The cyathane core framework was constructed in seven steps and 38% overall yield.

5.2 Tori's ($\pm$)-Allocyathin B₂

In a report published by Tori and coworkers in 1998, an intramolecular aldol strategy targeting the synthesis of the natural product allocyathin B₂ was described.²³

Starting from 3-methyl cyclohexenone (**61**), conjugate addition, ozonolysis, and oxidation yielded diketone **62** after five steps (Scheme 13). Intramolecular aldol condensation of cyclohexanone **62** afforded bicyclic enone **63**, a structure containing the completed five-membered A-ring of allocyathin B₂, including the requisite isopropyl group. Subsequent acylation of **63** with acid chloride **64** was followed by methylation and a highly optimized diastereoselective reduction employing Zn(BH₄)₂ to afford keto-alcohol **65**.²⁴ Additional transformations over seven steps provided access to allylic alcohol **66**, which was readily oxidized under Swern conditions to give bis-aldehyde **26**. Upon exposure of **26** to methanolic KOH, a final intramolecular aldol condensation completed the cyathane C-ring and produced allycyathin B₂ (**4**). Overall, both allocyathin B₂ and the cyathane core structure were synthesized in 19 steps and 0.5% yield, starting from **61**.

Subsequent to the preparation of allocyathin B₂, Tori also disclosed a different strategy toward the construction of the cyathane tricyclic core via ring-closing metathesis.²⁵ Modification of their route toward allocyathin B₂ allowed access to bicyclic intermediate **67** (Scheme 14). Upon treatment of this material with 20 mol% of Grubbs secondgeneration metathesis catalyst (**69**), completed cyathane tricycle **68** was obtained as the sole product of reaction.

5.3 Piers' ($\pm$)-Sarcodonin G

The first total synthesis of sarcodonin G was described by Piers in 2000.²⁶ Their approach to this cyathane diterpenoid targeted the tricyclic core with an alkylation and ring-expansion strategy, and employed late-stage installation of the peripheral functionality. Piers' synthesis began from known ketone **70** (introduced as a mixture of diastereomers),²⁷ which was subject

to hydrazone formation, epimerization at the ring fusion, and nucleophilic attack on alkyl iodide **71** to afford vinyl germane **72** (Scheme 15). Further transformation of germane **72** eventually produced the vinyl iodide species **24**.

Upon treatment of **24** with *n*-BuLi, lithium-halogen exchange and intramolecular trapping of the ketone moiety construct the five-membered A-ring of the natural product (Scheme 16). After deprotonation with KH and addition of Bu₃SnCH₂I, ether **73** was isolated as the major product. From this ether intermediate, a Still-Mitra [2,3]-sigmatropic rearrangement provides tricycle **74**, which contains the primary hydroxyl group at C(19) required for sarcodonin G.²⁸

Additional synthetic transformation of **74** over four steps yielded β -ketoester **75**, which bears an α -alkyl iodide group well suited for ring-expansion methodology developed by Hasegawa (Scheme 17).²⁹ Upon exposure to SmI₂ in THF, alkyl iodide **75** is converted in 71% yield to the one-carbon ring expanded product **76**. This process smoothly forms the seven-membered C-ring, and thus completes the tricyclic cyathane core. The total synthesis is thereafter concluded in six steps to yield sarcodonin G (**8**).

Overall the synthesis of sarcodonin G (**8**) was accomplished in a total of 21 steps and 4.0% yield. The cyathane tricyclic core was attained in 15 steps and 7.0% overall yield.

5.4 Ward's ($\pm$)-Allocyathin B₃

The total synthesis of allocyathin B₃ was achieved by Ward and coworkers in 2000 by leveraging an interesting cycloaddition strategy for rapid construction of the central B-ring.³⁰ The synthesis was initiated with a Diels–Alder cycloaddition between 2,5-dimethyl-*p*-benzoquinone (**78**) and 2,4-bis(trimethylsilyloxy)-1,3-pentadiene (**77**; Scheme 18). Subsequent [2 + 2] cycloaddition with allene and exposure to acidic conditions thereafter afforded a 4-6-6 tricyclic system as a 4:1 mixture of structural isomers (**28** to **79**), wherein each was produced as a single diastereomer.^{30a}

Though only isomer **28** was desired, the 4:1 mixture of **29** and **79** was epoxidized, reduced at the enone moiety, and then treated with benzenethiol at reflux under basic conditions to effect closure of the cyathane core A-ring. This furnished α -thiophenyl enone **80** in excellent yield, in a sequence that required only a single purification.^{30b}

Desulfurization, deoxygenation, protection, and epimerization over nine steps then allow access to 5-6-6 allylic benzoyl ester **81** (Scheme 19). The six-membered ring olefin of this intermediate provided the reactivity required to construct the seven-membered C-ring of the natural product. Ozonolysis of enone **81** in the presence of Sudan III indicator generated a sensitive keto-aldehyde intermediate which was subjected to aldol reaction, transacylation, and successive trapping via *O*-methylation to afford 5-6-7 tricycle **82**. This three step sequence completed the core framework and simultaneously established the trans-annular ketal bridge present in allocyathin B₃.

In order to install the necessary isopropyl side-chain appended to the A-ring of the natural product, intermediate **82** was then transformed to propargyl ether **83** (Scheme 20). Treatment of alkyl bromide **83** with AIBN in the presence of Ph₃SnH and subsequent hydrogenation produced the cyclized tetrahydrofuran **84**. After exposure to mild acid, tetracycle **84** opens to tricycle **85**, a structure which contains all of the carbon atoms required for completion of the A-ring. From this point, the total synthesis was completed in eleven steps to yield allocyathin B₃ (**3**).

Overall, Ward's strategy toward allocyathin B₃ comprises 34 steps and is concluded in 0.1% yield. The tricyclic cyathane core was attained in 18 steps and 11% overall yield.

5.5 Ward's (+)-Cyathin A₃

A variant of Ward's allocyathin A₃ strategy was later rendered enantioselective to achieve the total synthesis of the related diterpene natural product cyathin A.^{30,31}

The initial Diels–Alder reaction between 2,5-dimethyl-*p*-benzoquinone (**78**) and 2,4-bis(trimethylsilyloxy)-1,3-pentadiene (**77**) was made asymmetric by employing Mikami's titanium-based BINOL catalyst (Scheme 21).³³ After extensive optimization of this cycloaddition, Ward and coworkers found that addition of Mg powder and silica gel afforded cycloadduct **86** in 90% yield and 93% ee. With **86** in hand, [2 + 2] cycloaddition and acidic hydrolysis proceeded as in the case of Ward's allocyathin B₃ synthesis to furnish **28** and **79** as a 4:1 mixture of structural isomers.

From the isomeric mixture of **28** and **79**, eight additional steps lead to enone **87**, and thus allow Ward to follow his previously reported route (Scheme 22). Advancement of tricycle **87** along fifteen steps (including ozonolysis and aldol reaction for ring expansion) afforded the completed cyathane core in the form of acetal **88**. From this enone, oxidation, triflate formation, reductive deoxygenation, and hydrogenation produced tricycle **89** with the A-ring completed. With this material in hand, Ward was able to conclude the synthesis with four additional reactions, thus furnishing cyathin A₃ (**2**) in 34 steps and 1.0% overall yield. The cyathane core structure was attained in 17 steps and 15% overall yield.

5.6 Nakada's (+)-Allocyathin B₂

The first enantioselective total synthesis of allocyathin B₂ was described by the laboratories of Nakada in a report published in 2004.³³ In order to target this cyathane diterpenoid molecule, Nakada envisioned a convergent strategy starting with aldehyde **90** and alkyl iodide **91** (Scheme 23). Both fragment **90** and fragment **91** were prepared in enantioenriched form based on previous work published by Nakada and coworkers.^{34,35} After lithium/halogen exchange was performed on iodide **91**, the resulting alkyl lithium was added to a solution of aldehyde **90** to provide access to alcohol **92**. Deoxygenation, deprotection, and oxidation of this structure afforded diketone **93**.

The tethered diketone **93** was subsequently cyclized via intramolecular aldol reaction upon treatment with potassium *tert*-butoxide (Scheme 24). This nucleophilic attack served to form the C(5)–C(4) bond, thus constructing the central B-ring of the natural product. After dehydration, tricycle **94** was isolated as the major product of the reaction sequence. In seven additional steps ketone **94** was transformed into alkyl iodide **95**, an intermediate designed to undergo the samarium-mediated one-carbon ring expansion developed by Hasegawa and similar to that employed by Piers for the synthesis of Sarcodonin G (Scheme 17, Section 5.3).²⁹ When **95** was exposed to SmI₂ in the presence of HMPA, the expected migratory ring-expansion occurred in excellent yield to generate the completed cyathane core structure in the form of γ -keto ester **96**.

The route toward allocyathin B₂ was thereafter concluded with three additional oxidation state manipulations, finishing the synthesis in a total of 18 synthetic operations and 7.6% overall yield. The tricyclic cyathane core was attained in 15 steps and 13.4% overall yield. Additionally, this synthetic approach has been adapted toward a number of other cyathane natural products.³⁶

5.7 Nakada's (–)-Erinacine B

In 2007 Nakada and coworkers extended their synthetic route targeting allocyathin B₂ toward the preparation of erinacine B.³⁷ Following their previous synthetic efforts, it was possible to couple enantioenriched alkyl iodide **91** and aldehyde **90** to furnish, after elaboration, hydroxy

ketone **97** (Scheme 25).³⁵ After dehydration and deprotection to generate intermediate enone **98**, the envisioned synthetic route called for reduction of the α,β -unsaturation present across the central B-ring. Though numerous Birch reduction methods were attempted, it was discovered that such techniques provided mostly over-reduced diol products instead of the desired hydroxy ketone. Ultimately, this challenge was overcome via diastereoselective olefin reduction with SmI_2 in the presence of HMPA, which provided access to the desired tricycle **99**. This reduction was found to produce a single diastereomer of product, establishing the critical C(5) α -H stereochemistry required for (–)-erinacine B. From this point Nakada was able to once again draw from his synthesis of allocyathin B₂. Over eight steps, including SmI_2 -mediated ring-expansion, tricycle **100** was prepared.

Continued synthesis from ester **100** over nine additional steps allowed access to the protected allylic alcohol **101** (Scheme 26). From this point, reduction of the ketone moiety of **101** was required to establish the stereochemistry found in (–)-erinacine B. Unfortunately, all achiral reagents employed for the purpose of diastereoselective reduction afforded only the undesired epimer at C(14). Because of this difficulty, the ketone of **101** was reduced using the (*R*)-CBS catalyst, which set the desired relative stereochemistry with high selectivity to afford allylic alcohol **102**.

This material was thereafter glycosylated with a previously prepared xylose analog **103** to furnish glycone **104** (Scheme 27). After full deprotection of the carbohydrate ring, this material was treated with triethylamine and lithium bromide to effect an S_{N}' addition into the allylic alcohol. This provided (–)-erinacine B (**6**) as the sole product in enantioenriched form.

Overall, the synthesis was accomplished in 33 steps and 3.0% yield from **90** and **91**. The cyathane core structure was established in 18 steps and 14% overall yield.

5.8 Nakada's (–)-Erinacine E

One year subsequent to their report of erinacine B, Nakada and coworkers disclosed a beautiful modification of their synthetic strategy to target (–)-erinacine E, arguably the most complex cyathane diterpenoid molecule isolated to date.³⁸ Starting from agylcone molecule **102**, glycosylation with thioglycoside **105** and deprotection provided access to intermediate **106** (Scheme 28). After additional protecting group manipulation, Swern oxidation generated ketone **107**, an intermediate which spontaneously undergoes conjugate addition-elimination to form pentacyclic structure **108**.

Treatment of enal **108** with DBU in benzene at room temperature initiated an intramolecular aldol reaction between the enolate generated at C(4') and the aldehyde present at C(15) (Scheme 29). This aldol reaction is followed by spontaneous benzoate ester migration, thus producing protected erinacine E analog **109** as the final product of the sequence. Further deprotection, oxidation, and diastereoselective carbonyl reduction thereafter complete the total synthesis of (–)-erinacine E (**7**).

Overall, erinacine E was prepared in 39 steps from iodide **91** and aldehyde **90**, in a total yield of 0.9%.

5.9 Trost's (+)-Allocyathin B₂

Trost and coworkers have recently reported a unique Rucatalyzed cycloisomerization strategy for the total synthesis of the cyathane diterpenoid molecule allocyathin B₂.³⁹ The synthesis was initiated with a Pd-catalyzed asymmetric allylic alkylation. Racemic ketone **110** was alkylated in the presence of $[(\eta^3\text{-C}_3\text{H}_7)\text{PdCl}]_2$ and chiral ligand (*S,S*)-**112** to afford enantioenriched α -quaternary cyclopentanone **111** (Scheme 30).⁴⁰ The high-yielding allylation establishes an all-carbon quaternary stereocenter in 95% ee. This transformation not

only set the configuration required at C(9) of the natural product, but also provided the stereochemical basis upon which all subsequent diastereoselective transformations were leveraged.

Further elaboration of the allyl side-chain of **111**, as well as manipulation of the ketone moiety, eventually produced propargyl ester **17** (Scheme 31). With this material, Trost and coworkers planned to seal the central B-ring of allocyathin B₂ via an intramolecular ruthenium-mediated cycloisomerization reaction. In the event, treatment of allylic alcohol **17** with CpRu(CH₃CN)₃PF₆ initiated cyclization of the conjugated alkyne onto the trisubstituted olefin with concomitant oxidation of the primary alcohol.⁴¹

Notably, the cycloisomerization reaction proceeds with high diastereoselectivity. Both products obtained from this transformation possessed the desired anti relationship between the two all-carbon quaternary stereocenters, and no products bearing a syn arrangement were observed. Trost hypothesizes that initial ruthenium complexation to both the alkyne and alkene moieties of **17** can occur to from either a syn-coplanar orientation (**115**) or an orthogonal orientation (**117**, Scheme 32). Because the orbital overlap of the syncoplanar arrangement is likely more conducive to cycloisomerization, ruthenacycle formation from intermediate **115** to give **116** is expected to be much faster than the alternative formation of **118** from **117**. After b-hydride elimination occurs from ruthenacycle **116**, reductive elimination forms **113** and **114** as the major products of reaction. Because only **113** was desired, ultimately it was determined that increasing the size of the alkynyl ester to a bulky *tert*-butyl provided preferential formation of the Z olefin isomer in a 6.7:1 ratio with isomer **114**.^{39b}

From **113**, a hydroxylative Knoevenagel reaction was performed to access lactone **119** as a single diastereomer (Scheme 33).⁴² The final stages of the synthesis involved hydrogenation and nitrile/lactone reduction, followed by an intramolecular aldol reaction to yield (+)-allocyathin B₂ (**4**). Following this sequence, the natural product allocyathin B₂ (and the cyathane core) was synthesized in sixteen steps and 5.2% overall yield.

5.10 Danishefsky's (-)-Scabronine G

In 2005, Danishefsky and coworkers reported the first total synthesis of the cyathane diterpenoid scabronine G (**9**).⁴³ In order to target this bioactive compound, they approached the synthesis with “the pleasingly simple idea that scabronine G can be viewed as an annulated, one-carbon ring-expanded version of the (-)-Wieland–Miescher ketone”. The initial transformations of the synthesis commenced from the protected ketone **120** (Scheme 34). Over five steps, including kinetic enolate trapping and palladium-catalyzed carbonylation, acetal **120** was converted to dieneone **23**. Construction of the five-membered A-ring of scabronine G was then accomplished via Lewis acid-promoted Nazarov reaction. Upon treatment of dienone **23** with FeCl₃, cyclization proceeded smoothly to afford enone **121** in good yield. Notably, this tetracyclic product was obtained as a single tetrasubstituted olefin isomer.

The enone of tricycle **121** was then subsequently leveraged in order to install the critical all-carbon quaternary stereocenter present at C(9) of scabronine G. Conjugate addition of Nagata's reagent was observed to occur with high axial diastereoselectivity, and the resulting enolate was trapped with TMSCl (Scheme 35).⁴⁴ The intermediate silyl enol ether obtained from this sequence was then converted to a vinyl triflate species to provide nitrile **122**. Further synthetic elaboration over seven steps allowed for the conversion of vinyl triflate **122** to thiopropylmethyldene **123**.

In order to access the 5-6-7 tricyclic cyathane core, vinylogous thioester **123** was first treated with the lithium anion of methoxymethyl phenyl sulfide to produce intermediate tertiary alcohol **124** as a combination of diastereomers (Scheme 36). Subsequent exposure of this

mixture to HgCl_2 effected a one-carbon ring expansion reaction to furnish tricycle **125** and the seven-membered C-ring, thus completing the cyathane framework.⁴⁵ With ringexpanded aldehyde **125** in hand Danishefsky and coworkers were able to complete (–)-scabronine G (**9**) in three additional transformations, accomplishing the total synthesis in 20 steps and 8.2% overall yield. The cyathane core framework was constructed in 17 steps and 11% yield.

5.11 Phillip's (+)-Cyanthiwigin U, (+)-Cyanthiwigin W, and (–)-Cyanthiwigin Z

The first report of the total synthesis of any member of the cyanthiwigin sub-class of natural products was published by Phillips and coworkers in 2005.⁴⁶ Phillips' strategy for targeting cyanthiwigin U focused upon construction of the tricyclic cyathane core via simultaneous construction of both the A- and C-rings, and employed late-stage installation of the peripheral functionality.

The synthesis began with an asymmetric Diels–Alder reaction between 1,4-dimethyl cyclohexadiene (**127**) and (–)-borneol-appended enone **126** (Scheme 37).⁴⁷ This cycloaddition reaction proceeded smoothly to give a single diastereomer of **128**, thus establishing both quaternary stereocenters of the natural product in a single synthetic operation. After cleavage of the chiral controller, deprotection, and further functional group manipulation, this sequence provided access to bridged bicyclic bis-aldehyde **22** (Scheme 38).

Addition of vinyl Grignard to bis-aldehyde **22** was followed oxidation to bis-enone **129**. The bicyclo[2.2.2]octene system of **129** was designed to be well suited to the planned synthetic strategy of “two-directional” tandem ring-opening, ring-closing metathesis developed by Phillips in prior reports.⁴⁸ By treating this bis-enone with 20 mol% of Grubbs' second-generation metathesis catalyst (**69**) under an atmosphere of ethylene, ring-opening metathesis of the strained bridging olefin was rapidly followed by two sequential ring-closing events. This reaction established both the five- and seven-membered rings of the cyathane core, providing tricycle **130** as the ultimate product of the cascade. From this point, the synthesis of (+)-cyanthiwigin U (**11**) was completed in four steps involving oxidation state manipulation, as well as addition of the isopropyl and methyl groups. Phillip's route produces the cyathane core in eight synthetic operations and 19% yield, and in total, the synthesis comprises only 12 steps and is accomplished in 17% overall yield.

Very recently, the laboratories of Phillips have reported the synthesis of additional cyanthiwigin natural products based on an extension of their route toward cyanthiwigin U.⁴⁹ Treatment of cyanthiwigin U (**11**) under Luche reduction conditions afforded the natural product cyanthiwigin W in 9:1 dr (**131**, Scheme 39). Protection of the resulting secondary hydroxyl group was then followed by allylic transposition and alcohol oxidation via PCC. After deprotection, this sequence furnished cyanthiwigin Z (**132**).

5.12 Reddy's (+)-Cyanthiwigin AC

The structure of cyathane diterpenoid cyanthiwigin AC is unique in comparison to other cyathane molecules in that it does not possess a 5-6-7 fused tricyclic core. Instead, the natural product contains both a 5-6 fused bicycle and a 6-6 spirocyclic junction. The first total synthesis of this natural product was reported in 2006 by Reddy and coworkers.⁵⁰

Starting from the (+)-Hajos–Parrish ketone derivative **29**, double-alkylation with bis-mesylate **133** was effective to install the spiro-annulation of the natural product (Scheme 40). After methylenation of the resulting compound, spirocyclic intermediate **134** was isolated as the major product of this two-step sequence. Deprotection, oxidation, and isomerization then furnished enone **135** in preparation for installation of the isopropyl sidechain to the five-membered A-ring.

Conjugate addition of isopropenyl cyanocuprate to tetracycle **135** was followed by diastereoselective reduction of the exocyclic methylene to set the tertiary methyl stereocenter of the natural product (Scheme 41). This yielded ketone **136** as a 2:1 mixture of epimers at C (6). Further elaboration of this material provided access to thioether **137**, which upon exposure to IBX in toluene underwent oxidation to generate a mixture of six-membered ring enones.⁵¹ After sulfoxide formation and dehydrosulfonylation of this intermediate mixture, enones **138** and **139** were obtained in a 1:1.2 ratio of isomers.

From this point, the addition of methyl lithium to dienone **139** produced the natural product as a 1:2 mixture of epimers at C(12), favoring the generation of (+)-cyanthiwigin AC (**12**, Scheme 42) over isomer **140**. The total synthesis of the natural product was concluded in 13 steps and 2.0% overall yield.

5.13 Stoltz's (–)-Cyanthiwigin F

The total synthesis of cyanthiwigin F via a double catalytic enantioselective alkylation was published by Stoltz in 2008.⁵² Tandem Claisen condensation and Dieckmann cyclization of diallyl succinate (**141**) were followed by methylation of both α positions to yield bis- β -ketoester **19** (Scheme 43). Treatment of this material with a palladium catalyst precursor and (*S*)-*t*-BuPHOX (**142**) as a ligand initiates a double catalytic enantioselective alkylation reaction to generate both quaternary stereocenters present in the natural product.⁵³ This reaction affords enantioenriched diketone **143** in 99% ee, as well as the corresponding meso diastereomer **144**, in a 4.4:1 diastereomeric ratio.

Diketone **143** was subsequently desymmetrized via triflate formation and Negishi cross-coupling with alkyl iodide **144** to furnish tetraene **145** (Scheme 44). This material was subjected to ring-closing metathesis by the action of modified Grubbs' catalyst **146** accompanied by cross-metathesis with vinyl boronate **147**. This transformation served to close the seven-membered C-ring of the core scaffold while subsequently functionalizing the remaining allyl side-chain of **145** toward further elaboration. After mild oxidative work-up with aqueous sodium perborate, bicyclic aldehyde **148** was obtained in a single synthetic operation from **145**. Closure of the final ring of the cyathane framework was then accomplished via an acyl radical cyclization in a reaction which yielded a single diastereomer of tricyclic diketone **151**.⁵⁴ A potential explanation for the diastereoselectivity of this reaction involves rapid formation of the five-membered A-ring through intermediate acyl radical **149**, followed by external hydrogen atom abstraction by tertiary radical **150** to give diketone **151**. After vinyl triflate formation from this diketone, palladium-catalyzed cross-coupling with a pregenerated isopropyl cyanocuprate species produced both (–)-cyanthiwigin F (**10**) and reduced product **152** in a 1.8:1 ratio.

The total synthesis of cyanthiwigin F was achieved in nine steps and 1.9% overall yield. The tricyclic cyathane core was accessed in seven steps and 6.2% overall yield.

6. Conclusions

The cyathane diterpenoid natural products have been the focus of numerous total synthetic efforts. Sixteen completed total syntheses of these compounds have been reported, thirteen of which have emerged in the last eight years alone. Because the cyathane natural products have been implicated as important biologically active molecules, particularly in regard to the stimulation of NGF synthesis, continued investigation into and refinement of their laboratory preparation is undoubtedly forthcoming.

7. Acknowledgements

The authors wish to thank NIH-NIGMS (R01GM080269-01), Amgen, Abbott, Boehringer Ingelheim, Merck, and Bristol-Myers Squibb for financial support. JAE extends his thanks to Roche for support in the form of the excellence in chemistry award.

Biographies

John Enquist was born in Santa Rosa, CA, USA in 1982. He received his BA in chemistry from the University of California, San Diego in 2004, where he worked in the laboratories of Prof. Yitzhak Tor. He has been a graduate student in the laboratories of Prof. Brian Stoltz since 2004, where his work has focused on the total synthesis of natural product molecules. His research interests include cascade reactions, synthesis of biologically active natural products, transition metal-catalyzed transformations, and asymmetric methodology with a focus on the formation of multiple stereocenters.

Brian M. Stoltz was born in Philadelphia, PA, USA in 1970. After spending a year at the Ludwig Maximilians Universität in München, Germany, he obtained his BS in Chemistry and BA in German from Indiana University of Pennsylvania in 1993. He then earned his Ph.D. in 1997 under the direction of Professor John L. Wood at Yale University specializing in synthetic organic chemistry. Following an NIH postdoctoral fellowship in the laboratories of Professor E. J. Corey at Harvard University (1998-2000), he joined the faculty at Caltech in 2000 where he is currently the Ethel Wilson Bowles and Robert Bowles Professor of Chemistry. His research focuses on the design and implementation of new synthetic strategies for the synthesis of complex molecules possessing important biological properties, in addition to the development of new synthetic methods including asymmetric catalysis and cascade processes. In addition to awards from a number of pharmaceutical companies (i.e., Abbott, Amgen, AstraZeneca, Boehringer Ingelheim, Bristol-Myers Squibb, Eli Lilly, GlaxoSmithKline, Johnson and Johnson, Merck, Novartis, Pfizer, Roche), Professor Stoltz is the recipient of the Camille and Henry Dreyfus New Faculty and Teacher-Scholar Awards, a National Science Foundation CAREER Award, the Research Corporation Research Innovation and Cottrell Scholars Awards, an A. P. Sloan Research Fellowship, an Arthur C. Cope Scholar Award from the American Chemical Society, and has received the Presidential Early Career Award in Science and Engineering (PECASE) from the White House. In 2006, he was elected a fellow of the American Association for the Advancement of Science (AAAS) and in 2008 was named a KAUST GRP Investigator. Additionally, Professor Stoltz was recognized by the Caltech Graduate Student Council with both a Classroom Teaching Award and a Mentoring Award in 2001 and by the Associated Students of the California Institute of Technology for their 30th Annual Award for Excellence in Teaching in 2006.

References

1. Wright DL, Whitehead CR. *Org. Prep. Proced. Int* 2000;32:309–330.
2. Allbutt AD, Ayer WA, Brodie HJ, Johri BN, Taube H. *Can. J. Microbiol* 1971;17:1401–1407. [PubMed: 5156938]
3. (a) Ayer WA, Taube H. *Tetrahedron Lett* 1972:1917–1920. (b) Ayer WA, Taube H. *Can. J. Chem* 1973;51:3842–3854.
4. Ayer WA, Ling SP. *Can. J. Chem* 1979;57:3332–3337.
5. (a) Kawagishi H, Shimada A, Shirai R, Okamoto K, Ojima F, Sakamoto H, Ishiguro Y, Furukawa S. *Tetrahedron Lett* 1994;35:1569–1572. (b) Kawagishi H, Shimada A, Hosokawa S, Mori H, Sakamoto H, Ishiguro Y, Sakemi S, Bordner J, Kojima N, Furukawa S. *Tetrahedron Lett* 1996;37:7399–7402.
6. Shibata H, Tokunaga T, Karasawa D, Hirota A, Nakayama M, Nozaki H, Taka T. *Agric. Biol. Chem* 1989;53:3373–3375.

7. (a) Ohta T, Kita T, Kobayashi N, Obara Y, Nakahata N, Ohizumi Y, Takaya Y, Oshima Y. *Tetrahedron Lett* 1998;39:6229–6232. (b) Kita T, Takaya Y, Oshima Y, Ohta T. *Tetrahedron* 1998;54:11877–11886. (c) Ma B-J, Zhu H-J, Liu J-K. *Helv. Chim. Acta* 2004;87:2877–2881.
8. Green D, Goldberg I, Stein Z, Ilan M, Kashman Y. *Nat. Prod. Lett* 1992;1:193–199.
9. (a) Peng J, Walsh K, Weedman V, Bergthold JD, Lynch J, Lieu KL, Braude IA, Kelly M, Hamann MT. *Tetrahedron* 2002;58:7809–7819. (b) Peng J, Avery MA, Hamann MT. *Org. Lett* 2003;5:4575–4578. [PubMed: 14627387] (c) Peng J, Kasanah N, Stanley CE, Chadwick J, Fronczek FR, Hamann MT. *J. Nat. Prod* 2006;69:727–730. [PubMed: 16724830]
10. Sennett SH, Pomponi SA, Wright AE. *J. Nat. Prod* 1992;55:1421–1429. [PubMed: 1453179]
11. Conner B, Dragunow M. *Brian Res. Rev* 1998;27:1–39.
12. Ayer WA, Ling SP, Nakashima TT. *Can. J. Chem* 1979;57:3338–3343.
13. Wender PA, Bi FC, Brodney MA, Gosseln F. *Org. Lett* 2001;3:2105–2108. [PubMed: 11418060]
14. For previous examples of [5 + 2] cycloadditions, see: (a) Wender PA, Takahashi H, Witulski B. *J. Am. Chem. Soc* 1995;117:4720–4721.4721; (b) Wender PA, Husfeld CO, Langkopf E, Love JA. *J. Am. Chem. Soc* 1998;120:1940–1941.1941; (c) Wender PA, Dyckman AJ, Husfeld CO, Kadereit D, Love JA, Rieck H. *J. Am. Chem. Soc* 1999;121:10442–10443.10443
15. (a) Thominiaux C, Chiarnoi A, Desmaële D. *Tetrahedron Lett* 2002;43:4107–4110. (b) Drège E, Morgant G, Desmaële D. *Tetrahedron Lett* 2005;46:7263–7266. (c) Drège E, Tominaux C, Morgant G, Desmaële D. *Eur. J. Org. Chem* 2006:4825–4840.
16. The ketoester **39** can be prepared via Michael addition to the imine obtained after treatment of 1-methylcyclopentanone with (*S*)-1-phenylethylamine. See: (a) Pfau M, Revial G, d'Angelo J, Guingant A. *J. Am. Chem. Soc* 1985;107:273–274.274; (b) d'Angelo J, Desmaële D, Dumas F, Guingant A. *Tetrahedron: Asymmetry* 1992;3:459–505.505
17. (a) Maruoka K, Concepcion AB, Yamamoto H. *J. Org. Chem* 1994;59:4725–4726. (b) Maruoka K, Concepcion AB, Yamamoto H. *Synthesis* 1994:1283–1290. (c) Yang S, Hungerhoff B, Metz P. *Tetrahedron Lett* 1998;39:2097–2098.
18. Takeda K, Nakane D, Takeda M. *Org. Lett* 2000;2:1903–1905. [PubMed: 10891187]
19. Takeda K, Nakajima A, Takeda M, Okamoto Y, Sato T, Yoshii E, Koizumi T, Shiro M. *J. Am. Chem. Soc* 1998;120:4947–4959.
20. (a) Snider BB, Vo NH, O'Neil SV, Foxman BM. *J. Am. Chem. Soc* 1996;118:7644–7645. (b) Snider BB, Vo NH, O'Neil SV. *J. Org. Chem* 1998;63:4732–4740.
21. Snider BB, Rodini DJ, van Straten J. *J. Am. Chem. Soc* 1980;102:5872–5880.
22. Snider, BB. *Comprehensive Organic Synthesis*. Vol. 2. Oxford; Pergamon: 1991. p. 527 (b) Snider BB, Yang KJ. *J. Org. Chem* 1990;55:4392–4399.
23. Tori M, Toyoda N, Sono M. *J. Org. Chem* 1998;63:306–313.
24. (a) Gensler WJ, Johnson F, Sloan ADB. *J. Am. Chem. Soc* 1960;82:6074–6081. (b) Crabbé P, Garcia GA, Rius C. *J. Chem. Soc., Perkin Trans* 1973;1:810–816.
25. Nakashima K, Fujisaki N, Inoue K, Minami A, Nagaya C, Sono M, Tori M. *Bull. Chem. Soc. Jpn* 2006;79:1955–1962.
26. Piers E, Gilbert M, Cook KL. *Org. Lett* 2000;2:1407–1410. [PubMed: 10814459]
27. Piers E, Yeung BKA, Fleming FF. *Can. J. Chem* 1993;71:280–286.
28. Still WC, Mitra A. *J. Am. Chem. Soc* 1978;100:1927–1928.
29. Hasegawa E, Kitazume T, Suzuki K, Tosaka E. *Tetrahedron Lett* 1998;39:4059–4062.
30. (a) Ayer WA, Ward DE, Browne LM, Delbaere LTJ, Hoyano Y. *Can. J. Chem* 1981;59:2665–2672. (b) Ward DE. *Can. J. Chem* 1987;65:2380–2384. (c) Ward DE, Gai Y, Qiao Q. *Org. Lett* 2000;2:2125–2127. [PubMed: 10891246]
31. Ward DE, Shen J. *Org. Lett* 2007;9:2843–2846. [PubMed: 17580883]
32. Mikami K, Motoyama Y, Terada M. *J. Am. Chem. Soc* 1994;116:2812–2820.
33. Takano M, Umino A, Nakada M. *Org. Lett* 2004;6:4897–4900. [PubMed: 15606094]
34. Honma M, Sawada T, Fujisawa Y, Utsugi M, Watanabe H, Umino A, Matsumura T, Hagihara T, Takano M, Nakada M. *J. Am. Chem. Soc* 2003;125:2860–2861. [PubMed: 12617637]
35. Iwamoto M, Kawada H, Tanaka T, Nakada M. *Tetrahedron Lett* 2003;44:7239–7243.

36. An analogous synthetic approach was reported by Nakada for progress toward the synthesis of (-)-scabronine A. See: Watanabe H, Nakada M. *Tetrahedron Lett* 2008;49:1518–1522.1522
37. Watanabe H, Takano M, Umino A, Ito T, Ishikawa H, Nakada M. *Org. Lett* 2007;9:359–362. [PubMed: 17217304]
38. Watanabe H, Nakada M. *J. Am. Chem. Soc* 2008;130:1150–1151. [PubMed: 18181632]
39. (a) Trost BM, Dong L, Schroeder GM. *J. Am. Chem. Soc* 2005;127:2844–2845. [PubMed: 15740107]
(b) Trost BM, Dong L, Schroeder GM. *J. Am. Chem. Soc* 2005;127:10259–10268. [PubMed: 16028937]
40. (a) Trost BM, Schroeder GM. *J. Am. Chem. Soc* 1999;121:6759–6760. (b) Trost BM, Schroeder GM, Kristensen J. *Angew. Chem., Int. Ed* 2002;41:3492–3495.
41. Trost BM, Toste FD, Pinkerton AB. *Chem. Rev* 2001;101:2067–2096. [PubMed: 11710241]
42. Trost BM, Mallert S. *Tetrahedron Lett* 1993;34:8025–8028.
43. Waters SP, Tian Y, Li Y-M, Danishefsky SJ. *J. Am. Chem. Soc* 2005;127:13514–13515. [PubMed: 16190712]
44. Nagata W, Mitsuru Y, Masayuk M. *J. Am. Chem. Soc* 1972;94:4644–4653.
45. Guerrero A, Parrilla A, Camps F. *Tetrahedron Lett* 1990;31:1873–1876.
46. Pfeiffer MWB, Phillips AJ. *J. Am. Chem. Soc* 2005;127:5334–5335. [PubMed: 15826167]
47. Palomo C, Oiarbide M, García JM, González A, Lecumberri A, Linden A. *J. Am. Chem. Soc* 2002;124:10288–10289. [PubMed: 12197725]
48. Minger TL, Phillips AJ. *Tetrahedron Lett* 2002;43:5357–5359.
49. Pfeiffer MWB, Phillips AJ. *Tetrahedron Lett* 2008;49:6860–6861.
50. Reddy TJ, Bordeau G, Trimble L. *Org. Lett* 2006;8:5585–5588. [PubMed: 17107078]
51. Nicolaou KC, Zhong Y-L, Baran PS. *J. Am. Chem. Soc* 2000;122:7596–7597.
52. Enquist JA Jr, Stoltz BM. *Nature* 2008;453:1228–1231. [PubMed: 18580947]
53. (a) Behenna DC, Stoltz BM. *J. Am. Chem. Soc* 2004;126:15044–15045. [PubMed: 15547998] (b) Mohr JT, Behenna DC, Harned AM, Stoltz BM. *Angew. Chem., Int. Ed* 2005;44:6924–6927.
54. Yoshikai K, Hayama T, Nishimura K, Yamada K, Tomioka K. *J. Org. Chem* 2005;70:681–683. [PubMed: 15651818]

Figure 1.
Representative cyathane diterpenoid natural products.

Scheme 1.

Biosynthesis of the cyantane core tricycle from geranylgeranyl phosphate.

Scheme 2.
Transition metal and Lewis acid-catalyzed retrosynthetic disconnections toward the cyathane diterpenoid tricyclic core.

Scheme 3.

(a) Alkylation and aldol retrosynthetic disconnections of the cyathane diterpenoid tricyclic core. (b) Reddy's retrosynthetic disconnection of the cyanthiwigin AC core.

Scheme 4.

Preparation of the critical [5 + 2] cycloaddition precursor.

Scheme 5.

Wender's [5 + 2] cycloaddition reaction to construct the cyathane core tricycle.

Scheme 6.

Synthesis of Desmaële's alkyl iodide coupling partner.

Scheme 7.
Preparation of the crucial Heck cyclization precursor.

Scheme 8.

Heck cyclization and aluminum-promoted ring expansion reactions to target the cyathane tricycle.

Scheme 9.

Takeda's key [4 + 3] annulation to target the C-ring of the cyathane core

Scheme 10.
Advancement of Takeda's cyathane core structure.

Scheme 11.
Snider's carbonyl-ene strategy toward the cyathane tricycle.

Scheme 12.

Completion of allocyathin B₂ and glycosylation to achieve erinacine A.

Scheme 13.
Tori's strategy toward allocyathin B₂.

Scheme 14.

Tori's ring-closing strategy toward construction of the seven-membered C-ring and completion of the cyathane core.

Scheme 15.

Vinyl iodide construction in Pier's sarcodonin synthesis.

Scheme 16.

Still-Mitra [2,3] sigmatropic rearrangement to construct the A-ring of sarcodonin G.

Scheme 17.
Ring expansion and endgame for sarcodonin G.

Scheme 18.

Ward's cycloaddition strategy toward tricyclic formation.

Scheme 19.

Ring expansion via ozonolysis and aldol reaction to target the cyathane core.

Scheme 20.
Ward's allocyathin endgame strategy.

Scheme 21. Catalytic enantioselective Diels-Alder approach to intermediates **86** and **29**.

Scheme 22.

Endgame synthesis for cyathin A₃. (TTBP = 2,6-di(*tert*-butyl)-4-methylpyridine.)

Scheme 23.
Fragment coupling to prepare the tethered aldol precursor.

Scheme 24.

Intramolecular aldol reaction and ring expansion to complete the total synthesis of allocyathin B₂.

Scheme 25.
Adaptation of Nakada's allocyathin B₂ route to erinacine B.

Scheme 26.
Establishing the desired stereochemistry at C(14).

Scheme 27.
Completion of (-)-erinacine B.

Scheme 28.

Synthesis of **108** from allocyathin precursor **102** via conjugate addition-elimination.

Scheme 29.

Intramolecular aldol reaction for Nakada's endgame of erinacine E.

Scheme 30.

Asymmetric allylic alkylation to establish stereochemistry at C(9).

Scheme 31.

Ru-catalyzed cycloisomerization to establish the central Bring of allocyathin B₂.

Scheme 32.

Diastereoselectivity considerations in the ruthenium catalyzed cycloisomerization.

Scheme 33.

Completion of allocyathin B₂ via hydroxylative Knoevenagel and intramolecular aldol reactions.

Scheme 34.

Danishefsky's Nazarov strategy for cyathane A-ring construction.

Scheme 35.

Advancement toward Danishefsky's ring-expansion precursor.

Scheme 36.

Ring expansion and endgame to complete (-)-scabronine G.

Scheme 37.

Auxiliary-mediated Diels–Alder reaction to establish the quaternary stereocenters of the natural product.

Scheme 38.
Completion of cyanthiwigin U via ring-opening ring-closing metathesis.

Scheme 39.

Preparation of cyanthiwigin W and cyanthiwigin Z from cyanthiwigin U.

Scheme 40.
Spiro-annulation of Hajos-Parrish ketone derivative **29**.

Scheme 41.

Further oxidation and transformation toward cyanthiwigin AC.

Scheme 42.
Endgame of cyanthiwigin AC.

Scheme 43.
Catalytic enantioselective double-alkylation for cyanthiwigin F.

Scheme 44.

Completion of the total synthesis of cyanthiwigin F.