Published in final edited form as:

Mov Disord. 2008 December 15; 23(16): 2423–2424. doi:10.1002/mds.22011.

Unawareness of Head Tremor in Essential Tremor: A Study of Three Samples of Essential Tremor Patients

Elan D. Louis, MD, MSc^{1,2,3,4}, Kathryn M. Pellegrino, BA¹, and Eileen Rios, BS¹

¹ *GH Sergievsky Center, College of Physicians and Surgeons, Columbia University, New York, NY, USA*

² *Department of Neurology, College of Physicians and Surgeons, Columbia University, New York, NY, USA*

³ *Taub Institute for Research on Alzheimer's Disease and the Aging Brain, College of Physicians and Surgeons, Columbia University, New York, NY, USA*

⁴ *Department of Epidemiology, Mailman School of Public Health, Columbia University, New York, NY, USA*

Keywords

essential tremor; head tremor; awareness

Patients with Huntington disease may be unaware of their chorea and patients with Parkinson disease often do not recognize or endorse their medication-induced dyskinesias.^{1, 2} Head tremor occurs in essential tremor (ET),^{3, 4} yet patients in our experience are often unaware of it. While this phenomenon is anecdotally noted, it has not been formally documented or studied systematically.

To broadly sample ET in different settings, we selected cases from three settings: a tertiary-referral center (largest sample),⁵ a population-based study in Manhattan,⁶ and a brain repository.⁷ As expected, cases differed in several respects (Table). Using the same clinical questionnaire, each case was asked whether he/she sometimes has a head tremor. A 20 minute videotaped tremor examination was performed, which included assessments of arm and head tremors (including sitting facing the camera, sustained phonation, reading aloud, finger-nose-finger maneuver, drinking, using a spoon, stand facing camera, and walking). The videotape was reviewed (E.D.L.); arm tremors were rated using a 0–3 scale.⁸ Head tremor was rated as absent (0), mild or equivocal (1), intermittent yet clearly present (2), moderate (3), or severe (4).

In the tertiary-referral center, 119/320 (37.2%) ET cases had head tremor on examination (≥ 1); 46/119 (38.7%) did not report having head tremor. Most false negatives asked the interviewer, “do you see a head tremor now?” When told “yes”, they uniformly commented that they were unaware of this (e.g., “I don’t notice it”). By contrast to head tremor, of 298 ET cases with dominant hand tremor while writing on examination (rating ≥ 1), only 60 (20.1%) did not report having hand tremor while writing (OR 0.40, 95% CI 0.25 – 0.64; i.e., ET cases were 2.5 times less likely to report head tremor than handwriting tremor). Patients may under-report tremor because of embarrassment; however, when we restricted analyses to 129 ET cases who reported that they were *not* embarrassed by tremor, 21/43 (48.8%) with head tremor did not report their head tremor. When we restricted analyses to 78 ET cases who had moderate or severe head tremor on examination, approximately one-quarter (18/78 [23.1%]) did not

report head tremor (as compared to only 12/150 [8.0%] cases with moderate to severe handwriting tremor who did not report their handwriting tremor, OR 3.45, 95% CI 1.56 – 7.61). Based on these findings, we consider that failure to report head tremor was not merely due to having mild head tremor.

In the population-based sample, 13/106 (12.3%) cases had head tremor on examination; 7/13 (53.8%) did not report having head tremor. In the brain repository, 92/170 (54.1%) cases had head tremor on examination, but 30/92 (32.6%) did not report having head tremor.

In each setting, one-third to one-half of ET cases did not report the presence of head tremor. These same cases were two to three times more likely to report their hand tremor than their head tremor. One possibility is ET cases failed to report head tremor because they were poor historians or were embarrassed. However, they reported hand tremor with reasonably high validity and, among those who were unembarrassed by their tremor, nearly one-half did not report their head tremor. A second consideration is cognitive impairment.¹ This is unlikely; in our largest sample (tertiary-referral clinic), cases with cognitive impairment (Telephone Interview for Cognitive Status⁹ <31) were excluded. Third, even when head tremor impedes a skilled task (e.g., while shaving), patients are often unsure whether this is due to shaky hands or a shaky head. A final possibility is that cases were actually unaware of their head tremor. Indeed, when their tremor was pointed out to them, false negatives uniformly stated that they were unaware of it. A lack of internal feedback about a movement may lessen self-awareness of that movement.^{1,2} Whether, from a proprioceptive vantage point, patients have a subjective experience of head tremor, is not clear. For example, with some types of oscillatory cranial movements (e.g., patients with congenital nystagmus, who rarely experience oscillopsia¹⁰) perceptual stability (i.e., lack of awareness of nystagmus) may be achieved through a reduced sensitivity to the motion or the use of other signals to cancel the effects of the movements (i.e., a spatial constancy feedback loop).¹¹ Whether a similar mechanism is operative in ET cases with head tremor deserves future investigation.

Acknowledgements

Supported by R01 NS39422, R01 NS042859, P30 ES09089 and RR00645 (General Clinical Research Center)(NIH, Bethesda, MD). We would like to thank Blair Ford M.D. for reading the manuscript and providing constructive comments.

References

1. Snowden JS, Craufurd D, Griffiths HL, Neary D. Awareness of involuntary movements in Huntington disease. *Archives of neurology* 1998;55:801–805. [PubMed: 9626771]
2. Vitale C, Pellecchia MT, Grossi D, et al. Unawareness of dyskinesias in Parkinson's and Huntington's diseases. *Neurol Sci* 2001;22:105–106. [PubMed: 11487181]
3. Zesiewicz TA, Encarnacion E, Hauser RA. Management of essential tremor. *Curr Neurol Neurosci Rep* 2002;2:324–330. [PubMed: 12044252]
4. Benito-Leon J, Louis ED. Clinical update: diagnosis and treatment of essential tremor. *Lancet* 2007;369:1152–1154. [PubMed: 17416247]
5. Louis ED, Zheng W, Applegate L, et al. Blood harmaline concentrations and dietary protein consumption in essential tremor. *Neurology* 2005;65:391–396. [PubMed: 16087903]
6. Louis ED, Ford B, Frucht S. Factors associated with increased risk of head tremor in essential tremor: a community-based study in northern Manhattan. *Mov Disord* 2003;18:432–436. [PubMed: 12671952]
7. Louis ED, Borden S, Moskowitz CB. Essential tremor centralized brain repository: diagnostic validity and clinical characteristics of a highly selected group of essential tremor cases. *Mov Disord* 2005;20:1361–1365. [PubMed: 16001407]
8. Louis ED, Rios E, Applegate LM, et al. Jaw tremor: prevalence and clinical correlates in three essential tremor case samples. *Mov Disord* 2006;21:1872–1878. [PubMed: 16941462]

9. Brandt JSM, Folstein M. The telephone interview for cognitive status. *Neuropsychi Neuropsychol Beh Neurol* 1988;1.
10. Lee AG, Brazis PW. Localizing forms of nystagmus: symptoms, diagnosis, and treatment. *Curr Neurol Neurosci Rep* 2006;6:414–420. [PubMed: 16928352]
11. Abadi RV, Whittle JP, Worfolk R. Oscillopsia and tolerance to retinal image movement in congenital nystagmus. *Invest Ophthalmol Vis Sci* 1999;40:339–345. [PubMed: 9950591]

Table
Demographic and clinical features of ET cases in each of the three ET case samples

TERTIARY-REFERRAL CENTER (N = 320)	
Age in years	67.4 ± 15.2
Women	165 (51.6%)
White race	301 (94.1%)
Total tremor score	18.9 ± 7.3
Tremor duration in years	22.9 ± 18.7
POPULATION-BASED STUDY (N = 106)	
Age in years	69.8 ± 18.4
Women	63 (59.4%)
White race	42 (39.6%)
Total tremor score	17.6 ± 6.5
Tremor duration in years	17.4 ± 19.7
BRAIN REPOSITORY (N = 170)	
Age in years	74.5 ± 9.5
Women	102 (60.0%)
White race	170 (100%)
Total tremor score	22.3 ± 6.9
Tremor duration in years	39.2 ± 20.2

Values are means ± SD and number (percentages)