Published in final edited form as:

Behav Brain Res. 2008 March 5; 187(2): 207–220. doi:10.1016/j.bbr.2007.09.009.

Gabrb3 gene deficient mice exhibit impaired social and exploratory behaviors, deficits in non-selective attention and hypoplasia of cerebellar vermal lobules: a potential model of autism spectrum disorder

Timothy M. DeLorey¹, Peyman Sahbaie¹, Ezzat Hashemi¹, Gregg E. Homanics², and J. David Clark^{3,4}

¹Molecular Research Institute, Palo Alto, CA 94303

²Departments of Anesthesiology and Pharmacology, University of Pittsburgh, PA 15261

³Veterans Affairs Palo Alto Health Care System, Palo Alto, CA 94304

⁴Stanford University School of Medicine, Department of Anesthesiology

Abstract

Objective—GABA_A receptors play an important regulatory role in the developmental events leading to the formation of complex neuronal networks and to the behaviors they govern. The primary aim of this study was to assess whether gabrb3 gene deficient (gabrb3^{-/-}) mice exhibit abnormal social behavior, a core deficit associated with autism spectrum disorder.

Methods—Social and exploratory behaviors along with non-selective attention were assessed in gabrb3^{-/-}, littermates (gabrb3^{+/+}) and progenitor strains, C57BL/6J and 129/SvJ. In addition, semi-quantitative assessments of the size of cerebellar vermal lobules were performed on gabrb3^{+/+} and gabrb3^{-/-} mice.

Results—Relative to controls, gabrb3^{-/-} mice exhibited significant deficits in activities related to social behavior including sociability, social novelty and nesting. In addition, gabrb3^{-/-} mice also exhibited differences in exploratory behavior compared to controls, as well as reductions in the frequency and duration of rearing episodes, suggested as being an index of non-selective attention. Gabrb3^{-/-} mice also displayed significant hypoplasia of the cerebellar vermis compared to gabrb3^{+/+} mice.

Conclusions—The observed behavioral deficits, especially regarding social behaviors, strengthens the face validity of the gabrb3 gene deficient mouse as being a model of autism spectrum disorder.

Keywords

Social behavior; cerebellar vermis; exploration; non-selective attention; nest building

Corresponding author: Timothy M. DeLorey, Ph.D., Molecular Research Institute, 1000 Elwell Court, Suite 105, Palo Alto, CA 94303, Tel: 650-210-0311, FAX: 650-210-0318, Email: E-mail: tim@molres.org.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Introduction

Autism spectrum disorder (ASD) is a complex neurodevelopmental disorder characterized by impaired social behaviors, stereotypical or restrictive behavioral patterns, and deficits in language and communication [31]. In addition to the three core deficits, a wide continuum of behavioral phenotypes have also been reported in association with ASD, including cognitive impairment, hyperactivity, epilepsy, motor deficits, attentional deficits, hypotonia and sleep disturbances [2,11,47,74]. To date, no single region of the brain or pathophysiological mechanism has yet been identified as being the cause of ASD. However, the cerebellum, frontal cortex, hippocampus, amygdala and the cerebello-thalamo-cortical circuit have all been implicated in ASD [68]. The neural underpinnings of ASD are also poorly understood, although, there is evidence for a strong genetic component [60], even though no candidate gene has been verified to date. Nevertheless, one part of the genome stands out, chromosomal region 15q11-q13, which has been strongly implicated in ASD via linkage and association studies [50,51,56,70]. Furthermore, maternal duplications of this region remain one of the most common cytogenetic abnormalities found in cases of idiopathic ASD, accounting for about 1-2% of the case reported [22]. In addition, deletion of this region results in either Angelman syndrome or Prader-Willi syndrome depending from which parent the deletion was inherited. Both syndromes often meet the diagnostic criteria for ASD as defined by DSM-IV [61,65, 69,73]. Within this chromosomal region there exists a gene cluster of GABA_A receptor subunit genes: GABRB3, GABRA5, and GABRG3, encoding for the GABA_A receptor subunits β_3 , α_5 , and γ_3 , respectively.

GABA_A receptors are heterooligomeric GABA-gated chloride channels constructed from eight classes of subunits exhibiting varying amino acid sequence homologies (α_{1-6} , β_{1-4} , γ_{1-3} , δ , ρ_{1-2} , π , ϵ and θ) that produce multiple GABA_A receptor isoforms with various GABA sensitivities and associated pharmacologies [14,58]. These subunits exhibit unique regional and temporal distribution within the Central Nervous System. During development GABA_A receptors play a role in proliferation, migration, and differentiation of precursor cells that orchestrate the development of the embryonic brain [9]. A developmental deficiency in any of these roles would adversely effect the temporal ordering of neurogenesis and synaptogenesis, thereby affecting maturation of circuits that are later involved in complex behaviors.

The high prevalence rate of ASD, 1 out of 150 births [20] has prompted an urgency to develop animal models as a fundamental step toward comprehending the complex molecular underpinnings associated with this disorder. A suitable animal model should meet three fundamental criteria 1) *face validity*: behavioral characteristics should mirror those present in the human disorder; 2) *construct validity*: similarities in the underlying etiology should exist between the human disorder and the animal model; and 3) *predictive validity*: the outcome of a treatment regime applied to the animal model should reflect the likely impact on humans with the disorder. While no model can be expected to replicate the full complexity of the human behavioral phenotype, an animal model that exhibits specific behavioral and morphological characteristics typically associated with ASD would be invaluable.

Numerous studies, employing a diverse set of approaches including autoradiographic, molecular biological, and genetics, provide overwhelming support for the role of GABAergic mechanisms in the etiology of ASD [13,17,38,70]. In light of these observations, mice that have a targeted disruption of the mouse equivalent of the human GABRB3 gene, which encodes the β_3 subunit of the GABA_A receptor, can be argued to possess construct validity in reference to ASD. In addition, gabrb3 gene disrupted (gabrb3^{-/-}) mice exhibit numerous behavioral abnormalities, including many often reported in association with ASD [26,27,39]. In this present study we sought to further extend the face validity inherent to the gabrb3^{-/-} mouse relative to behaviors associated with ASD. Given the primacy of its deficit in ASD, special

emphasis was placed on the assessment of social behavior. In addition, we assessed other behaviors often impaired in ASD, including exploratory behavior and non-selective attention. Lastly, as the cerebellar vermis has been cited as being abnormal in ASD [64] we performed a simple morphological assessment of this region in *gabrb3*^{-/-} and *gabrb3*^{+/+} mice. In addition, this region has been reported to be crucial to the consolidation of aversely motivated contextual memory, an elementary form of spatial learning [66] and a feature previously found to be disrupted in *gabrb3*^{-/-} mice [27]. The cerebellum has also been reported to be involved in exploratory behaviors [62], shifting attention [1] and spatial orientation [41], features likely to impact the behaviors being assessed within the current study. In light of the overwhelming evidence implicating the GABAergic system in ASD and the numerous parallels between the ASD phenotype and the phenotype observed in *gabrb3*^{-/-} mice, this timely study provides a crucial connection between the disruption of the *gabrb3* gene and the impairment of social behavior, a key diagnostic component of the ASD phenotype. Insights gained from the current efforts aid in bridging gaps in our understanding of the interconnectiveness between genetics, development and behavioral outcome.

Methods

Mice

All mice used in this study were male, consistent with the 4:1 male-female ratio prevalent in autism [29]. *Gabrb3* gene knockout mice (*gabrb3*^{-/-}) and wild-type littermates *gabrb3*^{+/+} were produced at the University of Pittsburgh or the Veterans Affairs Palo Alto Health Care System. Techniques used to disrupt the *gabrb3* gene have been previously described [39]. C57BL/6J and 129/SvJ mice were obtained from Jackson Laboratory (Davis, CA) at 6-8 weeks of age. All animal protocols conformed to the guidelines determined by the National Institute of Health (USA) Office for Protection from Research Risks and were approved by the Animal Care and Use Committee of the Veterans Affairs Palo Alto Health Care System. All mice were housed in groups of 4-8 on a 12/12-h light/dark cycle under controlled temperature with food and water provided *ad libitum*. A number of studies report that the phenotype of a given single gene mutation is modulated by the genetic background of the inbred strain(s) in which the mutation is maintained (for review see [24]). As the genetic background of the *gabrb3*^{+/+} and *gabrb3*^{-/-} mice is a hybrid between 129Sv/SvJ and C57BL/6J mice, similar progenitor strains were also assessed in this study alongside the wildtype and knockout mice to assure that observed differences were not merely the result of strain variations within the two progenitor strains. The age of mice used in this study ranged from 16-52 weeks. The precise age range used for each assessment is listed under each method subsection. Due to a 90% mortality rate at birth of mice lacking the *gabrb3* gene [39], the number of *gabrb3*^{-/-} mice available to pursue the goals of this study represent the minimal amount of mice required to perform adequate statistical analysis. This low survival rate was a contributing factor in the wide age range of mice available for this study. In an effort to reduce the impact of age differences on the results, we age matched all controls to the available *gabrb3*^{-/-} mice.

Social interaction and preference for social novelty

The social interaction apparatus used in this study was a modified conditioned place preference apparatus (Med Associates Inc., St. Albans, VT), consisting of a polycarbonate box with two partitions dividing the apparatus into three separate chambers (two interaction chambers and one neutral chamber). Attached to each interaction chamber was a stimulus cage separated from the interaction chamber by a wire mesh wall with a 1 cm² grid mesh (Fig 1). With the partitions open, a test mouse is able to move freely between chambers. Partitions are automatically opened and closed by a computer controlled interface (Med PC for Windows) and the output interface ENV-313/BD (Med Associates Inc.). The three chambers were equipped with infrared photo-sensors spaced 2 cm apart and 1 cm above the chamber floor

with data collected via a Med Associates input module ENV-2561 and interface SG-6080/C. Both interaction chambers are identical in size, shape, color, floor texture and lighting with removable flooring for cleaning between experiments. The neutral chamber is similar in color and floor texture to the interaction chambers so as not to be aversive. During testing, “stranger” mice were placed in one or both stimulus cages. The wire mesh separating the interaction chamber from the stimulus cage allows nose contact between test mouse and stranger mouse, while preventing aggressive interactions and also prevents the “stranger” mouse from initiating the social contact.

Sociability—All test mice were experimentally naïve to the social interaction test and were habituated to the test room for at least 30 min prior to assessment. Testing was done on all four mouse genotypes: C57BL/6J, 129/SvJ, *gabrb3*^{-/-} and *gabrb3*^{+/+}. The age range of the mice tested in these social assessments were between 16-52 weeks of age. Stranger mice were male C57BL/6J housed in groups of four per cage. The experimental design of both the social interaction and social novelty tests were adapted from Nadler et al, [53,54]. Briefly, baseline adaptation was collected by placing the test mouse in the central neutral chamber for a two min habituation period, followed by the partitions being raised and allowing the mouse to explore freely all three chambers (two interaction chambers and the central neutral chamber) for a 10 min period with data being collected in five min bins. During baseline assessments there were no “stranger” mice in either of the stimulus cages. Time spent in each chamber was used as baseline data to assess general activity, willingness to circumnavigate the testing apparatus and to assure that no mouse exhibited a side preference. Both ambulatory locomotor activity and the total number of entrances (transitions between chambers of the apparatus) during the 10 min baseline period were assessed using the photo-sensors in the chamber walls. An exclusion criterion required each mouse to explore each of the three chambers for a minimum of 90 sec (15 % of the total test time) each during the initial 10 min exploratory period. Mice that failed to meet this criterion were eliminated from further consideration. Following baseline data collection, each mouse was returned to the neutral central chamber for two min with partitions closed. During this time, an unfamiliar adult C57BL/6J mouse (Stranger #1) was placed within one of the stimulus cages that were attached to each interaction chamber. Again partitions were raised, allowing the test mouse to move freely throughout the three chambers for a 10 min test session. Location of the ‘stranger mouse’ was alternated between the left and right stimulus cages on consecutive sessions. Again, measurements were collected in five min time bins and broken down into time spent in each the three chambers (the interaction chamber adjacent to a stimulus cage containing ‘stranger mouse #1’, interaction chamber adjacent to an empty stimulus cage, and the neutral central chamber). Also measured was the number of entries made into each interaction chamber.

Preference for social novelty—To quantify preference for social novelty, a second 10 min test session was performed immediately following the 10 min sociability test. The original unfamiliar mouse (Stranger mouse #1) remained in its stimulus cage while a second unfamiliar mouse (stranger mouse #2) was added to the second stimulus cage, which was previously empty. Stranger #1 and stranger #2 mice originated from different home cages and had never come into physical contact with each other or with the test subject. After the partitions were again raised, automated measurements were collected in an identical fashion to the social interaction test.

To assure that the time spent in the interaction chamber adjacent to the stimulus cage containing an unfamiliar mouse was social in nature, a blinded observer scored, from videotape, each test session for explorations or sniffing directed towards a “stranger mouse” (nose within 2 cm of the wire mesh wall and directed toward the stranger mouse). This data was then compared to exploration and sniffing behavior exhibited by the test mouse towards the wire mesh wall separating the 2nd interaction chamber from a stimulus cage that contained no mouse. The

observer scores were compared to the automated scoring. The cumulative scores for the 10 min observer scored test sessions were expressed as total duration of direct social interactions in seconds.

Data Analysis—Baseline locomotor activity, number of entries to and the time spent in side chambers was examined by an overall repeated measure ANOVA for the four genotypes during the adaptation period followed by Bonferroni *post hoc* multiple comparison tests. The social data were initially analyzed by an overall repeated measures ANOVA with the factors of genotype and side preference (e.g. “stranger mouse #1” side versus the empty side in the sociability test and “stranger mouse #1” side versus stranger mouse #2” side in the preference for social novelty test). Subsequently, statistical significance was determined by within group repeated measures ANOVA with Bonferroni *post hoc* multiple comparison tests to determine side preferences following a significant overall ANOVA. All statistical analyses were performed using GraphPad Prism 4.0 (GraphPad, San Diego, CA).

Nest building and utilization

Prior to evaluation of nest building, each mouse was housed individually for one hr in a clean cage with fresh sawdust bedding. The age of the mice used in these assessments ranged between 16-52 weeks of age. The observation period took place during the light cycle and was initiated by the introduction of a nestlet, a 2.5 gr, 5.5 cm² wafer of compressed cotton (Ancare, Bellemore, NY) into the cage. Assessment of the total amount of time spent building a nest during the first hr was performed during three separate 10 min epochs each separated by 10 min. Time building the nest included the amount of time a mouse was involved in shredding the nestlet using either its mouth or forepaws as well as transporting the material around the cage. After the first hr a second nestlet was provided by hanging the nestlet from a clip attached to the food hopper and after 24 hours a third nestlet was introduced in the same manner. The amount of nest material utilized after 1, 24 and 48 hr of exposure was assessed by subtracting the weight of unutilized nestlet material from the total weight of material made available to the mouse. After 48 hr, the nest quality was scored and the nest location recorded [16,46,52, 72]. Nest quality was scored according to the following scale modified from Moretti et al., [52]: 1=nestlets largely unmodified; 2=nestlets modified (shredded) but no distinct nest shape; 3=flat nest with partially shredded nestlet and large segments of the nestlet wafer remaining intact; 4=shallow nest with partially shredded nestlets; 5=shallow nest with well shredded material; 6=nest with well developed walls; 7=nest in a shape of a cocoon with small opening of a diameter of <3 cm or complete roof. Nest building time and nestlet usage were analyzed for statistical significance by a one-way ANOVA with Bonferroni *post hoc* multiple comparison test. Nest quality scores were analyzed using Kruskal-Wallis nonparametric one-way ANOVA with Dunn's *post hoc* multiple comparison test. Lastly, during the light period on three separate days, the resting location of each mouse, relative to the location of the nest was noted on three separate occasions throughout the day.

Exploratory behavior

Methods used for assessing exploratory behavior were adapted from ‘open field’ protocols [30]. Briefly, mice in the age range of 24-52 weeks were placed individually into a brightly lit circular arena (diameter 54 cm, height 25 cm) for a period of 1 hr in order to acclimate to the novel environment. Twenty-four hours later each mouse was returned to the open field arena that now contained a novel object (circular mesh cage 3” in diameter and 2” in height) placed in the center of the open field chamber. During the 10 min test period several assessments were manually assessed including the time it took to first make contact with the object, the total time spent investigating the object, the number of contacts with the object and the total number of times each mouse reared. A rearing episode was defined as either rearing on hindlimbs or leaning with one or both forepaws against the wall or the novel object placed within the open

field chamber. Instances in which the mouse sat back on its hindlimbs in order to groom itself were not considered a rearing event. The frequency and duration of rearing episodes, suggested as being a measure of non-selective attention, were also assessed according to methods adapted from Asipe et al., [5]. Briefly, recordings taken during the other exploratory assessments were visually monitored in 1-min time blocks in order to determine the rearing frequency of each mouse when placed in a novel environment. These values were averaged within genotype for each 1-min time block. Subsequently, the mean duration of each rearing episode was also assessed for each mouse and averaged within genotype. The open field chamber was cleaned with a wet sponge in between trials. Each session was videotaped in order to verify all assessments and to produce diagrams of the exploratory path that each mouse pursued. Data were analyzed for statistical significance by a repeated measures ANOVA with Bonferroni *post hoc* multiple comparison test following a significant ANOVA. Differences in rearing frequency were analyzed by a two-way ANOVA genotype $\times$ time block, as the dependent variable.

Brain preparation

Mice from both the *gabrb3*^{-/-} and *gabrb3*^{+/+} groups were sacrificed between the ages of 51-52 weeks by CO₂ inhalation followed by rapid transcardial perfusion with 20 ml of 0.1M phosphate buffered saline (PBS) and then 20 ml of 4% paraformaldehyde (PFA) in 0.1M phosphate buffer (PB). Perfusion was performed using a 22-gauge 1 inch needle inserted into the left ventricle with a small cut made in the right atrium for fluid to exit the circulatory system. Following perfusion, brains were carefully removed from the skull and incubated for 5 hrs in 4% PFA and then cryoprotected by incubation in 30% sucrose in 0.1M PB overnight at 4°C. Brains were placed in a 1mm mouse brain matrix (Ted Pella, Redding CA) and separated along the midline using a razor blade. Both the right and left hemispheres were frozen separately in M1 imbedding agent (Shandon Lipshaw Inc., Pittsburgh, PA) at -20 °C and then stored at -80° C until used. Prior to sectioning, brains were moved to a freezer and kept at -20°C for 15 hr. Each hemisphere was then mounted on a specimen holder of a freezing stage microtome (Leica, CM1850) that was kept at -15°C. Tissue was cut in serial sagittal sections from the mediosagittal line at a thickness of 40 μ m then thaw-mounted on room temperature Superfrost plus slides (Fisher Scientific, Pittsburgh, PA) and dried for 4 hr at room temp. Slide mounted tissue sections were stained for 20 min with toluidine blue (0.5%, in 1.0% sodium tetraborate w/v in deionized H₂O) then destained by 15 dips in 70%, 90% and 100% ethanol followed by four xylene washes lasting three min each. Slides were then cover-slipped using Permount (Fisher Scientific, Pittsburgh, PA).

Images of the cerebellar vermis from each section were taken using a SPOT RT slider camera (software v4.0.9, Diagnostic Instruments Inc, Sterling Heights, MI) mounted on an Olympus BHTU-BH2 microscope (Scientific Instrument Company, Sunnyvale, CA). Images were superimposed onto a corresponding diagram of a sagittal section of mouse cerebellar vermis taken from Paxinos and Franklin [59] at the level of the vermis between 0.36-0.60 mm lateral from the midline. Three separate sections were examined per animal (right or left vermal lobules) with surface areas being averaged per animal then pooled within genotype (*gabrb3*^{-/-} and *gabrb3*^{+/+}). Semi-quantitative measurements were done at a magnification of 4 $\times$ by two independent observers that were blind to the mouse genotype. The entire lobule (molecular layer, granule layer and white matter) area (see Fig. 9 for example) was included in the measurement. Statistical significance was tested using a two-tailed Student's *t*-test (Prism 4, GraphPad, San Diego CA).

Results

Social behavior

A repeated measures ANOVA of the baseline locomotor activity of the four genotypes, *gabrb3*^{-/-} (n=6), *gabrb3*^{+/+} (n=6), C57Bl/6J (n=6) and 129/SvJ (n=6) mice indicated a significant effect of genotype ($F_{3,20}=9.55$, $p<0.001$), with *gabrb3*^{-/-} mice displaying a significantly higher degree of ambulatory activity than control mice (Table 1), in agreement with previous reports of hyperactivity in *gabrb3*^{-/-} mice [39]. In addition, following a significant ANOVA ($F_{11,60}=11.95$, $p<0.001$), *gabrb3*^{-/-} mice were found to exhibit a significantly higher number of entries relative to control mice in regards to the total transitions between all chambers of the social interaction apparatus during the adaptation period (Table 1). All mice, regardless of genotype, were found not to exhibit a significant difference between the amount of time spent in either interaction chamber ($F_{1,40}=0.52$, $p=0.48$) or the number of entries into either chamber ($F_{1,40}=0.04$, $p=0.84$), thereby demonstrating a lack of side preference (Table 1). Additionally, the number of entries made to either side during the 10 min sociability or social novelty assessments were likewise found not differ significantly within each mouse genotype (data not shown), similar to what has been previously reported [54]. The initial five min test epoch tended to be more informative than the second five min epoch, in regards to the duration of the interaction in both the sociability and social novelty assessments, as has been previously reported [54]. In order to verify both the robustness of the automated data collection procedure and to verify that the data reflected interactions that were social in nature, a blind observer also evaluated the interactions from videotape. Due to the fragility of the *gabrb3*^{-/-} mouse genotype (see methods) there was a wider than normal age distribution, three mice in the 16-22 weeks age range and the remaining three in the 50-52 week range. Therefore, within genotype analysis was also performed, in regard to each assessment, in order to determine whether the age difference was likely to influence the outcome of the assessment. Differences between the two age groups were not significant.

Automated Measurements of the Sociability Test—In agreement with Nadler et al. [54], we found that the number of chamber entries was independent of the time spent in the chambers, regardless of whether the chamber was adjacent to a cage containing an unfamiliar mouse or was empty. An overall repeated measures ANOVA of the initial five min period revealed a significant difference for side between the four strains; $F_{1,40}=62.66$, $p<0.001$. On further analysis it was found that there was no significant difference between the three control strains on measures of side duration ($F_{2,30}=0.65$, $p>0.05$). Control strains were found to exhibit a significant preference for spending time in the interaction chamber adjacent to the stimulus cage containing an unfamiliar mouse “stranger mouse #1” as opposed to spending time in the interaction chamber adjacent to an empty stimulus cage, C57Bl/6J: $F_{5,25}=6.55$, $p<0.01$; 129/SvJ: $F_{5,25}=5.21$, $p<0.05$ and *gabrb3*^{+/+}: $F_{5,25}=2.82$, $p<0.01$ (Fig. 2). Similar results have been previously reported for C57BL/6J mice in this task [54]. In contrast, the time *gabrb3*^{-/-} mice spent, during the first five min epoch, in either the chamber adjacent to the cage containing “stranger mouse #1” or the chamber adjacent to an empty stimulus cage was found not to differ significantly ($F_{5,25}=1.92$, $p>0.05$; Figure 2).

Automated Measurements of the Preference for Social Novelty Test—Ten min after exposure to stranger mouse #1 a second unfamiliar mouse “stranger mouse #2” was placed into the previously empty stimulus cage adjacent to interaction chamber #2 (see Fig. 1). An overall repeated measures ANOVA of the first five min epoch revealed a significant difference for side between the four strains; $F_{1,40}=52.5$, $p<0.001$. There was no significant difference between the three control strains on measures of side duration during the initial five min test epoch; $F_{2,30}=0.27$, $p>0.05$. *Post hoc* tests revealed the control strains to exhibit a preference for interacting with the new mouse (stranger mouse #2), as opposed to the now familiar mouse

(stranger mouse #1), C57BL/6J: $F_{5,25}=13.58$, $p<0.001$, 129/SvJ: $F_{5,25}=14.11$, $p<0.01$ and $gabrb3^{+/+}$: $F_{5,25}=7.19$, $p<0.001$ (Fig. 3). This preference was also detectable in the second five min test epoch in both the C57BL/6J and 129/SvJ mouse strains, but not in $gabrb3^{+/+}$ mice (Fig. 3). $Gabrb3^{-/-}$ mice failed to display a significant preference for the unfamiliar mouse over the now familiar mouse in either test epoch ($F_{5,25}=0.70$, $p>0.05$; Fig. 3). Furthermore, $gabrb3^{-/-}$ mice exhibited an increased preference for the middle/neutral chamber (175.3 ± 55.4 , mean $\pm$ SEM) compared to the control strains $gabrb3^{+/+}$ (115.3 ± 25.5), C57BL/6J (97.3 ± 32.5) and 129/SvJ (60.7 ± 11.4) during the 10 min social novelty test, though the difference between genotypes was not found to be statistically significant (compare grey bars in Fig. 3).

Observer scoring of the Sociability and Preference for Social Novelty Tests—

Analysis of the observer scored data during the 10 min test period of the sociability tests was found to agree favorably with the automated data for both control and $gabrb3^{-/-}$ mice (compare Fig. 3 and Fig. 4). Similar agreement between the automated and observer scored sociability data has been previously reported in C57BL/6J mice [54]. *Post hoc* tests revealed significant differences in the amount of time mice spent sniffing at the cage containing Stranger mouse #1 versus the empty stimulus cage for C57BL/6J: $F_{1,5}=11.07$, $p<0.05$; 129/SvJ: $F_{1,5}=16.78$, $p<0.01$ and $gabrb3^{+/+}$: $F_{1,5}=15.62$, $p<0.05$, but not for the $gabrb3^{-/-}$ mice: $F_{1,5}=1.10$, $p>0.05$ (Fig. 4).

Comparison between the automated and observer scoring of the social novelty test again found the control genotypes to behave in a similar manner under both assessments. However, subtle differences in the performance of $gabrb3^{-/-}$ mice were detected by observer scoring that were not detected by the automated scoring method. Whereas, the amount of time $gabrb3^{-/-}$ mice spent in the proximity of either stranger mouse #1 or stranger mouse #2 was not found to be significantly different when assessed by the automated method (Fig. 3), the observer scoring, which only takes note of direct interactions (nose within 2 cm of the wire mesh of the stimulus cage containing stranger mouse #1 or #2), revealed a significant difference in the amount of time involved in direct interactions with stranger mouse #2 versus stranger mouse #1 (Fig. 4). *Post hoc* tests indicated a significant preference for sniffing at Stranger mouse #2 over Stranger mouse #1, C57BL/6J: $F_{1,5}=64.48$, $p<0.001$; 129/SvJ: $F_{1,5}=29.29$, $p<0.01$, $gabrb3^{+/+}$: $F_{1,5}=20.68$, $p<0.01$ and $gabrb3^{-/-}$ mice: $F_{1,5}=20.16$, $p<0.01$. Upon closer examination of the observer data, in regards to the percentage of the total time spent sniffing the “stranger mice”, the nature of the contrasting results for the $gabrb3^{-/-}$ mice between the automated versus the observer scored methods was revealed. By examining the total time that each mouse genotype spends interacting with stranger mouse #1 in the sociability test, you will note a markedly reduced level of sniffing of stranger mouse #1 by $gabrb3^{-/-}$ mice in comparison to the three control strains ($F_{3,15}=5.38$, $p<0.01$, Fig. 5A). Observer assessment of the social novelty test reveals a similar trend in respect to the total amount of time $gabrb3^{-/-}$ mice spend sniffing at stranger mouse #2 compared to the time the control strains spend sniffing at stranger mouse #2 ($F_{3,15}=13.65$, $p<0.001$, Fig. 5B). Moreover, when considering total sniffing directed at both stranger mouse #1 and stranger mouse #2 in the social novelty test, again $gabrb3^{-/-}$ mice clearly displayed a significant reduction in the overall amount of time spent in sniffing directed at either stranger mouse #1 or #2, compared to control genotypes ($F_{3,15}=20.04$, $p<0.001$, Fig. 5C). Thereby revealing $gabrb3^{-/-}$ mice to exhibit low levels of social engagement towards other mice.

Nesting behavior was likewise assessed as it represents a form of homecage-activity often linked to social behavior [52]. The amount of time mice spent interacting with nesting material during the first hour of exposure to nestlets, was analyzed by a repeated measures ANOVA, which indicated strain differences: $F_{3,26}=3.73$, $p<0.05$ with *post hoc* tests revealing significant differences between $gabrb3^{-/-}$ mice and the control genotypes (Fig. 6A). Although all mouse genotypes demonstrated an increase in material usage over the 48 hr observation period, a

repeated measures ANOVA again revealed a significant strain difference in nestlet material usage at the 1 hr, 24 hr and 48 hr test intervals: $F_{3,26}=8.32, p<0.001$; $F_{3,26}=44.53, p<0.001$; $F_{3,26}=13.95, p<0.001$ respectively with *post hoc* tests revealing *gabrb3*^{-/-} mice to differ significantly from controls at all three time-intervals (Table 2). The nest quality scores (see methods for scoring) were assessed 48 hr after nestlets were first introduced to the cages. A Kruskal-Wallis analysis revealed a significant variation between mouse genotypes ($\chi^2=18.47, p<0.001$) with *post hoc* tests indicating *gabrb3*^{-/-} mice to be significantly impaired in their ability to achieve the nest complexity exhibited by the control genotypes (Fig. 6B). Whereas, control mice typically built well-assembled walls or a cocoon-like nest from fully shredded nestlets, *gabrb3*^{-/-} mice generally built flat nests consisting of mostly unshredded nesting material. It is noteworthy to mention that all control mice built their nests in either a corner or along the cage wall, while *gabrb3*^{-/-} mice often built their nests away from the corners and walls, if they constructed one at all. Utilization of the nests during resting periods also differed between control and *gabrb3*^{-/-} mice: the controls were found resting in their nest in 100% of the observations taken over a 3 day period, but the *gabrb3*^{-/-} mice were observed sleeping away from the nesting material in 50% of the observations.

Exploratory behavior

Figure 7 depicts representative examples of the exploratory paths followed by a *gabrb3*^{+/+} and a *gabrb3*^{-/-} mouse. The higher overall line density/length of the *gabrb3*^{-/-} mouse, compared to the *gabrb3*^{+/+} mouse, is reflective of the hyperactive nature of *gabrb3*^{-/-} mice. The relative proportion of the path density enclosed within the inner open circle, containing the novel object, is qualitatively higher in the *gabrb3*^{+/+} mouse compared to the *gabrb3*^{-/-} mouse and is associated with the significantly longer time *gabrb3*^{+/+} mice spent investigating the novel object compared to *gabrb3*^{-/-} mice (Table 3). These exploratory path diagrams also reveal the stereotypical circling pattern common to *gabrb3*^{-/-} mice [39] (see arrow, Fig. 7) that was not observed in the more random locomotor pathways exhibited by *gabrb3*^{+/+} mice.

Significant genotype differences were detected in both the time that elapsed before mice first made contact with a novel object placed in the open field chamber ($F_{3,26}=8.55, p<0.001$) and the amount of time mice spent investigating the novel object ($F_{3,26}=36.9, p<0.001$). *Post hoc* analysis revealed significance difference between *gabrb3*^{-/-} mice and control genotypes in both assessments (Table 3). In addition, the number of contacts mice made with the novel object also revealed an overall significant effect of strain; $F_{3,26}=19.4, p<0.001$ with *post hoc* tests indicating statistically that only the 129/SvJ mice differed significantly from *gabrb3*^{-/-} mice (Table 3).

The total number of rearings, another indicator of exploratory behavior [12], was also found to exhibit a significant strain difference; $F_{3,25}=8.57, p<0.001$, with *post hoc* tests demonstrating a significance difference between *gabrb3*^{-/-} mice and both *gabrb3*^{+/+} and C57Bl/6J mice, but not 129/SvJ mice (Table 3). As shown in Fig. 8A *gabrb3*^{-/-} displayed a lower frequency of rearing activity, as compared with the corresponding control mice. A two-way ANOVA revealed a significant effect of genotype in the frequency of rearing ($F_{3,225}=6.84, p<0.01$) with no genotype $\times$ time block interaction. A statistically significant reduction in the mean duration of a rearing event was also noted in the *gabrb3*^{-/-} mice compared to the control genotypes (Fig. 8B) following a significant repeated measures ANOVA in regards to duration; $F_{3,25}=16.64, p<0.001$.

Cerebellar vermis

Semi-quantitative measurements of the sagittal surface area of cerebellar vermal lobules (Fig. 9) of *gabrb3*^{+/+} and *gabrb3*^{-/-} mice revealed significant decreases in the surface areas of vermal lobules II, III, IV-V and VI-VII in *gabrb3*^{-/-} mice compared to *gabrb3*^{+/+} (Fig. 10). A non-

significant reduction in the surface areas of vermal lobules VIII, IX and X in *gabrb3*^{-/-} mice compared to *gabrb3*^{+/+} mice, were also noted (Fig. 10).

Discussion

The GABRB3 gene is vital to proper brain development and to mature brain function. The consequences of disrupting this gene have been demonstrated in mice, which exhibit numerous abnormalities, many of which have been observed in association with the neurodevelopmental disorders Angelman syndrome and ASD [26,27,39]. The present study provides a crucial addition to this body of work by demonstrating that *gabrb3*^{-/-} mice exhibit deficits in social behavior, a core feature of ASD. In addition, other inclinatory traits often associated with ASD were also detected in these mice, including hypoplasia of cerebellar vermal lobules and deficits in both exploratory behaviors and non-selective attentional processes related to the orientation of attention [5,35,64].

Gabrb3^{-/-} mice clearly exhibited social deficits, as indicated by a significant reduction in overall social engagement in both sociability and social novelty testing, however, some aspects of social curiosity appear to remain intact. While *gabrb3*^{-/-} mice spent less time in the proximity of unfamiliar mice than did their control counterparts (see Fig. 4), we were able to discern from closely examining the direct interactive contact that *gabrb3*^{-/-} mice exhibited towards either of the two Stranger mice in the social novelty test, that there was a preference for the novel mouse over the now familiar mouse (compare Figure 3 with Figure 4). The discrepancy between the two measurements can be partially explained by noting that the automated scoring method collects data on total time spent in all three chambers including the central/neutral chamber, whereas the observer scoring method only collects data pertaining to direct social engagement (nose touching or sniffing within 2 cm of the unfamiliar mouse) as opposed to just being in the social proximity of the unfamiliar mouse. If one examines the total amount of direct social engagement that each mouse genotype exhibits, one easily discerns that *gabrb3*^{-/-} mice displayed an attenuated level of interaction towards unfamiliar mice in general, compared to controls (Figure 5). We also observed differences in the nesting behavior of *gabrb3*^{-/-} mice, a behavioral feature associated with abnormalities in social behavior [52] and that has likewise been observed in other mouse models linked to ASD, including *Mecp2* gene mutation mice, a model of Rett syndrome and in mice deficient in the *Dvl1* gene [46,48,52].

As a rodent's social behavior is highly dependent on olfactory cues, one must consider whether olfaction differences between *gabrb3*^{-/-} and *gabrb3*^{+/+} may have contributed to the observed differences in social behavior. Indeed, *gabrb3*^{-/-} mice have been reported to exhibit subtle differences in olfactory discrimination, however, *gabrb3*^{-/-} mice were found to exhibit better odor discrimination than *gabrb3*^{+/+} mice in certain tasks [57]. We also noted that *gabrb3*^{-/-} mice were as adept as *gabrb3*^{+/+} mice in locating a buried piece of cookie (data not shown). Although these observations do not eliminate the possibility of an olfaction confound, they do shed doubt on olfactory irregularities as being the primary explanation for the observed social impairment exhibited by *gabrb3*^{-/-} mice. Taken together, the deficits that *gabrb3*^{-/-} mice exhibit in regards to sociability, social novelty, nest building and previous observations that *gabrb3*^{-/-} mouse mothers fail to care for their young [39], add up to strongly indicate that normal social behavior is significantly impaired in *gabrb3*^{-/-} mice.

Gabrb3^{-/-} mice were also found to exhibit significant deficits in a variety of exploratory parameters. The diagrams depicted in Figure 7 represent a typical path that either a *gabrb3*^{+/+} or *gabrb3*^{-/-} might pursue. Differences between *gabrb3*^{+/+} and *gabrb3*^{-/-} mice can be gleaned from such diagrams, including the hyperactivity typically exhibited by *gabrb3*^{-/-} mice (high path density in outer region), low exploration of the novel object (low path density in the inner ring) as well as the stereotypical circling pattern typical of these mice (see arrow Fig. 7).

Whereas, *gabrb3*^{+/+} mice exhibit the opposite pattern, lower path density in the outer region and a higher path density in the inner ring containing the novel object, reflecting the higher amount of time spent investigating the object. Compared to control mice, *gabrb3*^{-/-} mice took longer to initiate the investigation of a novel object placed in the center of the testing chamber and spent less time devoted to exploring the novel object. Whereas, the total number of contacts a mouse made with the novel object did not appear to be as informative as the total amount of time the mouse actually spent investigating the object (Table 3).

Another component of exploratory behavior is rearing, which is associated with a rodent's motivational state and arousal level when exposed to a novel environment [67]. Consequently, novel stimuli, in all sensory modalities, are highly effective in attracting and focusing attention. Significantly less rearing occurred in *gabrb3*^{-/-} compared to *gabrb3*^{+/+} and C57BL/6J, but not 129/SvJ mice (Table 3). However, the duration of an average rearing event in 129/SvJ mice was observed to be much longer than in the other genotypes (Figure 8B), therefore, potentially accounting for the lower rearing number. On the whole, 129/SvJ mice appeared to be more inquisitive, in regards to the exploratory parameters assessed, than the other mouse genotypes. *Gabrb3*^{-/-} mice also exhibited a lower rearing frequency and lower mean rearing duration compared to control mice (Fig. 8 A and B). Studies conducted by Aspide et al., [5,6] have demonstrated that the frequency and duration of rearing episodes exhibited by a rodent placed in a novel environment also indexes the scanning and orienting phase of attention (non-selective attention). The significant reduction in both the frequency and duration of rearing episodes exhibited by the *gabrb3*^{-/-} mice, relative to controls (Fig. 8A & B) suggests *gabrb3*^{-/-} mice exhibit a deficit in non-selective attention. This is of interest as individuals with ASD often display deficits across many attentional domains, including selective and sustained attention with the inability to shift and orient attention rapidly and accurately among spatial targets and between sensory modalities [2,35,75]. This attentional deficit likely contributes to an autistic individuals reduced tendency and motivation to thoroughly explore novel environments [64] and has been suggested to be a contributing factor in the limited social interactions that autistic individuals display toward others [31].

In addition to the above behavioral deficits, *gabrb3*^{-/-} mice have been previously found to be poor swimmers, exhibit difficulty walking on grid floors, often run in circles and do not perform as well as controls on the accelerating rotarod [27,39]. Therefore, one possible explanation for the previously observed deficits and reduction in rearing, may involve changes to the vestibular system of the inner ear, which controls balance and orientation in space. Subsequently, two studies have reported changes in components of the vestibular system of the *gabrb3*^{-/-} mouse [42,49]. While mindful of these observations, the current assessments made were not dependent on high levels of motor coordination or postural control. Furthermore, upon close observation of *gabrb3*^{-/-} mice, they were found to be quite capable of rearing and able to groom themselves for extended periods while sitting back on their hindlimbs. Therefore, the moderate motor coordination and potential postural control issues associated with *gabrb3*^{-/-} mice may contribute, but are unlikely to be the primary cause of reduced rearing behavior. Another potential confound to the behavioral assessments performed on *gabrb3*^{-/-} mice is that these mice are prone to seizures [27,39]. Therefore, an accumulated seizure load, especially in the older *gabrb3*^{-/-} mice could potentially contribute to the observed behavioral differences as well as differences in brain morphology. We are not able to comment on whether there would be differences in the cerebellar hypoplasia between younger and older *gabrb3*^{-/-} mice as the cerebellar assessments were all done on mice between 51-52 weeks of age. However, younger *gabrb3*^{-/-} mice were found to perform in a similar fashion to older *gabrb3*^{-/-} mice in regards to the behaviors assessed in this study (data not shown). Furthermore, as demonstrated in the *gabrb3*^{-/-} mouse, GABAergic deficits leads to numerous phenotypic traits including epilepsy, therefore, one would anticipate that ASD cases that exhibit GABAergic deficits may likewise present with epilepsy. Subsequently, about 25% of the ASD population exhibit epilepsy [34]

and would also be subject to an accumulated seizure load as they age. This increasing seizure load could also potentially influence the manifestation of the ASD phenotype over the lifetime of this population.

A variety of neuroimaging studies suggest that the cerebellum is involved in more than just motor coordination, but also in a diverse set of higher cognitive functions [45], many of which have been observed to be affected in ASD. These include the processing of spatial information [44,63], spatial orientation [41], exploratory behavior [62] and shifting attention [1]. The cerebellar vermis, especially lobules VI-VII, have been reported as being abnormal in ASD [64]. Semi-quantitative assessments of the cerebellar vermal regions of *gabrb3*^{-/-} mice relative to *gabrb3*^{+/+} mice found significant reductions in the surface areas of lobules II-VII (Fig. 10) with non-significant reductions occurring in lobules VIII-X. As large sections of lobules IX and X mature earlier than lobules VI-VII, the current observation suggests that the developmental aberrations leading to reductions in the surface areas of lobules VI-VII in *gabrb3*^{-/-} mice may have, in part, occurred later in vermal development [4,40]. Similarly, the L1 cell adhesion molecule knockout mouse, which also exhibit hypoplasia of the cerebellar vermis, especially in lobule VI, also displays a reduced tendency to explore unfamiliar environments, novel objects placed within their environments, are in continual motion and display stereotypical circling [30]. It is not clear what the circling pattern in these mice or *gabrb3*^{-/-} mice represents, though it is reminiscent of the peripheral circling exhibited by rodents with induced cerebellar lesions. Moreover, it has been noted that there is a proclivity towards spinning (circling) in the ASD population [15]. The observed hypoplasia of the cerebellar vermis of the *gabrb3*^{-/-} mice could conceivably contribute to a variety of behavioral deficits observed in *gabrb3*^{-/-} mice including poor exploratory performance, reduced rearing, impaired contextual memory, poor motor coordination, hyperactivity, reduced digging, stereotypical circling and hypotonia [27,37,39]. It is noteworthy that the reduced exploratory and presence of stereotypical patterns observed in ASD have both been significantly correlated with the magnitude of cerebellar hypoplasia in vermal lobules VI-VII [64]. Furthermore, lobules VI and VII are part of the oculomotor vermis, which controls saccadic eye and head movements [32] and are therefore essential for adequate exploration of the external world. Although, it is compelling to consider that the behavioral deficits exhibited by *gabrb3*^{-/-} mice are linked with its cerebellar hypoplasia, disrupting other brain regions can also lead to similar behavioral deficits. For example lesioning of the hippocampus in normal mice likewise results in poor nest building and reduced rearing and exploration [25]. Subsequently, *gabrb3*^{-/-} mice also exhibit abnormal GABA_A receptor binding and function in the hippocampus [39,43]. In addition, we recently discovered that the locus coeruleus (LC) in *gabrb3*^{-/-} mice have an expanded plexus of LC dendrites relative to *gabrb3*^{+/+}, C57Bl/6J and 129/SvJ mice [37]. This is of potential interest as the LC has also been implicated in processes involved in focusing, orienting, scanning and shifting attention [7,28,77]. However, while the cerebellum [3] and hippocampus [33] have both been implicated in ASD, to date, the LC has not been highly investigated in regards to ASD.

In summary, the current results clearly indicate that *gabrb3*^{-/-} mice exhibit abnormalities in social behavior. This coupled with an earlier observation that these mice exhibit stereotypical behavior [39], demonstrates these mice to possess two of the triad of core behavioral deficits required in humans in order to receive a diagnosis of ASD. We also report that *gabrb3*^{-/-} mice exhibit deficits in exploratory behavior and in non-selective attention along with hypoplasia of the cerebellar vermis, all features associated with ASD. These current findings are added to a lengthy list of abnormalities previously reported in *gabrb3*^{-/-} mice, which bear obvious parallels to symptoms often reported in association with ASD (Table 4) [26]. Taken together, these observations provide overwhelming support for the face validity of the *gabrb3*^{-/-} mouse as a model of ASD. In addition, strong construct validity of this mouse model is supported by an abundant body of evidence implicating the GABRB3 gene in ASD [18,21,70]. Therefore,

a compelling argument can be made for the strength of the *gabrb3*^{-/-} mouse as a model of ASD and that further investigations into the predictive validity of this model are highly warranted.

Acknowledgments

This research was supported by a grant to Timothy M. DeLorey, Ph.D. from the National Institute of Mental Health, MH065393 and a grant to Gregg E. Homanics, Ph.D. from the National Institute on Alcohol Abuse and Alcoholism, AA10422. We would like to acknowledge Thomas Fjallstam and Carolyn Ferguson for expert technical support.

References

1. Allen G, Buxton RB, Wong EC, Courchesne E. Attentional activation of the cerebellum independent of motor involvement. *Science* 1997;275:1940–3. [PubMed: 9072973]
2. Allen G, Courchesne E. Attention function and dysfunction in autism. *Front Biosci* 2001;6:D105–19. [PubMed: 11171544]
3. Allen G, Muller RA, Courchesne E. Cerebellar function in autism: functional magnetic resonance image activation during a simple motor task. *Biol Psychiatry* 2004;56:269–78. [PubMed: 15312815]
4. Altman J. Morphological and behavioral markers of environmentally induced retardation of brain development: an animal model. *Environ Health Perspect* 1987;74:153–68. [PubMed: 3319550]
5. Aspidre R, Fresiello A, de Filippis G, Carnevale UA, Sadile AG. Non-selective attention in a rat model of hyperactivity and attention deficit: subchronic methylphenydate and nitric oxide synthesis inhibitor treatment. *Neurosci Biobehav Rev* 2000;24:59–71. [PubMed: 10654662]
6. Aspidre R, Gironi Carnevale UA, Sergeant JA, Sadile AG. Non-selective attention and nitric oxide in putative animal models of Attention-Deficit Hyperactivity Disorder. *Behav Brain Res* 1998;95:123–33. [PubMed: 9754884]
7. Aston-Jones G, Rajkowski J, Cohen J. Role of locus coeruleus in attention and behavioral flexibility. *Biol Psychiatry* 1999;46:1309–20. [PubMed: 10560036]
8. Ayres AJ, Tickle LS. Hyper-responsivity to touch and vestibular stimuli as a predictor of positive response to sensory integration procedures by autistic children. *Am J Occup Ther* 1980;34:375–81. [PubMed: 6971048]
9. Barker JL, Behar T, Li YX, Liu QY, Ma W, Maric D, Maric I, Schaffner AE, Serafini R, Smith SV, Somogyi R, Vautrin JY, Wen XL, Xian H. GABAergic cells and signals in CNS development. *Perspect Dev Neurobiol* 1998;5:305–22. [PubMed: 9777645]
10. Barthelemy C, Roux S, Adrien JL, Hameury L, Guerin P, Garreau B, Fermanian J, Lelord G. Validation of the Revised Behavior Summarized Evaluation Scale. *J Autism Dev Disord* 1997;27:139–53. [PubMed: 9105965]
11. Battaglia A. The inv dup(15) or idic(15) syndrome: a clinically recognisable neurogenetic disorder. *Brain Dev* 2005;27:365–9. [PubMed: 16023554]
12. Belzung, C. Measuring rodent exploratory behavior. In: Crusio, WE.; Gerlai, RT., editors. *Handbook of molecular-genetic techniques for brain and behavior research*. Amsterdam: Elsevier; 1999. p. 738–49.
13. Blatt GJ, Fitzgerald CM, Gupitill JT, Booker AB, Kemper TL, Bauman ML. Density and distribution of hippocampal neurotransmitter receptors in autism: an autoradiographic study. *J Autism Dev Disord* 2001;31:537–43. [PubMed: 11814263]
14. Bonnert TP, McKernan RM, Farrar S, le Bourdelles B, Heavens RP, Smith DW, Hewson L, Rigby MR, Sirinathsinghji DJ, Brown N, Wafford KA, Whiting PJ. θ , a novel gamma-aminobutyric acid type A receptor subunit. *Proc Natl Acad Sci U S A* 1999;96:9891–6. [PubMed: 10449790]
15. Bracha HS, Livingston R, Dykman K, Edwards DR, Adam B. An automated electronic method for quantifying spinning (circling) in children with autistic disorder. *J Neuropsychiatry Clin Neurosci* 1995;7:213–7. [PubMed: 7626965]
16. Broida J, Svare B. Mice: progesterone and the regulation of strain differences in pregnancy-induced nest building. *Behav Neurosci* 1983;97:994–1004. [PubMed: 6651968]

17. Burd L, Stenehjem A, Franceschini LA, Kerbeshian J. A 15-year follow-up of a boy with pyridoxine (vitamin B6)-dependent seizures with autism, breath holding, and severe mental retardation. *J Child Neurol* 2000;15:763–5. [PubMed: 11108513]
18. Buxbaum JD, Silverman JM, Smith CJ, Greenberg DA, Kilifarski M, Reichert J, Cook EH Jr, Fang Y, Song CY, Vitale R. Association between a GABRB3 polymorphism and autism. *Mol Psychiatry* 2002;7:311–6. [PubMed: 11920158]
19. Carod FJ, Prats JM, Garaizar C, Zuazo E. Clinical-radiological evaluation of infantile autism and epileptic syndromes associated with autism. *Rev Neurol* 1995;23:1203–7. [PubMed: 8556618]
20. CDC. Prevalence of autism spectrum disorders--autism and developmental disabilities monitoring network, 14 sites, United States, 2002. *MMWR Surveill Summ* 2007;56:12–28. [PubMed: 17287715]
21. Cook EH Jr, Courchesne RY, Cox NJ, Lord C, Gonen D, Guter SJ, Lincoln A, Nix K, Haas R, Leventhal BL, Courchesne E. Linkage-disequilibrium mapping of autistic disorder, with 15q11-13 markers. *Am J Hum Genet* 1998;62:1077–83. [PubMed: 9545402]
22. Cook EH Jr, Lindgren V, Leventhal BL, Courchesne R, Lincoln A, Shulman C, Lord C, Courchesne E. Autism or atypical autism in maternally but not paternally derived proximal 15q duplication. *Am J Hum Genet* 1997;60:928–34. [PubMed: 9106540]
23. Courchesne E, Yeung-Courchesne R, Press GA, Hesselink JR, Jernigan TL. Hypoplasia of cerebellar vermal lobules VI and VII in autism. *N Engl J Med* 1988;318:1349–54. [PubMed: 3367935]
24. Crusio WE. Flanking gene and genetic background problems in genetically manipulated mice. *Biol Psychiatry* 2004;56:381–5. [PubMed: 15364034]
25. Deacon RM, Rawlins JN. Hippocampal lesions, species-typical behaviours and anxiety in mice. *Behav Brain Res* 2005;156:241–9. [PubMed: 15582110]
26. DeLorey, TM. GABRB3 Gene Deficient Mice: A Potential Model of Autism Spectrum Disorder. In: Dhossche, D., editor. *GABA in Autism and Related Disorders*. Vol. 71. New York: Elsevier Science Publishing Co., Inc.; 2005. p. 259-382.
27. DeLorey TM, Handforth A, Anagnostaras SG, Homanics GE, Minassian BA, Asatourian A, Fanselow MS, Delgado-Escueta A, Ellison GD, Olsen RW. Mice lacking the beta3 subunit of the GABAA receptor have the epilepsy phenotype and many of the behavioral characteristics of Angelman syndrome. *J Neurosci* 1998;18:8505–14. [PubMed: 9763493]
28. Devauges V, Sara SJ. Activation of the noradrenergic system facilitates an attentional shift in the rat. *Behav Brain Res* 1990;39:19–28. [PubMed: 2167690]
29. Folstein SE, Rosen-Sheidley B. Genetics of autism: complex aetiology for a heterogeneous disorder. *Nat Rev Genet* 2001;2:943–55. [PubMed: 11733747]
30. Fransen E, D'Hooge R, Van Camp G, Verhoye M, Sijbers J, Reyniers E, Soriano P, Kamiguchi H, Willemsen R, Koekkoek SK, De Zeeuw CI, De Deyn PP, Van der Linden A, Lemmon V, Kooy RF, Willems PJ. L1 knockout mice show dilated ventricles, vermis hypoplasia and impaired exploration patterns. *Hum Mol Genet* 1998;7:999–1009. [PubMed: 9580664]
31. Gillberg C, Nordin V, Ehlers S. Early detection of autism. Diagnostic instruments for clinicians. *Eur Child Adolesc Psychiatry* 1996;5:67–74. [PubMed: 8814412]
32. Godschalk M, Van der Burg J, Van Duin B, De Zeeuw CI. Topography of saccadic eye movements evoked by microstimulation in rabbit cerebellar vermis. *J Physiol* 1994;480(Pt 1):147–53. [PubMed: 7853218]
33. Guptill JT, Booker AB, Gibbs TT, Kemper TL, Bauman ML, Blatt GJ. [3H]-flunitrazepam-labeled benzodiazepine binding sites in the hippocampal formation in autism: a multiple concentration autoradiographic study. *J Autism Dev Disord* 2007;37:911–20. [PubMed: 17019626]
34. Hara H. Autism and epilepsy: A retrospective follow-up study. *Brain Dev* 2007;29:486–90. [PubMed: 17321709]
35. Harris NS, Courchesne E, Townsend J, Carper RA, Lord C. Neuroanatomic contributions to slowed orienting of attention in children with autism. *Brain Res Cogn Brain Res* 1999;8:61–71. [PubMed: 10216274]
36. Harrison J, Hare DJ. Brief report: assessment of sensory abnormalities in people with autistic spectrum disorders. *J Autism Dev Disord* 2004;34:727–30. [PubMed: 15679192]

37. Hashemi E, Sahbaie P, Davies MF, Clark JD, DeLorey TM. Gabrb3 gene deficient mice exhibit increased risk assessment behavior, hypotonia and expansion of the plexus of locus coeruleus dendrites. *Brain Res* 2007;1129:191–9. [PubMed: 17156762]
38. Hogart A, Nagarajan RP, Patzel KA, Yasui DH, Lasalle JM. 15q11-13 GABAA receptor genes are normally biallelically expressed in brain yet are subject to epigenetic dysregulation in autism-spectrum disorders. *Hum Mol Genet* 2007;16:691–703. [PubMed: 17339270]
39. Homanics GE, DeLorey TM, Firestone LL, Quinlan JJ, Handforth A, Harrison NL, Krasowski MD, Rick CE, Korpi ER, Makela R, Brilliant MH, Hagiwara N, Ferguson C, Snyder K, Olsen RW. Mice devoid of gamma-aminobutyrate type A receptor beta3 subunit have epilepsy, cleft palate, and hypersensitive behavior. *Proc Natl Acad Sci U S A* 1997;94:4143–8. [PubMed: 9108119]
40. Isumi H, Mizuguchi M, Takashima S. Differential development of the human cerebellar vermis: immunohistochemical and morphometrical evaluation. *Brain Dev* 1997;19:254–7. [PubMed: 9187474]
41. Joyal CC, Meyer C, Jacquart G, Mahler P, Caston J, Lalonde R. Effects of midline and lateral cerebellar lesions on motor coordination and spatial orientation. *Brain Res* 1996;739:1–11. [PubMed: 8955918]
42. Koo JW, Homanics GE, Balaban CD. Hypoplasia of spiral and Scarpa's ganglion cells in GABA(A) receptor beta(3) subunit knockout mice. *Hear Res* 2002;167:71–80. [PubMed: 12117532]
43. Krasowski MD, Rick CE, Harrison NL, Firestone LL, Homanics GE. A deficit of functional GABA (A) receptors in neurons of beta 3 subunit knockout mice. *Neurosci Lett* 1998;240:81–4. [PubMed: 9486477]
44. Lalonde R, Botez MI. The cerebellum and learning processes in animals. *Brain Res Brain Res Rev* 1990;15:325–32. [PubMed: 2289089]
45. Leiner HC, Leiner AL, Dow RS. Cognitive and language functions of the human cerebellum. *Trends Neurosci* 1993;16:444–7. [PubMed: 7507614]
46. Lijam N, Paylor R, McDonald MP, Crawley JN, Deng CX, Herrup K, Stevens KE, Maccaferri G, McBain CJ, Sussman DJ, Wynshaw-Boris A. Social interaction and sensorimotor gating abnormalities in mice lacking Dvl1. *Cell* 1997;90:895–905. [PubMed: 9298901]
47. Limoges E, Mottron L, Bolduc C, Berthiaume C, Godbout R. Atypical sleep architecture and the autism phenotype. *Brain*. 2005
48. Long JM, LaPorte P, Paylor R, Wynshaw-Boris A. Expanded characterization of the social interaction abnormalities in mice lacking Dvl1. *Genes Brain Behav* 2004;3:51–62. [PubMed: 14960015]
49. Maison SF, Rosahl TW, Homanics GE, Liberman MC. Functional role of GABAergic innervation of the cochlea: phenotypic analysis of mice lacking GABA(A) receptor subunits alpha 1, alpha 2, alpha 5, alpha 6, beta 2, beta 3, or delta. *J Neurosci* 2006;26:10315–26. [PubMed: 17021187]
50. McCauley JL, Olson LM, Delahanty R, Amin T, Nurmi EL, Organ EL, Jacobs MM, Folstein SE, Haines JL, Sutcliffe JS. A linkage disequilibrium map of the 1-Mb 15q12 GABA(A) receptor subunit cluster and association to autism. *Am J Med Genet B Neuropsychiatr Genet* 2004;131:51–9. [PubMed: 15389768]
51. Menold MM, Shao Y, Wolpert CM, Donnelly SL, Raiford KL, Martin ER, Ravan SA, Abramson RK, Wright HH, DeLong GR, Cuccaro ML, Pericak-Vance MA, Gilbert JR. Association analysis of chromosome 15 gabaa receptor subunit genes in autistic disorder. *J Neurogenet* 2001;15:245–59. [PubMed: 12092907]
52. Moretti P, Bouwknecht JA, Teague R, Paylor R, Zoghbi HY. Abnormalities of social interactions and home-cage behavior in a mouse model of Rett syndrome. *Hum Mol Genet* 2005;14:205–20. [PubMed: 15548546]
53. Moy SS, Nadler JJ, Perez A, Barbaro RP, Johns JM, Magnuson TR, Piven J, Crawley JN. Sociability and preference for social novelty in five inbred strains: an approach to assess autistic-like behavior in mice. *Genes Brain Behav* 2004;3:287–302. [PubMed: 15344922]
54. Nadler JJ, Moy SS, Dold G, Trang D, Simmons N, Perez A, Young NB, Barbaro RP, Piven J, Magnuson TR, Crawley JN. Automated apparatus for quantitation of social approach behaviors in mice. *Genes Brain Behav* 2004;3:303–14. [PubMed: 15344923]
55. Noens I, van Berckelaer-Onnes I. Making sense in a fragmentary world: communication in people with autism and learning disability. *Autism* 2004;8:197–218. [PubMed: 15165435]

56. Nurmi EL, Amin T, Olson LM, Jacobs MM, McCauley JL, Lam AY, Organ EL, Folstein SE, Haines JL, Sutcliffe JS. Dense linkage disequilibrium mapping in the 15q11-q13 maternal expression domain yields evidence for association in autism. *Mol Psychiatry* 2003;8:624–34, 570. [PubMed: 12851639]
57. Nusser Z, Kay LM, Laurent G, Homanics GE, Mody I. Disruption of GABA(A) receptors on GABAergic interneurons leads to increased oscillatory power in the olfactory bulb network. *J Neurophysiol* 2001;86:2823–33. [PubMed: 11731539]
58. Olsen, RW.; DeLorey, TM. GABA and Glycine. In: Siegel, GJ.; Agranoff, BW.; Albers, RW.; Fisher, SK.; Uhler, MD., editors. *Basic Neurochemistry*. New York: Lippincott-Raven; 1999. p. 335-346.
59. Paxinos, G.; Franklin, K. *The Mouse Brain: In stereotaxic coordinates*. San Diego: Academic Press; 2001. p. 350
60. Persico AM, Bourgeron T. Searching for ways out of the autism maze: genetic, epigenetic and environmental clues. *Trends Neurosci* 2006;29:349–58. [PubMed: 16808981]
61. Peters SU, Beaudet AL, Madduri N, Bacino CA. Autism in Angelman syndrome: implications for autism research. *Clin Genet* 2004;66:530–6. [PubMed: 15521981]
62. Petrosini L, Graziano A, Mandolesi L, Neri P, Molinari M, Leggio MG. Watch how to do it! New advances in learning by observation. *Brain Res Brain Res Rev* 2003;42:252–64. [PubMed: 12791443]
63. Petrosini L, Molinari M, Dell'Anna ME. Cerebellar contribution to spatial event processing: Morris water maze and T-maze. *Eur J Neurosci* 1996;8:1882–96. [PubMed: 8921279]
64. Pierce K, Courchesne E. Evidence for a cerebellar role in reduced exploration and stereotyped behavior in autism. *Biol Psychiatry* 2001;49:655–64. [PubMed: 11313033]
65. Rineer S, Finucane B, Simon EW. Autistic symptoms among children and young adults with isodicentric chromosome 15. *Am J Med Genet* 1998;81:428–33. [PubMed: 9754629]
66. Sacchetti B, Baldi E, Lorenzini CA, Bucherelli C. Cerebellar role in fear-conditioning consolidation. *Proc Natl Acad Sci U S A* 2002;99:8406–11. [PubMed: 12034877]
67. Sadile, AG. Long-term habituation of theta-related activity components of albino rats in the Lat-maze. In: Sanberg, P.; Ossenkopp, K.; Kavaliers, M., editors. *Motor Activity and Movement Disorders: Measurement and Analysis*. New York: Humana; 1995. p. 1-54.
68. Santangelo SL, Tsatsanis K. What is known about autism: genes, brain, and behavior. *Am J Pharmacogenomics* 2005;5:71–92. [PubMed: 15813671]
69. Schroer RJ, Phelan MC, Michaelis RC, Crawford EC, Skinner SA, Cuccaro M, Simensen RJ, Bishop J, Skinner C, Fender D, Stevenson RE. Autism and maternally derived aberrations of chromosome 15q. *Am J Med Genet* 1998;76:327–36. [PubMed: 9545097]
70. Shao Y, Cuccaro ML, Hauser ER, Raiford KL, Menold MM, Wolpert CM, Ravan SA, Elston L, Decena K, Donnelly SL, Abramson RK, Wright HH, DeLong GR, Gilbert JR, Pericak-Vance MA. Fine mapping of autistic disorder to chromosome 15q11-q13 by use of phenotypic subtypes. *Am J Hum Genet* 2003;72:539–48. [PubMed: 12567325]
71. Sinkkonen ST, Homanics GE, Korpi ER. Mouse models of Angelman syndrome, a neurodevelopmental disorder, display different brain regional GABA(A) receptor alterations. *Neurosci Lett* 2003;340:205–8. [PubMed: 12672542]
72. Sluyter F, Marican CC, Crusio WE. Further phenotypical characterisation of two substrains of C57BL/6J inbred mice differing by a spontaneous single-gene mutation. *Behav Brain Res* 1999;98:39–43. [PubMed: 10210520]
73. Steffenburg S, Gillberg CL, Steffenburg U, Kyllerman M. Autism in Angelman syndrome: a population-based study. *Pediatr Neurol* 1996;14:131–6. [PubMed: 8703225]
74. Teitelbaum P, Teitelbaum O, Nye J, Fryman J, Maurer RG. Movement analysis in infancy may be useful for early diagnosis of autism. *Proc Natl Acad Sci U S A* 1998;95:13982–7. [PubMed: 9811912]
75. Townsend J, Courchesne E, Covington J, Westerfield M, Harris NS, Lyden P, Lowry TP, Press GA. Spatial attention deficits in patients with acquired or developmental cerebellar abnormality. *J Neurosci* 1999;19:5632–43. [PubMed: 10377369]
76. Ugarte SD, Homanics GE, Firestone LL, Hammond DL. Sensory thresholds and the antinociceptive effects of GABA receptor agonists in mice lacking the beta3 subunit of the GABA(A) receptor. *Neuroscience* 2000;95:795–806. [PubMed: 10670447]

77. Van Bockstaele EJ. Morphological substrates underlying opioid, epinephrine and gamma-aminobutyric acid inhibitory actions in the rat locus coeruleus. *Brain Res Bull* 1998;47:1–15. [PubMed: 9766384]
78. Vig S, Jedrysek E. Autistic features in young children with significant cognitive impairment: autism or mental retardation? *J Autism Dev Disord* 1999;29:235–48. [PubMed: 10425586]
79. Wisor JP, DeLorey TM, Homanics GE, Edgar DM. Sleep states and sleep electroencephalographic spectral power in mice lacking the beta 3 subunit of the GABA(A) receptor. *Brain Res* 2002;955:221–8. [PubMed: 12419540]

Figure 1.

Diagram of the automated apparatus used in assessing social behavior in mice. Side view showing location of infra-red photo-sensors and chamber partitions. Partitions can be automatically opened to create a 5 cm wide opening, allowing the test mouse to pass freely from one chamber to the next. A wire mesh wall separates each of the interaction chambers from an adjacent stimulus cage, which will either be empty or contain a “stranger mouse”.

Figure 2. Sociability histograms of the time that mice spent in each of the 3 chambers of the social interaction apparatus. The black bars represent the time spent in interaction chamber #1, adjacent to the stimulus cage containing an unfamiliar mouse (stranger mouse #1), grey bars represent time in the neutral central chamber and white is time in interaction chamber #2, adjacent to an empty stimulus cage. Data presented as the mean $\pm$ SEM. C57BL/6J n=6, 129/SvJ n=6, gabrb3^{+/+} n=6, gabrb3^{-/-} n=6. * p <0.05, ** p <0.01.

Figure 3.

Social novelty histograms of the mean time (±SEM) mice spent in the interaction chamber adjacent to the now familiar stranger mouse #1 (black bar), the unfamiliar stranger mouse #2 (white bar) or in the neutral central chamber (grey bar). As expected all three control genotypes exhibited a significant preference for the unfamiliar mouse (stranger mouse #2) over the now familiar mouse (stranger mouse #1). The preference for stranger mouse #2 remained significant into the second five min epoch for both C57BL/6J and 129/SvJ mice. C57BL/6J n=6, 129/SvJ n=6, *gabrb3*^{+/+} n=6, *gabrb3*^{-/-} n=6. ***p*<0.01, ****p*<0.001.

Figure 4.

Histograms of the sociability and the preference for social novelty assessed from videotape, by an observer, blind to the genotype of the test mice. Observer scores in the sociability test are presented as the total time, during a 10 min test period, that test mice spent with their noses within 2 cm of the wire mesh of the stimulus cage containing an unfamiliar mouse (stranger mouse #1) versus the total time the test mice spent with their noses within 2 cm of the wire mesh of the stimulus cage that was empty. In the preference for social novelty test the data is presented as the amount of time mice were observed to have their noses within 2 cm of the stimulus cage containing stranger mouse #1 (the now familiar mouse) versus the stimulus cage containing stranger mouse #2 (the new unfamiliar mouse). * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$.

Figure 5. Histogram of the percentage of the total 10 min test period that each mouse genotype spent sniffing at the stimulus cage containing **A.)** Stranger mouse #1 in the sociability test. **B.)** Stranger mouse #2 in the social novelty test and **C.)** % of total time spent sniffing at both Stranger mouse #1 and Stranger mouse #2 in the social novelty test. * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$

Figure 6.

A.) Histogram of the mean time ($\pm$ SEM) that each mouse genotypes spent interacting with freshly introduced nesting material (nestlet) during three separate 10 min observations taken over a 1 hr period immediately after the introduction of the nestlet. **B.)** Histogram of average nest scores (mean $\pm$ SEM), taken 48 hr after nestlet introduction. * p <0.05, ** p <0.01, *** p <0.001. *Gabrb3*^{-/-} n =6, *gabrb3*^{+/+} n =8, C57BL/6J n =8, 129/SvJ n =8.

Figure 7.

Representative examples of the exploratory paths taken by a **gabrb3^{+/+}** and **gabrb3^{-/-}** mouse in an open field chamber that contained a novel object (an empty round cage, depicted as a solid black disc) placed in the center, that was not present during the acclimation phase of the test. The arrow indicates an example of the stereotypical circling often exhibited by **gabrb3^{-/-}** mice.

Figure 8.

A.) Average rearing frequency (mean $\pm$ S.E.M.) demonstrated by each genotype, when placed in an open field chamber containing a novel object. *Post hoc* analysis revealed a significant difference ($p < 0.001$) between gabrb3^{-/-} mice and each control genotype in rearing frequency over the entire 10 min test period. **B.)** Average duration of a rearing event for each mouse genotype, averaged over the entire 10 min testing period in a novel environment. Gabrb3^{-/-} n=7, gabrb3^{+/+} n=7, C57BL/6J n=8, 129/SvJ n=7. * $p < 0.05$, *** $p < 0.001$

Figure 9.

Representative examples of toluidine blue stained midsagittal sections through the cerebellar vermis (0.36-0.60 mm lateral from the midsagittal line) from *gabrb3*^{+/+} and *gabrb3*^{-/-} mice. A roman numeral identifies each cerebellar vermal lobule. Lobules VI-VII on the *gabrb3*^{+/+} brain section have been outlined using a white dashed line to indicate the area being measured and compared.

Figure 10.

Histogram of the measured surface area of cerebellar vermal lobules in *gabrb3*^{+/+} (n=4) and *gabrb3*^{-/-} (n=5) mice. **p*<0.05, ***p*<0.01.

Table 1

Baseline locomotor activity

Mouse genotype	Ambulatory activity [†]	Total transitions between chambers	Chamber entrances		Time spent in each chamber (sec)	
			Left	Right	Left	Right
gabrb3 ^{-/-}	1936 ± 129	186 ± 40	49 ± 8	57 ± 19	208 ± 30	213 ± 33
gabrb3 ^{+/+}	1278 ± 92 ^{***}	132 ± 14 [*]	38 ± 3	31 ± 7	245 ± 29	173 ± 26
C57BL/6J	1459 ± 45 ^{**}	106 ± 15 ^{**}	29 ± 4	26 ± 6	229 ± 33	159 ± 28
129/SvJ	1342 ± 100 ^{***}	84 ± 14 ^{***}	23 ± 4	20 ± 4	237 ± 39	309 ± 37

Baseline locomotor data collected on test mice during the acclimation phase of the social behavior assessment, prior to introducing a “stranger mouse” to the test apparatus. Data was collected over a 10 min period. Data presented as the mean ± SEM. Gabrb3^{-/-} n=6, gabrb3^{+/+} n=6, C57BL/6J n=6, 129/SvJ n=6.

* $p < 0.05$,
** $p < 0.01$,
*** $p < 0.001$.
[†] arbitrary units

Table 2**Material utilization in nest construction**

Genotype	1 hr	24 hr	48 hr
gabrb3 ^{-/-}	0.3 ± 0.3	1.6 ± 0.4	4.1 ± 0.9
gabrb3 ^{+/-}	1.9 ± 0.4 ***	4.6 ± 0.2 ***	7.0 ± 0.3 ***
C57BL/6J	2.0 ± 0.2 ***	4.9 ± 0.0 ***	7.4 ± 0.1 ***
129/SvJ	2.0 ± 0.2 ***	5.0 ± 0.0 ***	7.5 ± 0.0 ***

Data is presented as the mean ± SEM of material usage, in grams, measured at three time intervals following initial introduction of nestlet material. Bonferroni *post hoc* comparison of material usage between each control mouse genotype and gabrb3^{-/-} mice at each time interval. Gabrb3^{-/-} n=6, gabrb3^{+/-} n=8, C57BL/6J n=8, 129/SvJ n=8.

p<0.001.

Table 3**Exploratory behavior**

Mouse genotype	Time to first contact with novel object (sec)	Time spent investigating novel object (sec)	Number of contacts with novel object	Total number of rearings
gabrb3 ^{-/-}	54.9 ± 12.0	63.6 ± 10.5	15.9 ± 4.5	14.2 ± 6.6
gabrb3 ^{+/+}	12.1 ± 1.7 ^{***}	236.1 ± 24.6 ^{***}	20.4 ± 1.4	51.6 ± 7.9 ^{**}
C57BL/6J	18.9 ± 3.9 ^{**}	180.8 ± 8.8 ^{**}	21.8 ± 0.7	53.0 ± 6.4 ^{***}
129/SvJ	15.0 ± 5.2 ^{***}	426.4 ± 39.7 ^{***}	46.2 ± 4.2 ^{***}	30.8 ± 3.1

Data was quantified from 10 min videotape sessions of mice placed in an open-field chamber containing a novel object, scored by an observer blind to the mouse genotype. Data represents the mean ± SEM. Gabrb3^{+/+} n=7, gabrb3^{-/-} n=7, C57BL/6J n=8, 129/SvJ n= 8.

**
p<0.01,

p<0.001.

Table 4
Overlap between ASD characteristics and phenotypic traits of *gabrb3*^{-/-} mice

Characteristic	ASD	<i>gabrb3</i> ^{-/-} mice	Reference
<i>Core deficits</i>			
Impaired social behavior	✓	✓ [*]	[31]
Repetitive, stereotypical behavior	✓	✓	[10,39]
Deficits in communication	✓	?	[55]
<i>Inclinator deficits</i>			
Hyperactivity	✓	✓	[10,27]
Poor motor coordination	✓	✓	[10,27]
Tactile hypersensitivity	✓	✓	[8,76]
Temperature hypersensitivity	✓	✓	[36,76]
Cognitive impairment	✓	✓	[27,78]
Sleep disturbances	✓	✓	[10,79]
Epilepsy	✓	✓	[19,27]
Abnormal exploratory behavior	✓	✓ [*]	[64]
Deficits in orientation of attention	✓	✓ [*]	[35]
Reduced cerebellar vermis size	✓	✓ [*]	[23]
Reduced benzodiazepine binding in hippocampus	✓	✓	[13,71]

✓
 = feature is present.

*
 Reported in the current study.