Published in final edited form as:

Pharmacol Biochem Behav. 2009 May ; 92(3): 480–487. doi:10.1016/j.pbb.2009.01.022.

Blocking GABA-A receptors in the Medial Septum Enhances Hippocampal Acetylcholine Release and Behavior in a Rat Model of Diencephalic Amnesia

Jessica J. Roland* and Lisa M. Savage

Behavioral Neuroscience Program, Department of Psychology, Binghamton University-State University of New York

Abstract

Wernicke-Korsakoff syndrome (WKS), a form of diencephalic amnesia caused by thiamine deficiency, results in severe anterograde memory loss. Pyridoxamine-induced thiamine deficiency (PTD), an animal model of WKS, produces cholinergic abnormalities including decreased functional hippocampal acetylcholine (ACh) release and poor spatial memory. Increasing hippocampal ACh levels has increased performance in PTD animals. Intraseptal bicuculline (GABA_A antagonist) augments hippocampal ACh release in normal animals and we found it (0.50µg/µl & 0.75µg/µl) also increased in-vivo hippocampal ACh release in PTD animals. However, the 0.75 µg/µl dose produced a greater change in hippocampal ACh release in control animals. The 0.50µg/µl dose of bicuculline was then selected to determine if it could enhance spontaneous alternation performance in PTD animals. This dose of bicuculline significantly increased hippocampal ACh levels above baseline in both PTD and control rats and resulted in complete behavioral recovery in PTD animals, without altering performance in control rats. This suggests that balancing ACh-GABA interactions in the septohippocampal circuit may be an effective therapeutic approach in certain amnesic syndromes.

Keywords

septohippocampal pathway; bicuculline; microdialysis; spontaneous alternation

1. Introduction

Wernicke-Korsakoff Syndrome (WKS), most often diagnosed in patients with chronic alcoholism, also is seen in disorders in which malnourishment – specifically thiamine deficiency-- is a component (AIDS, chronic feeding tube, cancer, etc.). WKS has an acute and chronic phase that often culminates in severe explicit memory deficits (Harper and Kril, 1986; Kopelman, 1995; Squire, 1981). The animal model of this disorder, known as pyridoxamine-induced thiamine deficiency (PTD), produces damage in both the mammillary bodies and several thalamic nuclei (anterior, medial dorsal, intralaminar, internal medullary lamina) that are interconnected with the limbic system (Langlais and Savage, 1995; Langlais et al., 1996; Mair, 1994; Victor et al., 1971).

*Corresponding Author, New Jersey Veterans Hospital, 385 Tremont Ave, East Orange, NJ 07014, Phone: (973) 676-1000 x2558, FAX: (973) 395-7114, E-Mail: E-mail: jessica.roland80@gmail.com.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Beyond damage to diencephalic structures, a number of studies have demonstrated dysfunction in the septohippocampal pathway. Both WKS patients (Reed et al., 2003; Sullivan and Marsh, 2003) and PTD-treated animals (Langlais and Zhang, 1997) display extensive degeneration in both the mammillothalamic tract and the fornix which interconnect with the septum and hippocampus (Aggleton and Brown, 1999). In addition, PTD-treated rats display a 25-30% loss of choline acetyltransferase (ChAT)-positive neurons in the medial septum/ diagonal band (MS/DB) (Pitkin and Savage, 2004, 2001; Savage et al., 2007). Furthermore, our laboratory has demonstrated that PTD animals have a selective decrease in hippocampal ACh efflux that parallels their poor performance (Roland and Savage, 2007). Similar results have been found when cortical ACh release was stimulated using potassium: Rats feed on a thiamine deficient (TD) diet alone had significantly less hippocampal and frontal cortex ACh release, compared to saline injected controls (Pires et al., 2005). Together these results demonstrate a septohippocampal cholinergic dysfunction that occurs after thiamine deficiency. Drugs that enhance cholinergic activity, such as acetylcholinesterase (AChE) inhibitors have been shown to improve the cognitive functioning in WKS and animal models of the disorder (Nakagawasai et al., 2000b)-- as well as increase hippocampal ACh release (Roland et al., 2008). It has been suggested that AChE inhibitors, such as physostigmine, are having their cognitive enhancing effects in the medial septal region of the basal forebrain (Mulder et al., 2005; Wu et al., 2003b).

However, the medial septum contains not only cholinergic, but also GABAergic and glutamatergic neurons that have local and diffuse projections (Manseau et al., 2005; Wainer et al., 1985). The interactions of the cholinergic and GABAergic populations in the MS/DB are important for modulating hippocampal-dependent learning and memory performance. The MS/DB cholinergic and GABAergic neurons contain both muscarinic and GABA_A receptors (Gao et al., 1995; Van der Zee and Luiten, 1994), therefore medial septal infusions of cholinergic and GABAergic agonists/antagonists will have an effect on both receptor systems. In turn, hippocampal activation is modulated by both of these MS/DB neurotransmitter systems. The presence of a muscarinic agonist in the medial septal region decreases the activity of the hippocampal GABAergic interneurons *in vitro* (Alreja et al., 2000b); increasing overall activity of the hippocampal pyramidal neurons through disinhibitory mechanisms (Toth et al., 1997).

The role of GABA_A receptor activation in the MS on the modulation of hippocampal activity and behavior has began to be elucidated. Medial septal infusions of a GABA_A antagonist increase hippocampal ACh release, whereas a GABA_A agonist decreases hippocampal ACh release (Moor et al., 1998a; Moor et al., 1998b). Furthermore, there is some evidence that the behavioral effects of activation of MS GABA_A receptors are further modulated by the action of GABA in the hippocampus. When animals were given septal muscimol (GABA_A agonist) combined with hippocampal bicuculline (GABA_A antagonist), the memory impairing effects normally produced by septal muscimol were attenuated (Krebs-Kraft et al., 2007). However, this study does not rule out the possibility that MS GABA_A receptor activation also influences cholinergic projection neurons.

The MS/DB cholinergic neurons supply an excitatory drive to the MS/DB GABAergic neurons (Alreja et al., 2000b); therefore a loss of cholinergic neurons, as occurs in PTD-treatment, could reduce activity in both the cholinergic and GABAergic septohippocampal pathways. Within the PTD model, studies have focused primarily on the cholinergic system; however, long-term changes in the GABAergic system have been reported. Reductions in midline-thalamic GABA levels have been seen up to 9 weeks after recovery from PTD-treatment (Langlais et al., 1988). However, the septohippocampal GABAergic system has not been examined for neuronal damage after PTD-treatment. Given the role of septohippocampal GABA in cognitive processes and the interaction between septohippocampal ACh and GABA, a change in

GABAergic functioning following recovery from PTD treatment could be contributing to the learning and memory impairments associated with this model.

Therefore, in Experiment 1 a dose-response curve was conducted using two different doses (0.5µg/µl and 0.75µg/µl) of the GABA_A antagonist bicuculline that have been shown to increase hippocampal ACh levels in normal animals (Chang et al., 2006). Increasing functional hippocampal ACh levels in PTD animals has been shown to increase spatial memory performance (Roland et al., 2008). Previous work has demonstrated that either i.c.v. or intrahippocampal bicuculline administration increases memory retention on a passive avoidance task (Zarrindast et al., 2002; Zarrindast et al., 2004). If bicuculline increased hippocampal ACh levels, it was expected that it would also increase behavioral performance in PTD-treated animals. In Experiment 2, bicuculline was infused into the medial septum while in-vivo microdialysis was performed for ACh in the ventral hippocampus and animals were tested on spontaneous alternation. Our main aim of these experiments was to determine if antagonizing septal GABA-A receptors to enhance hippocampal ACh efflux would recover spatial performance in PTD rats.

2. Experimental Procedures

2.1. Subjects

A total of 32 adult (3-4 months old) male Sprague-Dawley rats were used for these experiments (12 for Exp. 1 and 20 for Exp. 2; Harlan, Indianapolis, IN). All animals were maintained throughout the experiments in a 12hr light/dark cycle and housed in standard laboratory Plexiglas cages (dimensions: 30 × 45 × 18 cm). Animals were pair housed until surgery, after which they were single housed. All behavioral and microdialysis testing was performed during the light cycle. All procedures involving rats were approved by the Binghamton University Institutional Animal Care and Use Committee (IACUC).

2.2. PTD Treatment

At the onset of the treatment phase, both PTD (Exp 1: n = 6; Exp 2: n = 10) and pair-fed (PF) animals (Exp 1: n = 6; Exp 2: n = 10) had ad libitum access to water and thiamine-deficient chow (Harlan-Teklad Mills, WI). PTD treatment consisted of daily injections of pyrithiamine HBr (0.25 mg/kg, i.p. Sigma Chem. Corp., MO). Pyrithiamine is a thiamine antagonist that inhibits thiamine pyrophosphokinase, the enzyme that catalyzes the production of the coenzyme form of thiamine (Butterworth and Heroux, 1989). Within 14 -16 days of treatment, PTD subjects developed symptoms of anorexia, ataxia, loss of righting reflexes, and eventually display seizure activity. Animals were monitored bi-hourly for these neurological changes starting on day 13. Within 4 – 4.5 hours of the appearance of seizure activity, all PTD-treated rats were reversed with a large dose of thiamine (100mg/kg). The described protocol produces reliable diencephalic damage (Langlais and Savage, 1995). The administration of thiamine and return to normal chow completely reversed the acute symptoms within 8 hours and animals typically were fully recovered within 24 hours. The PF control rats were fed daily with thiamine deficient chow equal to the average amount eaten by the PTD-treatment group (to mimic the anorexia effects) and were given daily injections of thiamine HCl (.04mg/kg i.p.). All rats were given two weeks to recover and return to a normal free-feeding weight. Weight was monitored daily and any animals that did not show signs of recovery, such as failure to gain weight, were euthanized with a solution of sodium pentobarbital and isopropyl solution (Sleepaway, Fort Dodge Laboratories, Fort Dodge IA, 1mg/kg, i.p.).

2.3. Stereotaxic surgery

Three weeks after PTD/PF treatment and one week prior to microdialysis, all subjects were anesthetized with a Ketamine (8.25ml)/ Xylazine (1.75ml) mixture (50mg/kg i.p.) in

preparation for stereotaxic surgery. All animals were implanted with a microdialysis guide cannula (CMA/11) in the hippocampus and a drug cannula that is 5.0mm in length (22 gauge; Plastics One, Roanoke, VA) in the medial septum. Within each group, subjects were matched in terms of hippocampal microdialysis cannula placement (right vs. left hemisphere). Stereotaxic coordinates were obtained from the atlas of Paxinos and Watson (1986) and are based on previous studies (see Chang and Gold, 2003a). The coordinates for the hippocampus microdialysis cannula were: Anterior-Posterior (AP): -5.1mm; Medial-Lateral (ML): 5.0mm; Dorsal-Ventral (DV): 4.2mm. The coordinates for the medial septum drug cannula were: AP: +0.30mm; ML: 0; DV: 5.0mm. Acrylic cement and 2 skull screws were used to hold the cannulae in place. Immediately after surgery, subjects were placed in a warm incubator until they regained an upright posture. All subjects received a subcutaneous injection of the analgesic buprenorphine (.1ml/ 100g) directly after surgery and another injection 24 hours later. All subjects were allowed to recover for 1 week with ad libitum access to food and water.

2.4. High performance liquid chromatography (HPLC)

Acetylcholine output was assayed by HPLC (Epison, BAS, West Lafayette, IN) along with an enzyme reactor. The assay system includes an ion-exchange microbore analytical column (BAS, MR-8904), a microbore ACh/choline immobilized enzyme reactor containing acetylcholinesterase and choline oxidase (BAS MF-8903), an auxiliary electrode with a radical flow electrochemical thin-layer cell and 13 mm thin layer gasket, a wired enzyme electrode kit (a redox polymer film containing horseradish peroxidase coated in the surface of a 3 mm glassy carbon working electrode), and a low dispersion injected value with a 10 μ l polyetherethketone loop. The mobile phase is a 50 mM dibasic potassium phosphate buffer (pH = 8.5) containing Kathon (BAS). The mobile phase was delivered at a rate of 140 μ l/min by a PM-91 pump (BAS). The detection level was about 10 femtomoles. ACh standards (5 μ l of 20 and 100 nM ACh + Ch) were injected before and after samples to verify detection stability.

2.5. Histology

After the completion of testing, all animals were deeply anesthetized with Sleep Away (1mg/kg, i.p.) and sacrificed. The brains were removed and placed in a 10% formalin solution for 1 week followed by emersion in a 30% sucrose solution for an additional week. Brains were then be frozen and sliced (60 μ m) with a sliding microtome from the anterior commissure to the posterior pontine tegmentum. Sections were evaluated for diencephalic damage and cannula location using cresyl violet staining. All animals in both Experiments 1 and 2 had accurate medial septal and hippocampal cannula placement, therefore all animals were included in final data analysis.

2.6. Experimental Design

2.6.1. Experiment 1. The effects of intraseptal bicuculline administration on hippocampal acetylcholine release in PTD and PF animals—Experiment 1 used a total of 12 animals (PF = 6; PTD = 6); all rats survived the PTD/PF-treatment and were given three weeks to recover and return to a normal free-feeding weight. Following recovery from PTD treatment and cannula surgery, all animals received saline preceding the two doses of drug in a single in-vivo microdialysis session. All animals received both drug doses with a 2-hour window between active dose deliveries.

In-vivo Microdialysis & Pharmacological Challenge: Animals received intraseptal infusions of saline (1 μ l) and bicuculline (0.50 μ g/ μ l and 0.75 μ g/ μ l) during a single session of in vivo microdialysis. Within each group of 6 rats (PF and PTD), after an infusion of saline, 3 rats received the low dose (0.5 μ g/ μ l) of bicuculline first and the other 3 rats received the high dose (0.75 μ g/ μ l) of bicuculline first. This was done to control for an effect of drug dose.

Bicuculline doses were based on a study showing that in normal animals, similar doses of bicuculline in the septum increased hippocampal ACh levels without any adverse side effects (Chang et al., 2006). During the single microdialysis session, samples from the posterior ventral hippocampus were collected and later analyzed for ACh level. Subjects were transported to the testing room and placed in a microdialysis holding cage [acrylic cage (30 cm × 40 cm, depth 35 cm) with wood shavings at the bottom]. The microdialysis probe (CMA/11, 3mm) was connected to a microinfusion system and perfused continuously at a rate of 2 µl/min with artificial cerebrospinal fluid (aCSF; 127.6 mM NaCl, 4 mM KCl, 1.3 mM CaCl₂ dihydrate, 1.0 mM glucose, 0.9 mM MgCl₂, 0.9 mM NaH₂PO₄, and 2 mM Na₂HPO₄) plus neostigmine bromide (500nM) brought to pH 7.0. Following an initial period of habituation to the cage (60 minutes), dialysate samples (volume ~30 µl) were collected every 15 minutes for a period of 30 minutes to determine basal levels of ACh in awake rats. Changes in ACh levels caused by the drug were later determined as a change from baseline samples. After 30 minutes of baseline collection, an infusion needle (Plastics One, 15mm) was inserted into the septal drug cannula and 1 µl of saline was infused over 1 minute. The infusion needle extended 1mm beyond the drug cannula and was always left in place for an additional 2 minutes to allow for diffusion of drug into the site. After the saline infusion, samples continued to be collected for an additional 30 minutes (total of 2 samples). After which 1 µl of bicuculline (either 0.5µg or 0.75µg) was infused into the medial septum. The first drug phase lasted a total of 90 minutes (6-samples). Ninety minutes after the first dose of bicuculline, 1 µl of the second dose of bicuculline was infused. The second drug phase was also 90 minutes. Following the second drug phase, 2 – 15 minutes post-baseline samples were collected. Half the animals within each group (PF and PTD) received the low dose of bicuculline first while the other half received the high dose of bicuculline first. To test the reliability of the microdialysis probe, it was placed into a standard solution (100 nM concentration of ACh and choline) to assess recovery rate (RR). The minimal amount of ACh that can be detected in the system was 10 femtomoles. After collection, brain dialysis samples were frozen for later analysis using HPLC.

2.6.2. Experiment 2. The effects of intraseptal bicuculline administration on spontaneous alternation behavior in PF and PTD-treated rats—Experiment 2 used a total of 20 animals (PF = 10; PTD = 10); all rats survived the PTD/PF-treatment and were given three weeks to recover and return to a normal free-feeding weight. Prior to microdialysis and behavioral testing, every animal was handled for 4 days, 10 minutes each to reduce anxiety on the maze as well as food restricted overnight to increase maze exploration.

In-Vivo Microdialysis, Drug infusions & Behavioral testing: During the two days of in-vivo microdialysis and behavioral testing, samples from the posterior ventral hippocampus were collected and later analyzed for ACh levels. Subjects were transported to the same testing room and holding cage as Experiment 1. The microdialysis probe (CMA/11, 3 mm) was connected to a microinfusion system and perfused continuously at a rate of 2 µl/min with aCSF. Following an initial period of habituation to the cage (60 minutes), dialysate samples (volume ~12 µl) were collected every 6 minutes for a period of 18 minutes to determine basal levels of ACh in awake rats. After the 18 minutes of baseline sample were collection, the drug infusion needle (Plastics One, 15mm) was inserted into the drug cannula and infused over 1 minute with 1µl of either saline or bicuculline (0.5µg/µl). The drug infusion phase was a total of 18 minutes (3 samples). Within each group of 10 rats (PF and PTD), 5 rats received the saline infusion on the first day and bicuculline (0.5µg/µl) on the second session; the remaining 5 rats had the counterbalanced sequence (bicuculline on session 1, saline on session 2).

Eighteen minutes after the start of infusion, the animal was placed on the plus-maze to perform non-rewarded spontaneous alternation for 18 minutes. The plus maze used for behavioral testing was made of wood with clear Plexiglas sidewalls (12cm high) and a painted black floor with four arms of equal distance (55cm) and was elevated 80cm from the floor. The rat was

allowed to transverse the maze freely for an 18 minute period, the number and sequence of arms entered were recorded to determine alternation scores. The percent alternation score was equal to the ratio of: (actual alternation/possible alternations) $\times$ 100. The maze and post-baseline phases consisted of 3 – 6 minute samples for a total of 18 minutes each.

To test the reliability of the microdialysis probe, after the post baseline phase, it was placed into a standard solution (100 nM concentration of ACh and choline) to assess recovery rate (RR). After collection, brain dialysis samples were frozen for later analysis using HPLC.

3. Results

3.1. Experiment 1

Figure 1AB displays PTD thalamic neuropathology compared to a PF brain. Representative drug cannula placement in the medial septum and microdialysis cannula placement in the ventral hippocampus are shown in Figure 2AB. Before ACh levels were analyzed, it was confirmed that there was no overall order effect in terms of drug dose (low dose first vs. high dose first) on ACh release ($F [1, 10] < 1$).

3.1.1. Hippocampal ACh Levels—A single-factor (Group: PF vs. PTD) ANOVA revealed that basal amounts of ACh in the hippocampus were not different as a function of Group (PF = 34.67 ± 14.3 fmol; PTD = 56.04 ± 22.9 fmol; $F [1, 10] < 1$). A mixed design of one between-subjects (Group), one within-subjects (Phase: Baseline, Saline) factor ANOVA that compared ACh efflux during baseline and after saline infusion revealed that there was a non-significant rise above baseline (20%) in ACh levels after saline infusion across both groups. Therefore, the drug rise was calculated as a percent rise above saline to account for the effect of the injection alone. The mixed design of one between-subjects (Group), two within-subjects (Dose [0.5 ug/ul, 0.75 ug/ul]; Sample time [D1-D6]) factorial ANOVA revealed a significant main effect for Sample time ($F [5, 50] = 6.81$; $p < .0001$) and a significant interaction of Group $\times$ Dose ($F [1, 10] = 5.47$; $p < .05$). To determine the locus of this interaction, post-hoc Fisher's LSD t-tests contrasting the Group difference collapsed across Sample time were conducted at each dose. As shown in Figure 3, the groups did not differ in their ACh efflux levels to the challenge of 0.50 $\mu\text{g}/\mu\text{l}$ of bicuculline (main effect of Group: $t [10] < 1$). However, when subjects were administered the high dose, the rise in the PF rats, relative to the PTD rats, was greater $t [10] = 2.77$, $p < .05$). Given that the low dose (0.50 $\mu\text{g}/\mu\text{l}$) of bicuculline created an equal and significant rise of ACh in both PF and PTD animals, this was the only dose behaviorally tested in Experiment 2.

3.2. Experiment 2

Two PF and two PTD animals had misplaced cannula placement and were therefore not included in the final data analysis (final subject totals: PF = 8; PTD = 8). Once again, an analysis of order effects revealed that counterbalancing the drug condition across sessions did not affect alternation rates, number of arms entered or ACh levels in either group (all p 's $> .17$).

3.2.1. Behavioral Performance—Figure 4A displays a significant difference in percent alternation between PF and PTD-treated rats under saline, with bicuculline resulting in behavioral recovery in PTD rats. A two-factor ANOVA (2 $\times$ 2: Group: PF, PTD; Drug: Saline, Bicuculline) revealed a significant effect of Group (PF vs. PTD: $F [1, 14] = 4.89$; $p < .05$) and a significant effect of Drug $F [1, 14] = 5.58$; $p < .05$), but no interaction between Group and Drug ($F [1, 14] = 2.78$). The Group effect was primarily due to the PF rats alternating at a significantly higher rate than the PTD rats during the saline condition ($F [1, 14] = 9.78$; $p < .01$), as there was no significant difference between the Groups in the bicuculline condition ($F [1, 14] < 1$). The main effect of Drug was a result of bicuculline infusion into the septum

increasing alternation rates in both groups; however, follow-up analyses using an one within-subjects factor (Drug: Saline, 0.5 $\mu\text{g}/\mu\text{l}$) ANOVA for each Group condition revealed the effect of the drug was significant in the PTD rats (17.38%; $F [1,7] = 5.21$, $p = .05$), but not in the PF rats (3%, $F [1,7] < 1$).

As shown in Figure 4B, a two-factor (Group, Drug) ANOVA demonstrated that there were no differences in the number of arms entered as a function of Group ($F [1, 14] = 1.23$; $p = .29$), Drug ($F [1, 14] < 1$) or the interaction of Group $\times$ Drug ($F [1, 14] < 1$).

3.2.2. Hippocampal ACh Levels—Differences in basal amounts of ACh in the hippocampus (Saline: PF = 19.63 ± 5.7 , PTD = 22.42 ± 6.2 fmol; Bicuculline: PF = 19.2 ± 4.1 , PTD = 23.29 ± 4.7 fmol) were assessed with a one-factor (Group) ANOVA. This analysis revealed there was no differences in basal ACh levels between the Groups ($F [1, 10] < 1$). A mixed statistical model of one between-subjects (PF vs. PTD), three within-subjects [Drug (Saline, Bicuculline); Phase (Baseline, Infusion, Maze, After); Sample (One, Two, Three)] factorial ANOVA revealed a significant main effect for Phase ($F [3, 42] = 16.32$; $p < .0001$), a significant Drug $\times$ Phase interaction ($F [3, 42] = 4.37$; $p < .01$) and a significant Phase $\times$ Sample interaction ($F [6, 84] = 2.76$; $p < .05$). The overall main effect of Group was not significant and it did not interact with any variables.

Follow-up investigations of the Drug $\times$ Phase interaction were conducted using one between-subject (Group), one between-subject (Saline, Drug) factorial ANOVAs for each Phase. These analyses revealed that relative to the saline condition, intraseptal infusion of 0.50 $\mu\text{g}/\mu\text{l}$ of bicuculline increased hippocampal ACh efflux in both PF and PTD animals during the infusion phase and during maze testing (both F 's $[1,15] > 5.0$, p 's $< .05$), but not in the post-baseline phase ($F [1,15] < 1$).

Planned comparisons were conducted to assess whether the groups displayed differential changes in hippocampal ACh efflux to the drug challenge itself or to how the drug changed the ACh profile during behavior. Changes in ACh efflux were assessed separately in each group using a one within-subjects (Phase) factorial ANOVA. This analysis revealed that in the saline condition the PF rats had a significant rise above baseline in ACh levels during the Drug infusion phase ($F [1, 7] = 7.35$; $p < .05$) and Maze testing phase (see Figure 5A; $F [1, 7] = 7.01$; $p < .05$). However, after saline infusion PTD-treated rats only had a significant rise in ACh levels during the Maze testing phase ($F [1, 7] = 8.91$; $p < .05$; Figure 5B).

Intraseptal bicuculline administration increased hippocampal ACh efflux during Drug infusion (both F 's $[1, 7] > 12.70$; p 's $< .01$) and Maze testing (both F 's $[1, 7] = 12.30$; p 's $< .01$), relative to baseline ACh levels, in both PF and PTD rats. However, relative to the effect of the saline infusion, as assessed by one within-subjects factor ANOVAs for each group, the effect of increased ACh efflux during the bicuculline infusion phase was not significant in the PF rats ($F [1, 7] = 1.21$), but was in the PTD rats ($F [1, 7] = 6.34$; $p < .05$). This effect was a result of the increased reactivity of the PF rats to the saline infusion.

4. Discussion

The results from Experiment 1 demonstrated that intraseptal bicuculline administration increases hippocampal ACh levels in both PF and PTD animals. The doses and volume of bicuculline chosen for this study were based on a previous study in normal animals (Chang et al., 2006). In the PTD animals the low dose of bicuculline increased ACh levels by $167\% \pm 33.5$, whereas the high dose of bicuculline increased ACh levels by $131\% \pm 21.6$. However, in PF rats the low dose resulted in an increase of $177\% \pm 20.5$ and the high dose resulted in an increase of $225\% \pm 51.8$. There was less of a hippocampal ACh response to the intraseptal

administration of the high dose (0.75 $\mu\text{g}/\mu\text{l}$) of bicuculline in the PTD rats. Previous studies have shown that PTD animals can react differently to pharmacological manipulations than control animals (Nakagawasai et al., 2001; Savage et al., 1999). The 0.5 $\mu\text{g}/\mu\text{l}$ dose of bicuculline produced comparable changes in ACh efflux in both the groups and was therefore chosen for behavioral analysis. This suggests that antagonizing the GABA_A receptor could be an effective way to modulate ACh function and potentially lead to some behavioral recovery in amnesic subjects.

This is exactly what Experiment 2 illustrated: Intraseptal bicuculline led to complete behavioral recovery on the spontaneous alternation task without altering activity level on the maze. After bicuculline infusion, PTD animals alternated at levels comparable to saline-treated PF animals. Similar to Experiment 1, there was a comparable increase in hippocampal ACh levels in both groups in response to bicuculline administration. Although bicuculline infusion increased alternation performance in both groups, the percent of enhancement was only significant in the PTD group (PTD = 17%; PF = 3%). It should be noted that further increases in hippocampal ACh levels do not always lead to better performance. In normal rats increased hippocampal ACh efflux by an acetylcholinesterase inhibitor can actually lead to impaired behavior (Roland et al., 2008). Thus, there does appear to be an optimal range of cholinergic activity for optimal behavioral and hippocampal function.

Studies that have previously administered bicuculline into the medial septum of normal animals have produced mixed results. Post-training intraseptal bicuculline has been shown to increase memory retention in a passive avoidance task (Farr et al., 1999; Zarrindast et al., 2002) as well as performance in a stimulus-response task (Messier et al., 1999). However, others using young-normal rats have found that intraseptal bicuculline given either pre- or post-training can disrupt spatial working memory (Chrobak and Napier, 1992, 1991). Studies using various doses of intraseptal oxotremorine (Bunce et al., 2003), intraseptal carbachol (Bunce et al., 2004), intraseptal tacrine (Sabolek et al., 2005), and intrahippocampal physostigmine (Roland et al., 2008) have all found dose-dependent decreases in spatial memory performance in normal rats. In the current study, intraseptal bicuculline increased performance in cognitively impaired animals on a spontaneous alternation task—but had little effect in control PF rats. This difference in results may be due to the compromised integrity of the MS/DB cholinergic neurons within the PTD animal as well as task difficulty. Increasing the interval between trials on alternation can be a more sensitive assay for cognitive screening (Deacon and Rawlins, 2006).

Earlier reports have suggested that endogenous GABA could be inhibiting the MS/DB cholinergic neurons and affecting hippocampal ACh release by acting on the GABA_A receptor located on the cholinergic neurons (Degroot and Parent, 2001). PTD-treated animals display a degeneration of MS/DB cholinergic neurons (Pitkin and Savage, 2004, 2001; Savage et al., 2007) as well as a decrease in functional hippocampal ACh release (Roland et al., 2008; Roland and Savage, 2007; Savage et al., 2003), which could contribute to a reduction in hippocampal pyramidal cell activity.

In the present study, both saline and bicuculline increased hippocampal ACh release. However, factors such as handling can evoke significant ACh efflux (see Nilsson et al., 1990) and the handling during the saline injection procedure could have caused a rise in ACh levels in both groups. Thus, the ACh response to the saline infusion procedure may have masked the traditional efflux differences seen between PTD and PF rats during behavioral testing. However, the rise in ACh induced by either behavioral testing or bicuculline infusion was greater than that produced by the saline injection. Bicuculline produces antagonism of the GABAergic neurons as well as an increase in ACh release in cholinergic neurons (Moor et al.,

1998b); together these mechanisms could be beneficial in balancing the septo-hippocampal system within the PTD animal.

Wernicke-Korsakoff's syndrome is still a prevalent disease in many parts of the world and is widely under diagnosed in both adults and children (Harper, 2006; Sechi and Serra, 2007). The neuropsychiatric symptoms associated with WKS such as ocular abnormalities, changes in mental state and ataxia are usually treated with doses of thiamine when the patient is diagnosed. Nevertheless, treatment with thiamine alone is not effective at treating the anterograde amnesia related to WKS (Thomson and Marshall, 2006). Acetylcholinesterase inhibitors have been used with partial success for the treatment of the cognitive impairment associated with WKS (Angunawela and Barker, 2001; Cochrane et al., 2005; Luykx et al., 2008). Although bicuculline itself may not be the appropriate therapeutic tool, drugs influencing the GABAergic-cholinergic interactions in the septohippocampal pathway could be beneficial for the treatment of amnesic disorders.

Studies have demonstrated that increasing septohippocampal GABAergic pathway activity ultimately results in increased hippocampal pyramidal cell activity (Manns et al., 2001; Wu et al., 2003a). Therefore, drugs that increase the activity of the GABAergic septohippocampal pathway either alone or in combination with an AChE inhibitor could be more successful than an AChE inhibitor alone. In the past, GABA_A antagonists specific for the benzodiazepine binding site have improved cognition but have high pro-convulsant side effects (Duka et al., 1996; Little et al., 1984; Venault et al., 1986). Recently a new class of benzodiazepine inverse agonists specific for the $\alpha 5$ GABA receptor subunit have been shown to improve cognitive function with little to no pro-convulsant effects (Ballard et al., in press; Quirk et al., 1996; Sternfeld et al., 2004). These drugs are currently being investigated in humans and could be beneficial in WKS patients (Nutt et al., 2007). Overall, the involvement of the septohippocampal GABAergic neurons in the pathology of PTD-treatment and WKS requires further investigation to possibly determine more efficient treatment strategies.

Acknowledgments

This work was funded by NINDS Grant RO1 NS054272. The authors would like to thank Michelle Levinson, Steven Anzalone, and Jess Blackwolf for their help with behavioral testing and histology. Jessica Roland is now at the Department of Veterans Affairs in East Orange, NJ and the University of Medicine & Dentistry of New Jersey.

References

- Aggleton JP, Brown MW. Episodic memory, amnesia, and the hippocampal anterior thalamic axis. *Behav Brain Sci* 1999;22:425–489. [PubMed: 11301518]
- Alreja M, Wu M, Liu W, Atkins JB, Leranath C, Shanabrough M. Muscarinic Tone Sustains Impulse Flow in the Septohippocampal GABA But Not Cholinergic Pathway: Implications for Learning and Memory. *The Journal of Neuroscience* 2000b;20:8103–8110. [PubMed: 11050132]
- Angunawela II, Barker A. Anticholinesterase drugs for alcoholic Korsakoff syndrome. *Int J Geriatr Psychiatry* 2001;16:337–339. [PubMed: 11288170]
- Ballard TM, Knoflach F, Prinssen E, Borroni E, Vivian JA, Basile J, Gasser R, Moreau JL, Wettstein JG, Buettelmann B, Knust H, Thomas AW, Trube G, Hernandez MC. RO4938581, a novel cognitive enhancer acting at GABA-A α -5 subunit-containing receptors. *Psychopharmacology*. in press
- Bunce JG, Sabolek HR, Chrobak JJ. Intraseptal infusion of oxotremorine impairs memory in a delayed-non-match-to-sample radial maze task. *Neuroscience* 2003;121:259–267. [PubMed: 14521985]
- Bunce JG, Sabolek HR, Chrobak JJ. Intraseptal Infusion of the Cholinergic Agonist Carbachol Impairs Delayed-Non-Match-to-Sample Radial Arm Maze Performance in the Rat. *Hippocampus* 2004;14:450–459. [PubMed: 15224982]

- Butterworth RF, Heroux M. Effect of Pyridoxamine Treatment and Subsequent Thiamine Rehabilitation on Regional Cerebral Amino Acids and Thiamine-Dependent Enzymes. *J Neurochem* 1989;52:1079–1084. [PubMed: 2564421]
- Chang Q, Gold PE. Intra-hippocampal lidocaine injections impair acquisition of a place task and facilitate acquisition of a response task in rats. *Behav Brain Res* 2003a;144:19–24. [PubMed: 12946591]
- Chang Q, Savage LM, Gold PE. Microdialysis measures of functional increases in ACh release in the hippocampus with and without inclusion of acetylcholinesterase inhibitors in the perfusate. *J Neurochem* 2006;97:697–706. [PubMed: 16579834]
- Chrobak JJ, Napier TC. Antagonism of GABAergic Transmission within the Septum Disrupts Working/Episodic Memory in the Rat. *Neuroscience* 1992;47:833–841. [PubMed: 1579213]
- Chrobak JJ, Napier TC. Intraseptal Administration of Bicuculline Produces Working Memory Impairments in the Rat. *Behav Neural Biol* 1991;55:247–254. [PubMed: 1647762]
- Cochrane M, Cochrane A, Jauhar P, Ashton E. Acetylcholinesterase inhibitors for the treatment of Wernicke-Korsakoff Syndrome - three further cases show responses to donepezil. *Alcohol Alcohol* 2005;40:151–154. [PubMed: 15596427]
- Deacon RMJ, Rawlins JNP. T-maze alternation in the rodent. *Nature Protocols* 2006;1:7–12.
- Degroot A, Parent MB. Infusions of physostigmine into the hippocampus or the entorhinal cortex attenuate avoidance retention deficits produced by intra-septal infusions of the GABA agonist muscimol. *Brain Res* 2001;920:10–18. [PubMed: 11716806]
- Duka T, Ott H, Rohloff A, Voet B. The effects of a benzodiazepine receptor antagonist beta-carboline ZK-93426 on scopolamine-induced impairment on attention, memory and psychomotor skills. *Psychopharmacology* 1996;123:361–373. [PubMed: 8867876]
- Farr SA, Uezu K, Flood JF, Morley JE. Septo-hippocampal drug interactions in post-trial memory processing. *Brain Res* 1999;847:221–230. [PubMed: 10575091]
- Gao B, Hornung JP, Fritschy JM. Identification of distinct GABA-A receptor subtypes in cholinergic and parvalbumin-positive neurons of the rat and marmoset medial septum diagonal band complex. *Neuroscience* 1995;65:101–117. [PubMed: 7753393]
- Harper C. Thiamine (vitamin B1) deficiency and associated brain damage is still common throughout the world and prevention is simple and safe! *Eur J Neurol* 2006;13:1078–1082. [PubMed: 16987159]
- Harper C, Kril J. Pathological changes in alcoholic brain shrinkage. *The Medical Journal of Australia* 1986;144:3–4. [PubMed: 3941617]
- Kopelman MD. The Korsakoff Syndrome. *Br J Psychiatry* 1995;166:154–173. [PubMed: 7728359]
- Krebs-Kraft DL, Wheeler MG, Parent MB. The memory-impairing effects of septal GABA receptor activation involve GABAergic septo-hippocampal projection neurons. *Learn Memory* 2007;14:833–841.
- Langlais PJ, Mair RG, Anderson CD, McEntee WJ. Long-lasting changes in regional brain amino acids and monoamines in recovered pyridoxamine treated rats. *Neurochem Res* 1988;13:1199. [PubMed: 3237312]
- Langlais PJ, Savage LM. Thiamine deficiency in rats produces cognitive and memory deficits on spatial tasks that correlate with tissue loss in diencephalon, cortex and white matter. *Behav Brain Res* 1995;68:75–89. [PubMed: 7619308]
- Langlais PJ, Zhang SX. Cortical and Subcortical White Matter Damage without Wernicke's Encephalopathy after Recovery from Thiamine Deficiency in the Rat. *Alcohol Clin Exp Res* 1997;21:434–443. [PubMed: 9161603]
- Langlais PJ, Zhang SX, Savage LM. Neuropathology of thiamine deficiency: An update on the comparative analysis of human disorders and experimental models. *Metab Brain Dis* 1996;11:19–37. [PubMed: 8815388]
- Little HJ, Nutt DJ, Taylor SC. Acute and chronic effects of the benzodiazepine receptor ligand FG 7142: proconvulsant properties and kindling. *Br J Pharmacol* 1984;83:951–958. [PubMed: 6097329]
- Luykx HJ, Dorrestijn LDA, Haffmans PMJ, Bonebakker A, Kerkmeier M, Hendriks V. Rivastigmine in Wernicke-Korsakoff's syndrome: five patients with rivastigmine showed no more improvement than five patients without rivastigmine. *Alcohol Alcohol* 2008;43:70–72. [PubMed: 17984137]
- Mair RG. On the role of thalamic pathology in diencephalic amnesia. *Rev Neurosci* 1994;5:105–140. [PubMed: 7827707]

- Manns ID, Mainville L, Jones BE. Evidence for glutamate, in addition to acetylcholine and GABA, neurotransmitter synthesis in basal forebrain neurons projecting to the entorhinal cortex. *Neuroscience* 2001;107:249–263. [PubMed: 11731099]
- Manseau F, Danik M, Williams S. A functional glutamatergic neurone network in the medial septum and diagonal band area. *J Physiol* 2005;566:865–884. [PubMed: 15919710]
- Messier C, Wall PM, Ethier K. Contribution of cholinergic and gabaergic functions to memory processes in BALB / cAnNCrIBR mice. *Brain Res* 1999;818:583–592. [PubMed: 10082853]
- Moor E, DeBoer P, Westerink BHC. GABA receptors and benzodiazepine binding sites modulate hippocampal acetylcholine release in vivo. *Eur J Pharmacol* 1998a;359:119–126. [PubMed: 9832381]
- Moor E, Schirm E, Jacso J, Westerink BHC. Involvement of medial septal glutamate and GABA receptors in behaviour-induced acetylcholine release in the hippocampus: A dual probe microdialysis study. *Brain Res* 1998b;789:1–8. [PubMed: 9602020]
- Mulder J, Harkany T, Czollner K, Cremers TIFH, Keijser JN, Nyakas C, Luiten PGM. Galantamine-induced behavioral recovery after sublethal excitotoxic lesions to the rat medial septum. *Behav Brain Res* 2005;163:33–41. [PubMed: 15951032]
- Nakagawasai O, Tadano T, Hozumi S, Tan-No K, Nijima F, Kisara K. Immunohistochemical estimation of brain choline acetyltransferase and somatostatin related to the impairment of avoidance learning induced by thiamine deficiency. *Brain Res Bull* 2000b;52:189–196. [PubMed: 10822160]
- Nakagawasai O, Tadano T, Hozumi S, Taniguchi R, Yamadera F, Tan-No K, Nijima F, Arai H, Yasuhara H, Kinemuchi H, Kisara K. Involvement of Muscarinic Receptor on the Impairment of Avoidance Learning in Mice Fed a Thiamine-Deficient Diet. *Biogenic Amines* 2001;16:199–210.
- Nilsson OG, Kalen P, Rosengren E, Bjorklund A. Acetylcholine release in the rat hippocampus as studied by microdialysis is dependent on axonal impulse flow and increases during behavioural activation. *Neuroscience* 1990;36:325–338. [PubMed: 2215927]
- Nutt DJ, Beeson M, Wilson SJ, Dawson GR, Lingford-Hughes AR. Blockade of alcohol's amnesic activity in humans by an alpha-5 subtype benzodiazepine receptor inverse agonist. *Neuropsychopharmacology* 2007;53:810–820.
- Paxinos, G.; Watson, C. *The Rat Brain in Stereotaxic Coordinates*. San Diego, CA: Academic Press; 1986.
- Pires RGW, Pereira SRC, Oliveria-Silva IF, Franco GC, Ribeiro AM. Cholinergic parameters and the retrieval of learned and re-learned spatial information: A study using a model of Wernicke–Korsakoff Syndrome. *Behav Brain Res* 2005;162:11–21. [PubMed: 15922063]
- Pitkin SR, Savage LM. Age-related vulnerability to diencephalic amnesia produced by thiamine deficiency: the role of time of insult. *Behav Brain Res* 2004;148:93–105. [PubMed: 14684251]
- Pitkin SR, Savage LM. Aging potentiates the acute and chronic neurological symptoms of pyriithiamine-induced thiamine deficiency in the rodent. *Behav Brain Res* 2001;119:167–177. [PubMed: 11165332]
- Quirk K, Blurton P, Fletcher S, Leeson P, Tang F, Mellilo D, Ragan CI, McKernan RM. [3H]L-655,708, a novel ligand selective for the benzodiazepine site of GABA-A receptors which contain the alpha-5 subunit. *Neuropsychopharmacology* 1996;35:1331–1335.
- Reed LJ, Lasserson D, Marsden P, Stanhope N, Stevens T, Bello F, Kingsley D, Colchester A, Kopelman MD. FDP-PET findings in the Wernicke-Korsakoff syndrome. *Cortex* 2003;39:1027–1045. [PubMed: 14584565]
- Roland JJ, Mark K, Vetreno RP, Savage LM. Increasing Hippocampal Acetylcholine Levels Enhances Behavioral Performance in an Animal Model of Diencephalic Amnesia. *Brain Res* 2008;1234:116–127. [PubMed: 18706897]
- Roland JJ, Savage LM. Blunted hippocampal, but not striatal, acetylcholine efflux parallels learning impairment in diencephalic-lesioned rats. *Neurobiol Learn Mem* 2007;87:123–132. [PubMed: 16978888]
- Sabolek HR, Bunce JG, Chrobak JJ. Intraseptal tacrine-induced disruptions of spatial memory performance. *Behav Brain Res* 2005;158:1–7. [PubMed: 15680189]
- Savage LM, Chang Q, Gold PE. Diencephalic Damage Decreases Hippocampal Acetylcholine Release During Spontaneous Alternation Testing. *Learn Memory* 2003;10:242–246.

- Savage LM, Pitkin SR, Knitowski KM. Rats exposed to acute pyridoxamine-induced thiamine deficiency are more sensitive to the amnesic effects of scopolamine and MK-801: examination of working memory, response selection, and reinforcement contingencies. *Behav Brain Res* 1999;104:13–26. [PubMed: 11125732]
- Savage LM, Roland JJ, Klintsova AY. Selective septohippocampal – but not forebrain amygdalar – cholinergic dysfunction in diencephalic amnesia. *Brain Res* 2007;1139:210–219. [PubMed: 17289001]
- Sechi GP, Serra A. Wernicke's encephalopathy: new clinical settings and recent advances in diagnosis and management. *Lancet Neurology* 2007;6:442–455. [PubMed: 17434099]
- Squire LR. Two forms of human amnesia: an analysis of forgetting. *The Journal of Neuroscience* 1981;1:635–640. [PubMed: 7346571]
- Sternfeld F, Carling RW, Jelley RA, Ladduwahetty T, Merchant KJ, Moore KW, Reeve AJ, Street LJ, O'Connor D, Sohal B, Attack JR, Cook S, Seabrook G, Wafford K, Tattersall FD, Collinson N, Dawson GR, Castro JL, MacLeod AM. Selective, orally active GABA-A alpha-5 receptor inverse agonists as cognitive enhancers. *Journal of Medicinal Chemistry* 2004;47:2176–2179. [PubMed: 15084116]
- Sullivan EV, Marsh L. Hippocampal volume deficits in alcoholic Korsakoff's syndrome. *Neurology* 2003;61:1716–1719. [PubMed: 14694035]
- Thomson AD, Marshall EJ. The treatment of patients at risk of developing Wernicke's Encephalopathy in the community. *Alcohol Alcohol Suppl* 2006;41:159–167.
- Toth K, Freund TF, Miles R. Disinhibition of rat hippocampal pyramidal cells by GABAergic afferents from the septum. *J Physiol* 1997;500:463–474. [PubMed: 9147330]
- Van der Zee EA, Luiten PGM. Cholinergic and GABAergic neurons in the rat medial septum express muscarinic acetylcholine receptors. *Brain Res* 1994;652:263–272. [PubMed: 7953739]
- Venault P, Chapouthier G, de Carvalho LP, Simiand J, Morre M, Dodd RH, Rossier J. Benzodiazepine impairs and betacarboline enhances performance in learning and memory tasks. *Nature* 1986;321:864–866. [PubMed: 3724846]
- Victor, M.; Adams, RD.; Collins, GH. *The Wernicke-Korsakoff syndrome*. Philadelphia: F.A. Davis; 1971.
- Wainer BH, Levey AI, Rye DB, Mesulam MM, Mufson EJ. Cholinergic and noncholinergic septohippocampal pathways. *Neurosci Lett* 1985;54:45–52. [PubMed: 3974944]
- Wu M, Hajszan T, Leranth C, Alreja M. Nicotine recruits a local glutamatergic circuit to excite septohippocampal GABAergic neurons. *Eur J Neurosci* 2003a;18:1155–1168. [PubMed: 12956714]
- Wu M, Newton SS, Atkins JB, Xu C, Dunman RS, Alreja M. Acetylcholinesterase Inhibitors Activate Septohippocampal GABAergic Neurons via Muscarinic but Not Nicotinic Receptors. *The Journal of Pharmacology and Experimental Therapeutics* 2003b;307:535–543. [PubMed: 12966162]
- Zarrindast MR, Bakhsha A, Rostami P, Shafaghi B. Effects of intrahippocampal injection of GABAergic drugs on memory retention of passive avoidance learning in rats. *J Psychopharm* 2002;16:313–319.
- Zarrindast MR, Shamsi T, Azarmina P, Rostami P, Shafaghi B. GABAergic system and imipramine-induced impairment of memory retention in rats. *European Journal of Neuropsychopharmacology* 2004;14:59–64.

A. Thalamus of PF rat

B. Thalamus of PTD rat

Figure 1. Cresyl violet stained sections illustrating the intact thalamus of a PF control (**A**) and a midline thalamic lesion produced by PTD-treatment (**B**). The arrows indicate the prototypical PTD-induced lesion.

A. Medial Septum Drug Cannula **B. Hippocampal Microdialysis Cannula** **Figure 2.**

Cresyl violet stained section showing acceptable probe placement for the medial septal drug cannula (**A**) and the hippocampal microdialysis cannula (**B**). The arrows indicate the regions of infusion (**A**) or occupied by the microdialysis cannula (**B**).

Figure 3.

Experiment 1: Bicuculline dose response curve for both the PF (A) and PTD (B) animals. Rise in ACh release (Mean $\pm$ SEM) after bicuculline infusion was calculated as the percent rise above the ACh release caused by saline infusion. Saline infusion occurred at time point S and bicuculline infusion occurred at time point D1. Each sample (S, D1-D6) represents a 15-min time bin of microdialysis collection.

Figure 4.

Experiment 2: Behavioral data (Mean $\pm$ SEM) from spontaneous alternation testing displaying the percent alternation (**A**) and activity on the maze (**B**) after both saline and bicuculline administration in PF and PTD rats. Panel A displays the significant difference between PTD and PF rats in alternation behavior after saline infusion. Although intraseptal bicuculline infusion enhanced alternation rates, the effect was only significant in PTD animals. Panel B shows that maze activity did not change as a function of drug in either group.

Figure 5.

Experiment 2: Profiles of hippocampal ACh release (Mean percent rise above baseline $\pm$ SEM) after saline and bicuculline infusions in PF (A) and PTD (B) animals shown in 6-minute sample bins during baseline, drug infusion, maze testing and after-baseline phases. Under saline, PF rats had a significant rise above baseline in ACh levels during the drug infusion phase and maze testing phases (A), while PTD-treated rats only had a significant rise in ACh levels during the maze testing phase (B). After bicuculline administration, both PF (A) and PTD (B) animals showed a significant rise above baseline during the drug infusion and maze testing phases.