A. A. Lashkov,^a
A. G. Gabdoulkhakov,^a
A. A. Shtil^b and
A. M. Mikhailov^{a*}

^aA. V. Shubnikov Institute of Crystallography,
Russian Academy of Sciences, Leninskiy
Prospect 59, 119333 Moscow, Russia, and
^bBlokhin Cancer Center, Russian Academy of
Medical Sciences, Kashirskoe Shosse 24,
115478 Moscow, Russia

Correspondence e-mail: amm@ns.crys.ras.ru

Received 3 April 2009

Accepted 29 April 2009

Crystallization and preliminary X-ray diffraction analysis of *Salmonella typhimurium* uridine phosphorylase complexed with 5-fluorouracil

Uridine phosphorylase (UPh; EC 2.4.2.3) catalyzes the phosphorolytic cleavage of the N-glycosidic bond of uridine to form ribose 1-phosphate and uracil. This enzyme also activates pyrimidine-containing drugs, including 5-fluorouracil (5-FU). In order to better understand the mechanism of the enzyme–drug interaction, the complex of *Salmonella typhimurium* UPh with 5-FU was cocrystallized using the hanging-drop vapour-diffusion method at 294 K. X-ray diffraction data were collected to 2.2 Å resolution. Analysis of these data revealed that the crystal belonged to space group C2, with unit-cell parameters $a = 158.26$, $b = 93.04$, $c = 149.87$ Å, $\alpha = \gamma = 90^\circ$, $\beta = 90.65^\circ$. The solvent content was 45.85% assuming the presence of six hexameric molecules of the complex in the unit cell.

1. Introduction

5-Fluorouracil (5-FU) has been used in chemotherapeutic regimens in patients with gastrointestinal malignancies, including oesophageal, gastric, colon and pancreatic tumours, as well as breast cancer for several decades (Kemeny, 1987; Huang *et al.*, 2007). This drug competes with uracil for interaction with thymidylate synthase. 5-FU interferes with DNA and RNA synthesis, thereby causing cell death. Furthermore, 5-halogen-containing derivatives of uracil are also used as antimicrobial agents. Synergism of 5-FU with antibacterial agents such as ceftriaxone, ceftazidime, cefotiam, piperacillin and netilmicin has been shown (Price *et al.*, 1965; Gieringer *et al.*, 1986; Yamashiro *et al.*, 1986). However, toxicity towards haematopoietic cells, skin, heart and neurons can limit the therapeutic efficacy of 5-FU and structurally similar drugs (Peters & van Groenigen, 1991). Uridine phosphorylase (UPh) catalyzes the phosphorolytic cleavage of the N-glycosidic bond in uridine to produce ribose 1-phosphate and uracil (Leer *et al.*, 1977; Vita *et al.*, 1986). This enzyme activates pyrimidine-based drugs, including 5-FU (Cao *et al.*, 2002; Caradoc-Davies *et al.*, 2004). Therefore, studies of the structure of UPh bound to 5-FU should be instrumental in future modifications aimed at the design of 5-FU derivatives with lower general toxicity and retained antitumour and antimicrobial potencies (Iigo *et al.*, 1990; Temmink *et al.*, 2006; Matsusaka *et al.*, 2007).

2. Protein expression and purification

Salmonella typhimurium UPh (StUPh) was expressed in *Escherichia coli* and purified as reported elsewhere (Molchan *et al.*, 1998; Dontsova *et al.*, 2004). The *Escherichia coli* BL21 (DE3) strain was transformed with the recombinant plasmid and grown on solid LB agar for 12 h at 310 K. Protein synthesis was stimulated with 0.5 mM isopropyl β -D-1-thiogalactopyranoside. The biomass was sonicated, ammonium sulfate was added to precipitate the proteins and the pellets were dissolved in buffer (pH 7.2) containing 50 mM KH_2PO_4 and 0.5 mM β -mercaptoethanol. Further steps of StUPh purification were performed using chromatography on butyl Sepharose as a first step and Q-Sepharose as the final step. The enzymatic activity was 280 units per milligram of purified protein. The homogeneity of the purified StUPh was 96% as determined by nondenaturing gel electrophoresis.

Table 1

Statistics of X-ray data.

Values in parentheses are for the last resolution shell.

Wavelength (Å)	0.918
Temperature (K)	100
Oscillation (°)	0.5
Space group	C2
Unit-cell parameters (Å, °)	$a = 158.26$, $b = 93.04$, $c = 149.87$, $\alpha = \gamma = 90$, $\beta = 90.65$
Molecules per unit cell	6
V_M (Å ³ Da ⁻¹)	2.27
Solvent content (%)	45.85
Resolution limits (Å)	30.0–2.2 (2.25–2.2)
Completeness (%)	90.2 (79.3)
No. of reflections	324446
No. of unique reflections	99573 (7124)
R_{observed} (%)	6.8 (55.6)
R_{expected} (%)	7.0 (55.3)
$\langle I/\sigma(I) \rangle$	12.67 (2.16)
R_{meas}	0.08 (0.66)
R_{merge}	0.017 (0.47)
Redundancy	3.26 (3.24)

3. Crystallization of the StUph–5FU complex

Crystals of the complex of StUph with 5-FU (EBEWE Pharma, Austria) were obtained by cocrystallization (Fig. 1). Crystallization was performed on siliconized glass slides (Hampton Research, USA) in Linbro plates at 294 K using the hanging-drop vapour-diffusion method. The reservoir solution (0.5 ml) consisted of 0.34 ml 0.1 M Tris–maleate–NaOH buffer pH 5.5 and 0.16 ml 40% (w/v) polyethylene glycol 3350. The crystallization drop contained 2 µl StUph solution (11.3 mg ml⁻¹) in 10 mM Tris–HCl buffer pH 7.3, 2 µl H₂O, 1.3 µl reservoir solution, 2 µl 100 mM 5-FU and 0.3 µl 2-propanol. Crystals of dimensions 0.07 × 0.3 × 0.5 mm were obtained after 1–2 weeks and were used for X-ray diffraction analysis.

4. X-ray analysis

The X-ray data set (Table 1) was collected upon irradiation of StUph–5FU crystals under cryogenic conditions (100 K) on beamline 14.2 at BESSY, Berlin, Germany. The wavelength was 0.9184 Å. A CHESS CCD detector was used with an oscillation range $\Delta\phi$ of 0.5° and a crystal-to-detector distance of 240 mm. Prior to freezing in liquid nitrogen, the crystals were transferred into cryoprotectant solution containing 100 mM Tris–maleate–NaOH buffer pH 5.5, 25% (w/v) polyethylene glycol 3350 and 20% (v/v) anhydrous glycerol. All data were processed and scaled using XDS (Kabsch, 1988).

Figure 1
Crystal of StUph complexed with 5-FU.

300 images were obtained during the collection of X-ray diffraction data. All data were indexed, merged and processed using the XDS program with XYCORR, INIT, COLSPOT, IDXREF, DEFPIX, XPLAN, INTEGRATE and CORRECT options. For a semi-automatic determination of the space group, the minimal value of the XDS 'quality of fit' function was used. The crystals of the StUph–5-FU complex belonged to space group C2. Detailed data statistics are presented in Table 1.

The structure was resolved by the molecular-replacement technique using the Phaser program with rigid-body refinement option (McCoy, 2007). X-ray diffraction data from 10 to 2.5 Å resolution were used in this step. The X-ray structure of monomer A only of ligand-free StUph at 1.76 Å resolution (Timofeev *et al.*, 2007; PDB code 2oxf) was utilized as a search model. Water molecules were removed from the model. Six full homohexamers were found in the unit cell. The Matthews coefficient (Matthews, 1968) was 2.27 Å³ Da⁻¹ and the solvent content was 45.85% (Table 1). Only one solution was evident, with an *R* factor of 37.64% and a correlation coefficient *R*_{corr} of 76.95%.

To improve the phase model, one macrocycle of simulated annealing using the phenix.refine module of PHENIX (Adams *et al.*, 2002) was performed in the temperature range 12 000–300 K with 50 K steps and resolution 10–2.2 Å. Before refinement, 5% of the observations were chosen at random and set aside for cross-validation analysis and to monitor the various refinement strategies. Next, σ_A -weighted electron-density maps with coefficients $(2|F_o| - |F_c|)$ and $(|F_o| - |F_c|)$ were obtained using PHENIX. Using the $(|F_o| - |F_c|)$ electron-density map in the Coot program (Emsley & Cowtan, 2004), we identified 5-FU molecules and water molecules localized in the close vicinity of the 5-FU molecules. After two cycles of restrained refinement in REFMAC (Murshudov *et al.*, 1997) and the synthesis of σ_A -weighted $(2|F_o| - |F_c|)$ and $(|F_o| - |F_c|)$ electron-density maps the *R* factor was 26.42% and *R*_{free} was 29.24%.

One uracil-binding site in StUph complexed with 5-FU and water is shown in Fig. 2. The cartoon representation was generated with PyMOL (DeLano, 2008). The drug and water bind to the following atoms of the amino-acid residues in the uracil-binding site:

Figure 2
Preliminary structure of the uracil-binding site in the StUph–5-FU complex.

Arg223 NH1–water (2.85 Å), water–5-FU O4 (2.88 Å), Arg168 NH1–5-FU O4 (3.00 Å), Gln166 OE1–5-FU N3 (2.70 Å), Gln166 NE2–5-FU O2 (2.89 Å). Refinement of the structure is currently in progress and will be published elsewhere.

This work was supported partially by RFBR and Kaluga region administration (grant No. 09-02-97519).

References

- Adams, P. D., Grosse-Kunstleve, R. W., Hung, L.-W., Ioerger, T. R., McCoy, A. J., Moriarty, N. W., Read, R. J., Sacchettini, J. C., Sauter, N. K. & Terwilliger, T. C. (2002). *Acta Cryst. D* **58**, 1948–1954.
- Cao, D., Russell, R. L., Zhang, D., Leffert, J. J. & Pizzorno, G. (2002). *Cancer Res.* **62**, 2313–2317.
- Caradoc-Davies, T. T., Cutfield, S. M., Lamont, I. L. & Cutfield, J. F. (2004). *J. Mol. Biol.* **337**, 337–354.
- DeLano, W. L. (2008). *The PyMOL Molecular Graphics System*. <http://www.pymol.org>.
- Dontsova, M. V., Savochkina, Y. A., Gabdoulkhakov, A. G., Baidakov, S. N., Lyashenko, A. V., Zolotukhina, M., Errais Lopes, L., Garber, M. B., Morgunova, E. Y., Nikonov, S. V., Mironov, A. S., Ealick, S. E. & Mikhailov, A. M. (2004). *Acta Cryst. D* **60**, 709–711.
- Emsley, P. & Cowtan, K. (2004). *Acta Cryst. D* **60**, 2126–2132.
- Gieringer, J. H., Wenz, A. F., Just, H. M. & Daschner, F. D. (1986). *Chemotherapy*, **32**, 418–424.
- Huang, Z., Guo, K. J., Guo, R. X. & He, S. G. (2007). *Hepatobiliary Pancreat. Dis. Int.* **6**, 312–320.
- Kabsch, W. (1988). *J. Appl. Cryst.* **21**, 916–924.
- Kemeny, N. (1987). *Semin. Surg. Oncol.* **3**, 190–214.
- Leer, J. C., Hammer-Jespersen, K. & Schwartz, M. (1977). *Eur. J. Biochem.* **75**, 217–224.
- Iigo, M., Nishikata, K., Nakajima, Y., Szinai, I., Veres, Z., Szabolcs, A. & De Clercq, E. (1990). *Biochem. Pharmacol.* **39**, 1247–1253.
- Matsusaka, S., Yamasaki, H., Fukushima, M. & Wakabayashi, I. (2007). *Chemotherapy*, **53**, 36–41.
- Matthews, B. W. (1968). *J. Mol. Biol.* **33**, 491–497.
- McCoy, A. J. (2007). *Acta Cryst. D* **63**, 32–41.
- Molchan, O. K., Dmitrieva, N. A., Romanova, D. V., Errais Lopes, L., Debabov, V. G. & Mironov, A. S. (1998). *Biochemistry (Mosc.)*, **63**, 195–199.
- Murshudov, G. N., Vagin, A. A. & Dodson, E. J. (1997). *Acta Cryst. D* **53**, 240–255.
- Peters, G. J. & van Groeningen, C. J. (1991). *Ann. Oncol.* **2**, 469–480.
- Price, K. E., Bradner, W. T., Buck, R. E. & Lein, J. (1965). *Antimicrob. Agents Chemother.* **5**, 481–487.
- Temminck, O. H., de Bruin, M., Laan, A. C., Turksma, A. W., Cricca, S., Masterson, A. J., Noordhuis, P. & Peters, G. J. (2006). *Int. J. Biochem. Cell Biol.* **38**, 1759–1765.
- Timofeev, V. I., Pavlyuk, B. F., Lashkov, A. A., Seregina, A. G., Gabdulkhakov, A. G., Vainshtein, B. K. & Mikhailov, A. M. (2007). *Kristallografiya*, **52**, 1106–1113.
- Vita, A., Amici, A., Cacciamani, T., Lanciotti, M. & Magni, G. (1986). *Int. J. Biochem.* **18**, 431–435.
- Yamashiro, Y., Fukuoka, Y., Yotsuji, A., Yasuda, T., Saikawa, I. & Ueda, Y. (1986). *J. Antimicrob. Chemother.* **18**, 703–708.