Published in final edited form as:

Org Lett. 2008 December 18; 10(24): 5609–5612. doi:10.1021/ol802401a.

Chromium(0)-promoted $[6\pi+2\pi]$ cycloadditions of allenes with cycloheptatriene

James H. Rigby^{*}, Stéphane B. Laurent, Zeeshan Kamal, and Mary Jane Heeg[†]

Department of Chemistry, Wayne State University, Detroit, MI 48202-3489, USA

Abstract

Photoinitiated $[6\pi+2\pi]$ cycloadditions of allenes with $(\eta^6\text{-cycloheptatriene})\text{tricarboxylchromium(0)}$ (**1**) are described. An example of asymmetric induction obtained by reaction of **1** with a chiral 1,3-disubstituted allene is also reported.

Transition metal-mediated higher-order cycloaddition reactions have become important synthetic tools for the construction of medium-sized ring targets.¹ In particular, Cr(0)-promoted $[6\pi+4\pi]$ and $[6\pi+2\pi]$ cycloadditions with cyclic trienes have been extensively studied in the context of synthetic applications.^{2,3} These transformations can be initiated either photochemically or thermally; however, photochemical activation appears to be more compatible with sensitive reaction partners. Thus, isocyanates and ketenes have been successfully used as 2π partners in the photoinduced $[6\pi+2\pi]$ cycloaddition reactions with $(\text{cycloheptatriene})\text{tricarboxylchromium(0)}$ (**1**) whereas they were less successful under thermal conditions.⁴ In the light of these results, allenes were viewed as potentially viable 2π partners. Furthermore, the $[6\pi+2\pi]$ cycloaddition of chiral 1,3-disubstituted allenes could provide a simple route to optically active bicyclo[4.2.1]nonane ring systems (Scheme 1). To the best of our knowledge, the metal-mediated $[6\pi+2\pi]$ cycloaddition of allenes has not been described previously.⁵ Thermal, metal-free cycloadditions of allenes to cyclic trienes such as tropone and azaheptafulvenes have been studied; but only $[8+2]$ ⁶ or $[4+2]$ ⁷ cycloadducts were isolated depending on the nature of the allene.

Herein, results of a study on the photoinitiated $[6\pi+2\pi]$ cycloaddition of allenes with $(\eta^6\text{-cycloheptatriene})\text{-tricarboxylchromium(0)}$ (**1**) are reported.

All of the following cycloaddition reactions were performed in a pyrex reactor with a 450-W Canrad-Hanovia medium pressure Hg vapor lamp. Two different procedures were evaluated

E-mail: jhr@chem.wayne.edu.

[†]To whom inquiries about X-ray structures should be addressed.

in the study: (a) a mixture of complex **1** (1 equiv) and allene (2 equiv) in hexanes was directly irradiated (procedure A); (b) a solution of allene (2 equiv) in hexanes was added slowly to a solution of **1** in hexanes using a syringe pump under continuous irradiation (procedure B). A preliminary experiment was performed with complex **1** and 1,1-diphenyl-1,2-propadiene (**2a**) following procedure A. The metal-free [6+2]-cycloadduct **3a** was obtained in 50% yield as a single regioisomer (Table 1, entry 1). Similar results were observed with vinylidenecyclohexane (**2b**) and trisubstituted allene **2c** using procedure B, affording cycloadducts **3b** and **3c** respectively, in moderate yields (entries 2–3). In each case, cycloaddition occurred at the less sterically congested double bond of the 2η partner. Regioisomer **3c** was isolated as a single diastereomer resulting from an exclusive *endo* approach of the allene, paralleling the [6π+2π] cycloaddition with alkenes.^{8,9}

1,3-Disubstituted allenes were then examined. When allene **4a** was employed using procedure A, a single isomer **5a** was obtained in 59% yield (Table 2, entry 1). The addition occurred exclusively at the unsaturation bearing the CH₂OTBS substituent via an *endo* transition state. The phenyl group on the *exocyclic* double bond in **5a** was found to be *syn* coplanar with the (C-8)-(C-1) bond, and consequently, orientated away from the substituent at C-7.¹⁰ Subsequent X-ray crystallographic analysis performed on a derivative of **5a** (*p*-nitrobenzoate ester) confirmed the structural and stereochemical assignments (see Supporting Information). Surprisingly, when the same cycloaddition was performed using the ‘slow addition’ technique (procedure B), the *Z* isomer of **5a** with the phenyl group *syn* to C-7 could also be detected (*E/Z* = 85:15, entry 2). In rare cases, procedure B led to the virtually exclusive formation of the *E* isomer. For the moment, it is not evident how the rate of slow addition of allene promotes the formation of the *Z* isomer. The cycloaddition with **4b** (procedure A) behaved similarly to the cycloaddition with **4a** to give, after deprotection of the THP group, uniquely the alcohol (*E*)-*endo*-**5b** (56% over two steps, entry 3). Irradiation of **1** with **4c** (procedure B) gave a 88:12 mixture of regioisomers *endo*-**5c** and *endo*-**6c** in 64% overall yield (entry 4). Curiously, cycloaddition also occurred at the π bond bearing the phenyl group. The high regioselectivity observed when an ether group was present is consistent with a prior coordination of the oxygen atom to the chromium center which should favor bond formation at the π bond bearing the ether substituent. With a methyl group, no coordination is possible and the steric effects with **1** would be expected to control regioselectivity. Each of the regioisomers **5c** and **6c** was in turn isolated as a mixture of *Z* and *E* isomers. For **5c**, the *Z* isomer with the phenyl group orientated away from the methyl group at C-7 was found to be the major isomer in line with previous results (Table 2, entries 1–3), whereas for the major isomer of **6c**, the methyl group was found to be *syn* to the phenyl group at C-7.

Attention was then directed toward cycloaddition of **1** with monosubstituted allenes (Table 3). Preliminary experiments were carried out with aryl allenes **7a–d** bearing various substituents on the phenyl ring. Whenever an arylallene was used, the cycloaddition (procedure B) afforded the **8**-type regioisomer with excellent selectivity (ratios **8a–d**/**9a–d** > 97:3 by GC analysis of the crude mixtures) and with moderate yields (entries 1–4). Each regioisomer was isolated as an inseparable mixture of *Z* and *E* isomers in ratios of 50:50. In contrast to the case with 1,3-disubstituted allenes, the absence of a substituent at the C-7 position in the final products would reduce steric interactions and would allow the aryl substituent to be placed *syn* to C-7 more easily. When a mixture of **1** and ethyl buta-2,3-dienoate (**7e**) was irradiated, the cycloaddition occurred only at the terminal double bond of the allene to afford the regioisomer **8e** as a separable 65:35 mixture of *Z* and *E* isomers in 34% yield (entry 5). No addition to the electron-deficient double bond of the allene was observed. This result is in stark contrast to that obtained in the [6π+2π] cycloaddition with alkenes in which only electron-deficient alkenes participate in the cycloaddition process.^{8a} It is also noteworthy that **8e** isomerizes during the reaction to give **10** (ca. 4%). The yield of **8e** could be dramatically improved by using 5 equivalents of allene instead of the usual 2 equivalents (entry 6). Irradiation of **1** with protected buta-2,3-

dienol (**7f**) (procedure A or B) yielded a separable 40:60 mixture of regioisomers **8f** and *endo*-**9f** in 38% overall yield (entry 7). In contrast to the cycloadditions with arylallenes or allenic ester **7e**, there was a slight preference for reaction at the internal double bond of the allene. This might be explained by a prior coordination of the oxygen atom of the ether group to the metal center, which would favor attack at the substituted double bond. A similar behavior was observed when **7g** was examined (entry 8). Cycloaddition of **1** with methoxyallene (**7h**) (procedure B) took place preferentially at the internal and electron-rich double bond of the allene, to give a 25:75 mixture of **8h** and *endo*-**9h** in a 51% overall yield (entry 9). When the bulkier alkoxyallene **7i** was engaged, the regioselectivity of the cycloaddition was found to reverse relative to the cycloaddition with **7h**, giving mixtures of **8i** and *endo*-**9i** in 60–65:40–35 ratios (entries 10–12). This reversal might be due to the presence of the bulky TBS group, which would disfavor approach of the allene from the internal double bond. An optimum overall reaction yield of 64% was obtained when reaction was performed at 0 °C using procedure B (entry 12). It is also noteworthy that the *Z/E* selectivity for **8**-type adducts was the same when 1-alkoxypropa-1,2-dienes **7h,i** or 4-alkoxybuta-1,2-dienes **7f,g** were employed (substituent preferentially *syn* to C-7) whereas it was reversed when electron-deficient allene **7e** was used (substituent preferentially *anti* to C-7).

In the reactions that gave cycloadducts **8i** and **9i**, a new adduct identified as the tetracyclic **11** was also isolated as a single diastereomer (5–8%, Table 3, entries 10–12). The structure of **11** was unambiguously established by analysis of 1D and 2D NMR data (¹H, ¹³C, Dept, COSY, HMQC, Homodecoupling, INADEQUATE and NOE sequences). A final confirmation of the structural and stereochemical assignment was provided by an X-ray crystallographic analysis performed on its diol derivative (see Supporting Information). This structure appears to be novel and might arise from a second Cr(0)-mediated [6 π +2 π] cycloaddition between cycloadduct **9i** and allene **7i** (Scheme 2). Such a tandem cycloaddition process has precedent in metal-mediated cycloaddition.¹¹ Work is currently in progress to identify conditions that would favor the formation of this tetracyclic core and to obtain a better understanding of this unusual transformation.

Finally, to evaluate asymmetric induction in the [6+2] cycloaddition, a preliminary investigation was performed with a chiral, non-racemic allene.

Chiral propargylic alcohol (*R*)-**13** was prepared in 31% yield and 88% $\pm$ 5% ee¹² via enantioselective addition of terminal alkyne **12** to benzaldehyde using a procedure described by Carreira¹³ (Scheme 3). Compound (*R*)-**13** was then converted to allene (*S*)-**4a** (33%, 60% $\pm$ 5% ee) in a two-step one-pot process.¹⁴ The enantiomeric excess was determined by ¹H NMR using a chiral shift reagent (Ag(fod)-Yb(tfc)₃).¹⁵

Irradiation of **1** with (*S*)-**4a** using procedure A afforded cycloadduct (*E*)-**5a** (45%). Subsequent removal of the protecting TBS group gave the optically active alcohol **5b** in 54% $\pm$ 5% ee,¹² demonstrating a good transfer of the chiral information from the allene (Scheme 4).

In summary, we have validated the first Cr(0)-mediated [6 π +2 π] cycloaddition of allenes with cycloheptatriene under photochemical activation, and demonstrated its potential by constructing an enantioenriched bicyclo[4.2.1]nonane framework starting from a chiral allene. Regiochemistry and *E/Z* selectivity in this cycloaddition were found to be subtly dependent on competing steric and electronic effects of the allene substituents. Further studies of this cycloaddition and investigation of synthetic applications are currently underway.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgment

The authors would like to thank the National Institutes of Health (GM-30771) for their generous financial support of this research. We are grateful to Dr. B. Ksebati for performing the NOE, homodecoupling and INADEQUATE experiments.

References

1. (a) Lautens M, Klute W, Tam W. Chem. Rev 1996;96:49–92. [PubMed: 11848744] (b) Früchauf HW. Chem. Rev 1997;97:523–596. [PubMed: 11848882]
2. For reviews of Cr(0)-mediated $[6\pi+4\pi]$ and $[6\pi+2\pi]$ cycloaddition reactions, see: (a) Rigby JH. Acc. Chem. Res 1993;26:579–585.585 (b) Rigby JH, Liebeskind LS. Advances in Metal-Organic Chemistry 1995;Vol. 4GreenwichJAI Press:89–127.127 (c) Rigby JH. Tetrahedron 1999;55:4521–4538.4538
3. For synthetic applications of Cr(0)-mediated $[6\pi+4\pi]$ and $[6\pi+2\pi]$ cycloadditions, see: (a) Rigby JH, Harmata M. Advances in Cycloaddition 1999;Vol. 6GreenwichJAI Press:97–118.118 (b) Rigby JH, Meyer JH. Synlett 1999;(Eschenmoser Issue):860–862.862 (c) Rigby JH, Warshakoon NC, Payen AJ. J. Am. Chem. Soc 1999;121:8237–8245.8245 (d) Rigby JH, Niyaz NM. Tetrahedron 2001;57:5091–5095.5095 (e) Rigby JH, Niyaz NM, Bazin B. Tetrahedron 2002;58:4879–4885.4885 (f) Rigby JH, Laxmisha MS, Hudson AR, Heap CH, Heeg MJ. J. Org. Chem 2004;69:6751–6760.6760 [PubMed: 15387599]
4. Rigby JH, Ahmed G, Ferguson MD. Tetrahedron Lett 1993;34:5397–5400.
5. For an overview of cycloadditions of allenes, see: Murakami M, Matsuda T. Krause N, Hashmi ASK. Modern Allene Chemistry 2004Wiley-VCHWeinheim:727–815.815
6. (a) Gompper R, Wolf U. Liebigs Ann. Chem 1979:1388–1405. (b) Hayakawa K, Nishiyama H, Kanematsu K. J. Org. Chem 1985;50:512–517.
7. (a) Gandhi RP, Ishar MPS. Chem. Lett 1989:101–104. (b) Ishar MPS, Gandhi RP. Tetrahedron 1993;49:6729–6740.
8. (a) Rigby JH, Henshilwood JA. J. Am. Chem. Soc 1991;113:5122–5123. (b) Rigby JH, Ateeq HS, Charles NR, Henshilwood JA, Short KM, Sugathapala PM. Tetrahedron 1993;49:5495–5506.
9. When concerned, the relative configuration of the stereocenter at the C-7 position was established by ^1H NMR NOE difference experiments, and in each case studied, only the *endo*-cycloadduct was obtained. See Supporting Information for additional discussion of structural assignments.
10. All the E and Z geometries discussed in this study were supported by ^1H NMR NOE correlations. See Supporting Information for some discussed examples.
11. Related Cr(0)-mediated tandem $[6\pi+2\pi]$, homo $[6\pi+2\pi]$ cycloadditions of alkynes with cyclic trienes are reported in the literature: (a) Rigby JH, Warshakoon NC, Heeg MJ. J. Am. Chem. Soc 1996;118:6094–6095.6095 (b) Chen W, Chaffee K, Chung HJ, Sheridan JB. J. Am. Chem. Soc 1996;118:9980–9981.9981 (c) Rigby JH, Heap CR, Warshakoon NC, Heeg MJ. Org. Lett 1999;1:507–508.508 [PubMed: 10822590] (d) Rigby JH, Heap CR, Warshakoon NS. Tetrahedron 2000;56:2305–2311.2311 Ref 3f.
12. ee was determined by ^1H NMR after conversion to the Mosher ester (See Supporting Information).
13. Boyall D, Frantz DE, Carreira EM. Org. Lett 2002;4:2605–2606. [PubMed: 12123387] and references cited therein.
14. Konoike T, Araki Y. Tetrahedron Lett 1992;33:5093–5096.
15. Mannschreck A, Munninger W, Burgemeister T, Gore J, Cazes B. Tetrahedron 1986;42:399–408. See Supporting Information.

Scheme 1.

Scheme 2.

Scheme 3.

Scheme 4.

Table 1

Photoinitiated $[6\pi+2\pi]$ cycloaddition reaction of **1** with 1,1-disubstituted allenes **2a-c**

entry	allene	R ¹	R ²	procedure ^a	product	Yield (%) ^b
1	2a	Ph	H	A	3a	50
2	2b	(CH ₂) ₅	H	B	3b	51
3	2c	(CH ₂) ₅	CH ₂ OTBS	B	3c	35

^aProcedure A: a mixture of **1** (1 equiv) and allene (2 equiv) in hexanes was irradiated at rt in a pyrex tube for periods of 4–6 h. Procedure B: to a solution of **1** in hexanes in a pyrex tube was introduced under irradiation at rt slowly a solution of allene (2 equiv) in hexanes via a syringe pump, then additional irradiation was continued for 4–6 h. See Supporting Information for more details.

^bYield of isolated product after purification by flash chromatography.

Table 2
Photoinitiated $[6\pi+2\pi]$ cycloaddition reaction of **1** with 1,3-disubstituted allenes **4a–c**

$1 + R^1-CH=CH-R^2 \xrightarrow[h\nu]{\text{pyrex}}$
4a $R^1 = Ph, R^2 = CH_2OTBS$
4b $R^1 = Ph, R^2 = CH_2OTHP$
4c $R^1 = Ph, R^2 = CH_3$

entry	allene	procedure ^a	product	yield(%) ^b	isomeric ratio ^c	
					5 (E/Z)	6 (E/Z)
1	4a	A	5a	59	100 (>99:<1)	0
2	4a	B	5a	50	100 (85:15 ^d)	0
3	4b	A	5b^e	56 ^f	100 (>99:<1)	0
4	4c	B	5c, 6c	64	88 (10:90)	12 (23:77)

^aSee footnote a of Table 1.

^bOverall yield (**5**+**6**) of isolated product after purification.

^cRatios (**5**/**6**, E/Z) determined by GC analysis on the crude product before purification.

^dSometimes E/Z ~ 98:2.

^eIsolated as an alcohol after removal of THP group.

^fYield over two steps after removal of THP group.

Table 3
Photoinitiated $[6\pi+2\pi]$ cycloaddition reaction of **1** with monosubstituted allenes **7a-i**

entry	R	procedure ^a	product	yield (%) ^b	isomeric ratio ^c
1	Ph, 7a	B	8a, 9a	50	> 97 (45:55)
2	p-MeOC ₆ H ₄ , 7b	B	8b, 9b	42	> 97 (48:52)
3	p-EtO ₂ CC ₆ H ₄ , 7c	B	8c, 9c	23	> 97 (48:52)
4	p-O ₂ NC ₆ H ₄ , 7d	B	8d, 9d	56	> 97 (40:60)
5	CO ₂ Et, 7e	A or B	8e	34 ^d	100 (65:35)
6	CO ₂ Et, 7e	A ^e or B ^e	8e	56 ^d	100 (60:40)
7	CH ₂ OTBS, 7f	A or B	8f, 9f	38	40 (15:85)
8	CH ₂ OTHP, 7g	A or B	8g^f, 9g^f	41 ^g	40 (35:65)
9	OMe, 7h	B	8h, 9h	51	25 (35:65)
10	OTBS, 7i	A	8i, 9i	23 ^h	65 (45:55)
11	OTBS, 7i	B	8i, 9i	50 ^h	65 (40:60)
12	OTBS, 7i	B ⁱ	8i, 9i	64 ^h	60 (40:60)

^a See footnote a of Table 1.

^b Overall yield (**8-9**) of isolated product after flash chromatography.

^c Ratios (**8-9**, Z/E) determined by GC analysis on the crude product before purification or by ¹H NMR.

^d Adduct **10** (ca. 4% yield) was also formed.

^e 5 equiv of allene were used.

^f Isolated as an alcohol after removal of THP group.

^g Overall yield over two steps after removal of THP group.

^h Tetracycle **11** (5-8% yield) was also isolated.

The reaction mixture was cooled down at 0 °C during irradiation, and additional irradiation was continued for 1 h after addition of allene.
i