Melanotic neuroectodermal tumour of infancy

Scott Hamilton MD¹, Duncan MacRae MD¹, Sumit Agrawal MD², Damir Matic MD³

S Hamilton, D MacRae, S Agrawal, D Matic. Melanotic neuroectodermal tumour of infancy. *Can J Plast Surg* 2008;16(1):41-44.

Melanotic neuroectodermal tumour of infancy is an extremely rare neoplasm arising in newborns and young children, typically involving the face or cranium. A case arising from the maxilla, requiring extensive resection with a near-total maxillectomy, is presented. A thorough review of the literature on this unusual tumour is provided, with emphasis on prognostic factors and appropriate treatment.

Key Words: *Maxillectomy; Melanotic; MNTI; Pediatric; Pigmented*

Melanotic neuroectodermal tumour of infancy (MNTI) is a rare, pigmented neoplasm generally arising in infants during the first year of life (1-3). This benign tumour originates from the soft tissue overlying the maxilla in more than 70% of cases, although mandibular, cranial, cerebral and genital involvement have also been described (4,5). The mean age of patients at diagnosis is 4.3 months, with a near-equal male to female ratio of 6:7 (6-8). The present report describes a case of MNTI requiring a near-total maxillectomy, along with an overview of the current pathological knowledge and therapeutic recommendations.

CASE PRESENTATION

A two-month-old male presented with a one-month history of a smooth, firm, rapidly growing mass involving the superior maxillary alveolar ridge. There was no history of airway or feeding difficulties. Maternal, birth and family history were all within normal limits. On physical examination, there was a dark mass involving the premaxilla that was nontender and nonpulsatile (Figure 1).

Both computed tomography and magnetic resonance imaging (MRI) scans revealed an extensive mass filling the maxilla, crossing the midline and involving the orbital floor (Figures 2A and 2B). A fine-needle aspiration biopsy was performed and revealed a dual population of small neuroblastic cells and large melanin-containing epithelial cells. A provisional diagnosis of MNTI was made and confirmed on final histology (Figure 3) following an incisional biopsy.

The child underwent a complete resection of the mass using a modified Weber-Ferguson approach, with the nasal floor, septum and two-thirds of the alveolar ridge resected (Figure 4). The orbital floor was also removed, although the periorbital was preserved. A 5 mm margin was included around the palpable

Tumeur mélanotique neuroectodermique de l'enfance

La tumeur mélanotique neuroectodermique de l'enfance, qui touche surtout la face et le crâne, est extrêmement rare chez les nouveau-nés et les jeunes enfants. Il sera question, dans l'article, d'un cas de tumeur du maxillaire, qui a nécessité une résection étendue, voire une maxillectomie quasi totale. Un examen exhaustif de la documentation sur ce type peu fréquent de tumeur, plus particulièrement sur les facteurs pronostiques et le traitement approprié, complètera l'étude du sujet.

Figure 1) A two-month-old male with a tumour arising from the maxilla

tumour. Nasal stents were placed to prevent stenosis (Figures 5A and 5B). A palatal appliance impression was taken intraoperatively; however, it was not used because of difficulty in retaining the obturator due to early granulation.

The child was followed every three months for the first year, then annually afterwards, with an MRI obtained each year. There is no evidence of recurrence at three years following resection. Visual acuity is normal without strabismus or diplopia.

¹Department of Otolaryngology, University of Western Ontario, London, Ontario; ²Department of Otolaryngology, Stanford University, Stanford, California, USA; ³Division of Plastic Surgery, University of Western Ontario, London, Ontario

Correspondence: Dr Damir Matic, University of Western Ontario, Childrens Hospital of Western Ontario, 800 Commissioners Road East, Room E1-332, London, Ontario N6A 5W9. Telephone 519-685-8557, fax 519-685-8405, e-mail damir.matic@lhsc.on.ca

Figure 2) A Preoperative three-dimensional computed tomography scan showing an extensive tumour involving the maxilla and orbital floor. B Sagittal magnetic resonance imaging demonstrating depth of tumour invasion. Arrow shows anterior extent of tumour

Figure 3) Histology demonstrating both small neuroblastic cells and large melanin-containing epithelial cells

A class I skeletal relationship exists. The palate has completely mucosalized except for a 5 mm × 10 mm anterior fistula (Figure 6). Mild signs of velopharyngeal insufficiency are

Figure 4) Intraoperative photograph demonstrating the Weber-Ferguson approach

present and were thought to be secondary to the fistula, until further evaluation revealed a submucous cleft. Dental rehabilitation and closure of the fistula is planned near the time of permanent dental eruption, with possible surgical correction of the cleft should symptoms persist.

DISCUSSION AND LITERATURE REVIEW

This is the first reported case in which a patient was diagnosed both with an MNTI involving the maxilla and a submucous cleft. The concomitant presence of the submucous cleft together with a postoperative palatal fistula complicated the diagnosis and treatment of his velopharyngeal insufficiency. This will require further evaluation before definitive management.

Since first described in 1918, there have been approximately 250 cases of MNTI reported in the world literature (9). The variety of other names assigned to the tumour, including melanocarcinoma, melanotic epithelial odontoma and melanotic prognoma, reflects its confusing histogenesis (10). Patients typically present with a rapidly growing, nonulcerated mass, affecting the craniofacial region in 90% of cases (11,12). Although melanin is produced by the tumour, pigmentation may not be clinically evident (13,14). Local invasion may be accompanied by bony destruction, tooth displacement and feeding difficulties (6,15,16).

Despite this locally aggressive behaviour, MNTI is generally classified as a benign tumour (11,17). A review by Cutler et al (18) demonstrated malignant features in only 1.9% of MNTI cases, with estimates from more recent publications as high as 6% (17,19). Unfortunately, it is difficult to determine the potential for malignancy or recurrence based on clinical assessment, imaging or histopathology (20).

Computed tomography imaging typically reveals a well-demarcated, hyperdense lesion with contrast enhancement and hyperostosis of adjacent bone (21-24). On MRI scans, these lesions tend to be hyperintense on T1 and hypointense on T2 (21,22,24). Despite these characteristic findings, imaging will seldom be diagnostic and tissue biopsy is therefore required.

The histological profile of MNTI demonstrates a biphasic cell population within a stroma of moderately vascularized fibrous tissue (1). Large, epithelioid, melanin-containing cells are arranged in alveolar or tubular formations around clusters

Figure 5) A Nasal stents placed. **B** Immediately postoperatively

of small neuroblastic cells (15,25,26). A differential diagnosis would include other small round cell tumours occurring in infancy, such as rhabdomyosarcoma, neuroblastoma, melanoma and lymphoma. The diagnosis can be especially difficult, given the rarity of MNTI.

Immunohistochemical stains can greatly assist in establishing the diagnosis. The larger, epithelioid cells stain positive most frequently with cytokeratin, epithelial membrane antigen, vimentin and HMB-45, reflecting epithelial and melanocytic differentiation (3,27). These larger cells are usually nonreactive with S-100 protein, aiding in differentiation from tumours such as melanoma. (14,28). The nests of smaller cells in MNTI are often positive for neurogenic markers such as

Figure 6) Patient three years postoperatively with no evidence of tumour recurrence

synaptophysin, neuron-specific enolase and glial fibrillar acidic protein (15,25,27). The variability seen in staining results likely represents the multiphenotypical character of this tumour (1).

Ultrastructural studies with electron microscopy have been used alongside histochemical staining to support the theory that MNTIs belong to a group of neuroectodermal tumours arising from neural crest origin (17). Tumours in this group also include neuroblastoma, Wilm's tumour, Ewing sarcoma and primitive neuroectodermal tumour. Elevated urinary vanillyl-mandelic acid and the ability to differentiate primitive neural crest cells to neuroblastic and melanocytic lineages provide additional evidence (29-32). Increased alpha-fetoprotein levels and positive staining for the c-myc antigen, characteristic of other neural crest tumours, has also been described (5,27,29,33,34). Despite these microscopic and immunohistochemical similarities, there are no shared genetic changes linking MNTI with the other neuroectodermal tumours (14,35).

Surgical excision remains the mainstay of treatment for patients with MNTI, as the only modality with proven efficacy (2,15). Given the benign nature of most tumours, there has been some controversy over the optimal surgical management and necessity of adequate margins. Views range from conservative enucleation, to curettage of adjacent tissue, to recommended margins of up to 5 mm (2,18,20,36,37). Efforts should be directed toward the preservation of vital structures and organs (38). Lesion pigmentation and preoperative imaging often assist in delineating tumour margins (14). Adjuvant chemotherapy and radiation are ineffective in controlling recurrences where total excision has not been achieved, and their role is therefore extremely limited (39,40). Authors have advocated their role in metastatic disease and in cases of recurrence after a second resection, although there remains little evidence of added benefit (38,39).

The local recurrence rate following excision ranges from 10% to 15%, although it has been reported as high as 45% (11,12,25,41,42). Shaia et al (38) examined the risk of recurrence over the past 20 years and found that it did not differ from historical data since the tumour was first diagnosed 90 years previously. More than 80% of recurrences occur in the

first four months, (25) with the majority of recurrent tumours occurring in patients initially diagnosed before 12 weeks of age (38). Most patients can achieve cure with repeated surgery (16).

Metastatic spread is rare, occurring in only 3% of patients (5). When metastases develop, the smaller neuroblastic cells predominate in the secondary deposits (14,41,43,44). Histology therefore resembles neuroblastoma more than MNTI (14).

In recent years, a great deal of scientific effort has been put into identifying prognostic markers for MNTI, with little success. Histology has been proven to be an unreliable guide to clinical outcome (15). Flow cytometry has been attempted, again with little success (25,42). Promise may lie with staining

for CD99 and Ki-67, both of which have been shown to be associated with aggressive tumours in single studies (15,45). The best indicator of prognosis continues to be close postoperative follow-up with frequent imaging.

CONCLUSIONS

MNTI is a rare pediatric tumour of neural crest origin. The vast majority of tumours occur in the head and neck, predominantly with maxillary involvement. As in our case, imaging, histology and immunohistochemical staining contribute to making a challenging diagnosis. The tumours are treated surgically, with little benefit from adjuvant treatments. Careful follow-up is required given high recurrence rates.

REFERENCES

- Kaya S, Unal OF, Sarac S, Gedikoglu G. Melanotic neuroectodermal tumor of infancy: Report of two cases and review of literature. *Int J Pediatr Otorhinolaryngol* 2000;52:169-72.
- Franchi G, Sleilati F, Soupre V, et al. Melanotic neuroectodermal tumor of infancy involving the orbit and maxilla: Surgical management and follow-up strategy. *Br J Plast Surg* 2002;55:526-9.
- Adad SJ, Pinheiro SW, Marinho EO, Rua AM, Rodrigues DB. Melanotic neuroectodermal tumor of infancy (MNTI). *Diagn Cytopathol* 2004;30:67-9.
- Morin S, Bergevin M, Derkay C. Melanotic neuroectodermal tumor of infancy (MNTI). *Arch Otolaryngol Head Neck Surg* 1992;118:664-7.
- Carnevale GG, Mortelliti AJ. The operating microscope in the management of melanotic neuroectodermal tumor of infancy. *Am J Otolaryngol* 2001;22:76-9.
- Puchalski R, Shah UK, Carpentieri D, McLaughlin R, Handler SD. Melanotic neuroectodermal tumor of infancy (MNTI) of the hard palate: Presentation and management. *Int J Pediatr Otorhinolaryngol* 2000;53:163-8.
- Antunes AC, Freitas RM, Oliveira PP, Reboucas RG. Melanotic neuroectodermal tumor of infancy: Case report. *Arq Neuropsiquiatr* 2005;63:670-2.
- Tan O, Atik B, Ugras S. Melanotic neuroectodermal tumor in a newborn. *Int J Pediatr Otorhinolaryngol* 2005;69:1441-4.
- Rustemeyer J, Thieme V, Loeschke S, Bremerish A, Kossling FK. Melanotic neuroectodermal tumor of infancy. *Klin Pediatr* 2001;213:69-73.
- Agarwal P, Agarwal V, Raina VK. Melanotic neuroectodermal tumor of infancy: Case report of an unusual tumor. *Indian J Plastic Surg* 2003;36:123-5.
- Eckardt A, Swennen G, Teltzow T. Melanotic neuroectodermal tumor of infancy involving the mandible: 7-year follow-up after hemimandibulectomy and costochondral graft reconstruction. *J Craniofac Surg* 2001;12:349-54.
- Al-Marzooq YM, Al-Bagshi MH, Chopra R, Hashish H. Melanotic neuroectodermal tumor of infancy in the soft tissues of the arm: Fine needle aspiration biopsy and histologic correlation-a case report. *Diagn Cytopathol* 2003;29:352-5.
- Ijiri R, Onuma K, Ikeda M, et al. Pigmented intraosseous odontogenic carcinoma of the maxilla: A pediatric case report and differential diagnosis. *Hum Pathol* 2001;32:880-4.
- Gaiger de Oliveira M, Thompson LD, Chaves AC, Rados PV, da Silva Lauxen I, Filho MS. Management of melanotic neuroectodermal tumor of infancy. *Ann Diagn Pathol* 2004;8:207-12.
- Barrett AW, Morgan M, Ramsay AD, Farthing PM, Newman L, Speight PM. A clinicopathologic and immunohistochemical analysis of melanotic neuroectodermal tumor of infancy. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod* 2002;93:688-98.
- Liu HH, Chen TW, Chang HS. Melanotic neuroectodermal tumor of infancy in the maxilla: a case report. *Int J Paediatr Dent* 2004;14:371-5.
- Haider N, Mc Dermott M, Fitzgerald RJ. Melanotic neuroectodermal tumor of infancy arising in skull. *Med Pediatr Oncol* 2003;41:495-6.
- Cutler LS, Chaudhry AP, Topazian R. Melanotic neuroectodermal tumor of infancy: An ultrastructural study, literature review, and reevaluation. *Cancer* 1981;48:257-70.
- Gupta A, Trehan A, Marwaha RK, Sharma RK, Nijhawan R. Melanotic neuroectodermal tumor in an infant. *Indian J Pediatr* 2002;69:725-6.
- Mast BA, Kapadia SB, Yunis E, Bentz M. Subtotal maxillectomy for melanotic neuroectodermal tumor of infancy. *Plast Reconstr Surg* 1999;103:1961-3.
- Atkinson GO Jr, Davis PC, Patrick LE, Winn KJ, Ball TI, Wylly JB. Melanotic neuroectodermal tumor of infancy. MR findings and a review of the literature. *Pediatr Radiol* 1989;20:20-2.
- Nitta T, Endo T, Tsunoda A, Kadota Y, Matsumoto T, Sato K. Melanotic neuroectodermal tumor of infancy: A molecular approach to diagnosis. *J Neurosurg* 1995;83:145-8.
- George JC, Edwards MK, Jakacki RI, Kho-Duffin J. Melanotic neuroectodermal tumor of infancy. *AJNR Am J Neuroradiol* 1995;16:1273-5.
- Nishio S, Morioka T, Murakami N, Fukui M, Inamitsu T, Ishihara S. Melanotic neuroectodermal tumor of infancy at the anterior fontanelle. *Neuroradiology* 1999;41:202-4.
- Kapadia SB, Frisman DM, Hitchcock CL, Ellis GL, Popek EJ. Melanotic neuroectodermal tumor of infancy. Clinicopathological, immunohistochemical, and flow cytometric study. *Am J Surg Pathol* 1993;17:566-73.
- Galera-Ruiz H, Gomez-Angel D, Vazquez-Ramirez FJ, Sanguino-Fabre JC, Salazar-Fernandez CI, Gonzalez-Hachero J. Fine needle aspiration in the pre-operative diagnosis of melanotic neuroectodermal tumor of infancy. *J Laryngol Otol* 1999;113:581-4.
- Metwally H, Cheng J, Maruyama S, et al. Establishment and characterization of new cell lines derived from melanotic neuroectodermal tumor of infancy arising in the mandible. *Pathol Int* 2005;55:331-42.
- Rickert CH, Probst-Cousin S, Blasius S, Gullotta F. Melanotic progenoma of the brain: a case report and review. *Childs Nerv Syst* 1998;14:389-93.
- Anil S, Raji MS, Beeana VT. Melanotic neuroectodermal tumor of infancy. *Indian J Dent Res* 1993;4:65-7.
- Lahav R, Ziller C, Dupin E, Le Douarin NM. Endothelin 3 promotes neural crest cell proliferation and mediates a vast increase in melanocyte number in culture. *Proc Natl Acad Sci USA* 1996;93:3892-7.
- Stone JG, Spirling LI, Richardson MK. The neural cell population responding to endothelin-3 in vitro includes multipotent cells. *J Cell Sci* 1997;110:1673-82.
- Watabe H, Soma Y, Kawa Y, et al. Differentiation of murine melanocyte precursors induced by 1,25-dihydroxyvitamin D3 is associated with the stimulation of endothelin B receptor expression. *J Invest Dermatol* 2002;119:583-9.
- Moriuchi S, Shimizu K, Miyao Y, Hayakawa T. A novel PNET cell line with melanotic differentiation. *Anticancer Res* 1996;16:779-84.
- Kim YG, Oh JH, Lee SC, Ryu DM. Melanotic neuroectodermal tumor of infancy. *J Oral Maxillofac Surg* 1996;54:517-20.
- Khoddami M, Squire J, Zielenska M, Thorner P. Melanotic neuroectodermal tumor of infancy: A molecular genetic study. *Pediatr Dev Pathol* 1998;1:295-9.
- Irving RM, Parikh A, Coumbe A, Albert DM. Melanotic neuroectodermal tumor of infancy. *J Laryngol Otol* 1993;107:1045-8.
- Dashti SR, Cohen ML, Cohen AR. Role of radical surgery for intracranial melanotic neuroectodermal tumor of infancy: case report. *Neurosurgery* 1999;45:175-8.
- Shaia WT, Dinardo LJ, Underhill TE, Cesca CE. Recurrent melanotic neuroectodermal tumor of infancy. *Am J Otolaryngol* 2002;23:249-52.
- Mirich DR, Blaser SI, Harwood-Nash DC, Armstrong DC, Becker LE, Posnick JC. Melanotic neuroectodermal tumor of infancy: Clinical, radiologic, and pathologic findings in five cases. *AJNR Am J Neuroradiol* 1991;12:689-97.
- Woessmann W, Neugebauer M, Gossen R, Blutters-Sawatzki R, Reiter A. Successful chemotherapy for melanotic neuroectodermal tumor of infancy in a baby. *Med Pediatr Oncol* 2003;40:198-9.
- Johnson RE, Scheithauer BW, Dahlin DC. Melanotic neuroectodermal tumor of infancy. A review of seven cases. *Cancer* 1983;52:661-6.
- Pettinato G, Manivel JC, d'Amore ES, Jaszcz W, Gorlin RJ. Melanotic neuroectodermal tumor of infancy. A reexamination of a histogenetic problem based on immunohistochemical, flow cytometric, and ultrastructural study of 10 cases. *Am J Surg Pathol* 1991;15:233-45.
- Block JC, Waite DE, Dehner LP, Leonard AS, Ogle RG, Gatto DJ. Pigmented neuroectodermal tumor of infancy. An example of rarely expressed malignant behavior. *Oral Surg Oral Med Oral Pathol* 1980;49:279-85.
- Navas Palacios JJ. Malignant melanotic neuroectodermal tumor: Light and electron microscopic study. *Cancer* 1980;46:529-36.
- van Oijen MG, Medema RH, Slootweg PJ, Rijksen G. Positivity of the proliferation marker Ki-67 in noncycling cells. *Am J Clin Pathol* 1998;110:24-31.