Published in final edited form as:

J Am Chem Soc. 2009 May 13; 131(18): 6332–6333. doi:10.1021/ja901313v.

Biosynthesis of the Sesquiterpene Antibiotic Albaflavenone in *Streptomyces coelicolor*. Mechanism and Stereochemistry of the Enzymatic Formation of Epi-isozizaene

Xin Lin and David E. Cane*

Department of Chemistry, Box H, Brown University, Providence, Rhode Island 02912-9108

The tricyclic sesquiterpene albaflavenone (**1**), which has been isolated from the gram-positive soil bacteria *Streptomyces coelicolor* and *S. albidoflavus*, has an earthy, camphor-like odor and exhibits antibacterial activity.^{1,2} The biosynthesis of albaflavenone in *S. coelicolor* A3(2) is under the control of a two-gene cluster, *sco5222* and *sco5223*.^{1a} The 1086-bp *sco5222* gene encodes a protein of 361 amino acids that catalyzes the cyclization of farnesyl diphosphate (**2**, FPP) to the unusual sesquiterpene hydrocarbon (+)-epi-isozizaene (**3**).^{1b} The downstream *sco5223* gene, which shares a four-nucleotide ATGA translational overlap with *sco5222*, corresponds to a cytochrome P450 that catalyzes the two-step allylic oxidation of epi-isozizaene to albaflavenone (**1**).^{1a}

The cyclization of FPP to epi-isozizaene catalyzed by the SCO5222 protein has been proposed to involve the initial ionization and isomerization of FPP (**2**) to the tertiary allylic isomer, (3*R*)-nerolidyl diphosphate (**4**, NPP) (Scheme 2).^{1b} Rotation about the newly generated C-2-C-3 single bond would generate the corresponding *cisoid* NPP conformer that can undergo ionization and cyclization to generate the intermediate bisabolyl cation **5**. Following 1,2-hydride shift and spirocyclization, the resultant acorenyl cation **6** can undergo further cyclization, ring contraction, 1,2-methyl migration, and final deprotonation to yield (+)-epi-isozizaene (**3**). This proposed mechanism has been supported by incubation of recombinant epi-isozizaene synthase with individual samples of deuterated FPP, [1,1-²H₂]FPP (**2a**), (1*R*)-[1-²H]FPP (**2b**), and (1*S*)-[1-²H]FPP (**2c**) followed by NMR analysis of the resultant labeled **3**.^{1b} The combined labeling results are completely consistent with the proposed stereochemical model (Scheme 2), confirming the predicted net retention of configuration in the displacement of the diphosphate group of FPP that reflects the intermediacy of NPP. We now report the direct demonstration of the role of (3*R*)-NPP (**4**) in the formation of epi-isozizaene, including the determination of the detailed stereochemistry of the isomerization-cyclization-rearrangement reaction cascade. We also report the results of site-directed mutagenesis of the universally conserved Mg²⁺-binding domains and the isolation of abortive cyclization products that reinforce the proposed cyclization mechanism.

Incubation of (3*RS*)-(Z)-[1-²H]NPP ((±)-**4b**)³ with purified recombinant epi-isozizaene synthase resulted in the formation of [11-²H]epi-isozizaene, *m/z* 205, as determined by GC-MS analysis.⁴ To establish the absolute configuration of the natural NPP intermediate, we carried out competitive incubations between deuterated and undeuterated samples of both (3*S*)- and (3*R*)-(Z)-NPP,³ using selected ion monitoring GC-MS to measure the deuterium content of the resulting epi-isozizaene. Thus incubation of a 1:2 mixture of (3*S*)-(Z)-[1-²H]

E-mail: E-mail: David_Cane@brown.edu.

Supporting Information Available: Experimental procedures, GC-MS and NMR data, and supplemental Figures and Tables. This material is available free of charge via the Internet at <http://pubs.acs.org>.

NPP and unlabeled (3*RS*)-NPP gave epi-isozizaene that was devoid of deuterium (>95% d₀). Had only the (3*S*)-NPP been utilized, the observed d₀:d₁ ratio of **3** would have been 1.0, while equal conversion of both enantiomers of NPP would have yielded a d₀:d₁ ratio in the product of 0.5. Complementary incubation of a 1:2 mixture of unlabeled (3*S*)-NPP and (3*RS*)-(Z)-[1-²H]NPP with epi-isozizaene synthase gave [²H]epi-isozizaene whose deuterium content indicated a >93:7 preference for (3*R*)-NPP over (3*S*)-NPP. To determine the stereochemistry of the cyclization of the (3*R*)-NPP, we carried out preparative-scale incubations with (3*RS*)-(Z)-[1-²H]NPP. The resulting [11-*anti*-²H]epi-isozizaene (**3b**) lacked the characteristic ¹H NMR signal at δ 1.47 previously assigned^{1b} to H-11-*anti* while displaying a broad singlet for H-11-*syn* that was shifted 0.03 ppm upfield to δ 1.36 due to the isotope effect of the geminal D-11-*anti*. Similarly, the H-11-*anti*/C-11 cross-peak was absent from the HSQC spectrum, while a clear cross-peak for H-11-*syn*/C-11 was evident at δ 1.36 and 36.5 ppm.

These results are completely consistent with the stereochemical model illustrated in Scheme 2, confirming the predicted intermediacy of (3*R*)-NPP (**4**) which is generated by *syn* allylic isomerization of FPP (**2**) and then undergoes *anti*-S_N' cyclization to the bisabolyll cation. This sequence of *syn* isomerization/*anti*-S_N' cyclization to form the bisabolyll cation, with its necessarily *cis* ring double bond, is typical of terpene synthases.^{3,5}

To further test the mechanism of Scheme 2, individual samples of [12,12,12-²H₃]FPP (**2d**)⁶ and [13,13,13-²H₃]FPP (**2e**)⁶ were each incubated with recombinant epi-isozizaene synthase. The labeling of the resulting epi-isozizaene was assigned by ¹H and ²H NMR. Thus [14,14,14-²H₃]epi-isozizaene (**3d**), derived from [12,12,12-²H₃]FPP (**2d**), lacked the characteristic allylic methyl singlet for H-14 at δ 1.42, while showing the corresponding peak for D-14 in the ²H NMR spectrum. By contrast, cyclization of [13,13,13-²H₃]FPP (**2e**) gave [12,12,12-²H₃]epi-isozizaene (**3e**) with only a minor singlet at δ 0.99 (<10%), corresponding to geminal methyl H-12, while displaying a singlet at δ 0.99 for D-12 in the ²H-NMR spectrum.⁶

It is therefore established that in the cyclization of the intermediate acorenyl cation **6**, electrophilic attack engages exclusively the *si* face of the 2-propyl cation side chain to give the postulated tricyclic intermediate cation **7** (Scheme 2).⁷ The ²H-labeling of the geminal methyl groups in **7** and **8** can be inferred from the mandatorily *suprafacial* 1,2-methyl migration. After ring contraction, intermediate **7** undergoes *syn*-1,2-methyl migration and *syn*-deprotonation of **8** to yield [12,12,12-²H₃]epi-isozizaene (**3e**).⁸

Detailed GC-MS analysis of the organic extracts resulting from the incubation of FPP with wild-type epi-isozizaene synthase revealed the formation of at least 6 additional sesquiterpene hydrocarbons **9-14**, *m/z* 204, which together comprised ~20% of the total product mixture, with the main components being β-farnesene (**9**, 5%) and zizaene (**10**, 10%) (Chart 1, Table S2). All but compound **14** were identified by comparison of their individual mass spectra and retention indices with those of authentic compounds in the MassFinder 3.0 Database.^{9,10} Each aberrant product results from premature quenching or derailment of the normal intermediates of the natural cyclization cascade (Scheme S1).

Epi-isozizaene synthase harbors two universally conserved Mg²⁺-binding domains typical of all sesquiterpene synthases, the apartate-rich 99**DDRHD** motif and the characteristic NSE triad ²⁴⁰**NDLCSLPKE**.¹¹ Consistent with the importance of each of these domains, the corresponding D99E, D100N, N240D, S244A, and E248D mutants of epi-isozizaene synthase each retained <5% of the FPP cyclase activity of the wild-type enzyme, while the D99N mutant was completely inactive. In common with the behavior of active site mutants of paralogous sesquiterpene synthases,¹² the five epi-isozizaene mutants with residual cyclase activity generated perturbed ratios of the aberrant FPP cyclization products **9-14**, as well as an

additional sesquiterpene, α -neocallitropsene (**15**) (Table S1).¹⁰ The modified product distributions are thought to reflect compromised binding of the divalent metal ions of the cofactor, with resultant changes in protein conformation and relaxed control over the conformation of the bound substrate FPP and derived intermediates.^{12,13}

A BLAST search using as query the *S. coelicolor* epi-isozizaene sequence (SCO5222; UniProt ID Q9K499) reveals likely orthologous terpene synthases in *S. sviveus* (B5HN36, 84% identity), *S. avermitilis* (Q82IV1, 82% identity), and *Streptomyces* sp. SPB74 (B5GD17, 57% identity), each of which has identical conserved Mg²⁺-binding domains. Moreover, all three epi-isozizaene synthase orthologs have 4-nt ATGA translational overlaps with downstream close homologs of CYP170A1 encoded by *sco5223*, indicating that the albaflavenone biosynthetic pathway may well be widely distributed and highly conserved among Streptomycetes.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

This work was supported by NIH grant GM30301 to DEC.

References and Notes

- (1). (a) Zhao B, Lin X, Lei L, Lamb DC, Kelly SL, Waterman MR, Cane DE. *J. Biol. Chem.* 2008;283:8183–8189. [PubMed: 18234666] (b) Lin X, Hopson R, Cane DE. *J. Am. Chem. Soc.* 2006;128:6022–6023. [PubMed: 16669656]
- (2). Scholler CE, Gurtler H, Pedersen R, Molin S, Wilkins K. *J. Agric. Food Chem.* 2002;50:2615–2621. [PubMed: 11958631] Gurtler H, Pedersen R, Anthoni U, Christophersen C, Nielsen PH, Wellington EMH, Pedersen C, Bock K. *J. Antibiot.* 1994;47:434–439. [PubMed: 8195043]
- (3). (a) Cane DE, Ha HJ. *J. Am. Chem. Soc.* 1988;110:6865–6870. Cane DE, Ha HJ. *J. Am. Chem. Soc.* 1986;108:3097–3099. (b) Cane DE, Tandon M. *J. Am. Chem. Soc.* 1995;117:5602–5603.
- (4). Incubation of (3*RS*)-NPP gave a~1:1 mixture of epi-isozizaene (**3**) and β -farnesene (**9**). Control experiments showed that >95% of this coproduct **9** was derived from the unnatural enantiomer (3*S*)-NPP, indicating that only (3*R*)-NPP can be cyclized by epi-isozizaene synthase.
- (5). Wise, ML.; Croteau, R. *Comprehensive Natural Products Chemistry. Isoprenoids Including Carotenoids and Steroids.* Cane, DE., editor. Vol. 2. Elsevier; Oxford: 1999. p. 97-153.
- (6). Cane DE, Prabhakaran PC, Oliver JS, McIlwaine DB. *J. Am. Chem. Soc.* 1990;112:3209–3210.3210. The sample of [13,13,13-²H₃]FPP (**2e**) contained ~10% of the isomeric [12,12,12-²H₃]FPP (**2d**).
- (7). The stereochemistry of the H-4 bridgehead-proton in **6**, **7** and **8** was deduced based on the assumption of least motion of the 2-propyl cation side chain during the formation of the new C-C bond in **7**.
- (8). For analogous sequential 1,2-methyl migration and *syn*-deprotonation, cf. aristolochene synthase (ref 3a) and epi-aristolochene synthase: Schenk DJ, Starks CM, Manna KR, Chappell J, Noel JP, Coates RM. *Arch. Biochem. Biophys.* 2006;448:31–44.44 [PubMed: 16309622].
- (9). Harangi J. *J. Chromatogr. A* 2003;993:187–195. [PubMed: 12735452]
- (10). The absolute configurations of **10-13** and **15** were provisionally assigned on the basis of the cyclization mechanism and the known absolute configuration of epi-isozizaene.
- (11). Christianson DW. *Chem. Rev.* 2006;106:3412–3442. [PubMed: 16895335]
- (12). Trichodiene synthase: Vedula LS, Zhao Y, Coates RM, Koyama T, Cane DE, Christianson DW. *Arch. Biochem. Biophys.* 2007;466:260–266.266 [PubMed: 17678871]. Pentalenene synthase: Seemann M, Zhai G, de Kraker JW, Paschall CM, Christianson DW, Cane DE. *J. Am. Chem. Soc.* 2002;124:7681–7689.7689 [PubMed: 12083921]. Aristolochene synthase: Felicetti B, Cane DE. *J. Am. Chem. Soc.* 2004;126:7212–7221.7221 [PubMed: 15186158].

- (13). Vedula LS, Rynkiewicz MJ, Pyun HJ, Coates RM, Cane DE, Christianson DW. *Biochemistry* 2005;44:6153–6163. [PubMed: 15835903] Rynkiewicz MJ, Cane DE, Christianson DW. *Biochemistry* 2002;41:1732–1741. [PubMed: 11827517]

Scheme 1.

Biosynthesis of Albaflavenone (1) from Farnesyl Diphosphate (2) and Epi-isozaene (3)

Scheme 2.

Mechanism and Stereochemistry of the Cyclization of FPP (**2**) to Epi-isozizaene (**3**) via $(3R)$ -Nerolidyl Diphosphate (**4**).

9 β -farnesene **12** sesquisabinene **13** sesquiphellandrene **10** zizaene **11** α -cedrene **15** α -neocallitropsene

Chart 1.
Additional sesquiterpenes produced by wild-type and mutant epi-isozizaene synthases.