Published in final edited form as:

Org Lett. 2009 March 19; 11(6): 1293–1295. doi:10.1021/ol9000685.

Pd(0)/InI-Mediated Allylic Additions to 4-Acetoxy-2-Azetidinone: New Route to Highly Functionalized Carbocyclic Scaffolds

Cara Cesario and Marvin J. Miller*

Department of Chemistry and Biochemistry, University of Notre Dame, 251 Nieuwland Science Hall,
Notre Dame, Indiana 46556

Abstract

Acylnitroso-derived hetero-Diels-Alder cycloadducts are susceptible to C–O bond cleavage with Pd (0) and InI to form allylic indium (III) species. The in situ prepared allylindium compounds readily react at room temperature with Eschenmoser's salt. Allylation of 4-acetoxy-2-azetidinone provides derivatized cyclopentenones in high regio- and diastereoselectivity.

Indium-mediated allylations have gained considerable attention due to their ability to achieve carbon-carbon bond formation under mild conditions with high stereo- and regioselectivities.¹ Although many indium-promoted allylations have been successful with diverse electrophilic substrates (including aldehydes,² imines,³ hydrazones,⁴ epoxides,⁵ etc.), methodologies have mainly focused on simple allylindium reagents (generated in situ from allyl bromide and indium metal) as the reactive nucleophilic species. Herein, we report the diastereoselective allylation of 4-acetoxy-2-azetidinone from a unique allylindium precursor, acylnitroso-derived hetero-Diels-Alder adduct **1**, to incorporate highly functionalized cyclopentenones at the azetidinone C4 position.

Our group has optimized conditions for Pd(0)/InI-mediated allylations of aldehydes and ketones with several hetero Diels-Alder cycloadducts ($\pm$)**1** (Scheme 1).⁶ By investigating aromatic aldehydes, aliphatic aldehydes, and ketones as suitable electrophiles for the metal-mediated allylation reactions, a diverse set of 1,4-*syn* products **2** have been prepared in 39–73% yields (Scheme 1). The stereo- and regioselective outcome may be rationalized by initial Pd(0)-mediated ring opening of hetero-Diels-Alder derived cycloadduct **1** to form π -allylpalladium(II) complex **3**.⁷ The resultant π -allylpalladium(II) **3** may undergo reductive transmetalation with indium (I) iodide⁸ to produce allylic indium (III) species **4**. Diastereo- and regioselective allylation may occur through indium-aldehyde chelation in proposed transition state **5** and accounts for stereocontrol at the peripheral carbon to afford allylation product **2**.

We were eager to identify additional electrophiles to react with hetero-Diels-Alder cycloadduct ($\pm$)**1** under Pd(0)/InI-mediated allylation conditions. We turned our attention to iminium species which display similar electrophilic character as the successful carbonyl-containing electrophiles. Accordingly, *N,N*-dimethylmethyle-ammonium iodide (Eschenmoser's salt) was selected as an electrophile to probe the reactivity of allylindium species **4** (Scheme 2).

When Boc cycloadduct ($\pm$)**6** was exposed to Pd(0) and InI in the presence of Eschenmoser's salt, the reaction was complete within 1.5 h and revealed a mixture of *anti*-1,4- and *syn*-1,4 dimethylaminomethyl-cyclopentenones **8a** and **8b** (4:1 *anti:syn*) as an inseparable mixture of diastereomers in 45% overall yield. In contrast to previously studied reactions with carbonyls,⁶ the iminium species does not have electrons available to coordinate to indium in the transition state and *anti*-1,4 products result. Therefore, allylation is directed to the less hindered top face of proposed allylindium species **7** to afford major 1,4-*anti* product **8a**. Interestingly, the reaction of dimethylamine (hydrolysis product of Eschenmoser's salt) with π -allylpalladium(II) complex **3** is competitive with reductive transmetalation and thus affords undesired byproduct **9** in 11% yield.

Encouraged by our initial result with Eschenmoser's salt, we investigated acyl imine precursor,⁹ 4-acetoxy-2-azetidinone, as a suitable electrophile. The reaction of 4-acetoxy-2-azetidinone with allylindium and crotylindium species has been reported.¹⁰ Additionally, Lee and co-workers have disclosed indium-mediated allenylations of 4-acetoxy-2-azetidinones and subsequent cyclization to carbapenems.¹¹ Our research group has previously synthesized related carbapenems;¹² allylation products derived from azetidinones and cycloadduct ($\pm$)**1** would provide new carbocyclic platforms that would allow us to gain entry to tricyclic β -lactam antibiotics¹³ and related structures.

Phenylacetyl cycloadduct ($\pm$)**10** was treated with 4-acetoxy-2-azetidinone in the presence of Pd(0) and InI to afford β -lactam-derived carbocyclic scaffold **11**¹⁴ in 78% isolated yield as exclusively the *syn*-1,4-substituted diastereomer with additional stereocontrol at the C4 azetidinone center (Scheme 3). ¹H NMR and LCMS of the crude reaction mixture confirmed the absence of additional allylation products. When Boc cycloadduct ($\pm$)**6** was exposed to Pd(0) and indium (I) iodide in the presence of 4-acetoxy-2-azetidinone, derivatized azetidinone **12** was obtained as a single diastereomer in 58% isolated yield; an identical isolated yield was attained when the reaction was performed on large scale (33 mmol).

In order to identify the newly formed stereocenter at C4 of the azetidinone, we embarked on 2D NMR studies while concurrently growing suitable crystals of **11** for X-ray crystallography. We immediately identified product **11** as the 1,4-*syn* substituted diastereomer from diagnostic *J* values.¹⁵ Unfortunately, several resonances in the ¹H NMR spectrum were very broad and required manipulation of sinebell and line broadening functions (massaging) to extract *J* values. Even though we identified $J(\text{H}_c\text{--H}_f) = 6.8$ Hz, the relative stereochemistry could not be elucidated without additional structural information (Figure 1).

A series of 2D NMR experiments were performed (COSY, TOCSY, HMQC, and HMBC) to unambiguously assign all proton and carbon resonances.¹⁶ Finally, the 3D structure of **11** was determined by ROESY. The relative stereochemistry at the C4 center of azetidinone **11** was elucidated from nOes between H_a and H_c and H_e and H_g, respectively (Figure 1). Crystals of **11** were obtained by recrystallization from acetonitrile and the proposed structure was confirmed by X-ray crystallography (Figure 2).

The exclusive diastereoselectivity suggests that coordination to indium in the transition state is relevant. In order to test this hypothesis, phenylacetyl hydroxamic acid **13** was subjected to Pd(0)/InI in the presence of 4-acetoxy-2-azetidinone (Scheme 4). Identical product **11** was isolated as the exclusive allylation product in decreased yield (41%). However, non-hydroxamate containing starting materials, phenylacetyl amide **14** and Boc carbamate **15**, did not provide allylation products when exposed to Pd(0)/InI conditions in the presence of 4-acetoxy-2-azetidinone. Although complicated mixtures resulted, consumption of acetate starting materials **14** and **15** suggests that an allylindium species formed but did not react with 4-acetoxy-2-azetidinone.¹⁷

Although the exact origin of diastereoselectivity at the azetidinone center is unknown, an organized transition state involving coordination of the acyl imine electrophile may be appropriate. Additionally, acetic acid is produced during Lewis acid-mediated acyl imine formation and may play a role in the observed diastereoselectivity.

In summary, we have reported a new diastereoselective allylation of 4-acetoxy-2-azetidinone with an allylic indium (III) species generated from cycloadduct ($\pm$)**1**. The lack of C–C bond formation in the case of starting materials **14** and **15** indicates that hydroxamate is a crucial component in these Pd(0)/InI-mediated allylations.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgment

We would like to thank Dr. Jed Fisher (University of Notre Dame) for helpful discussions, Dr. Bill Boggess (University of Notre Dame) and Nonka Sevova (University of Notre Dame) for mass spectroscopic analyses, Dr. Jaroslav Zajicek (University of Notre Dame) for NMR assistance and Dr. Bruce Noll (University of Notre Dame), Dr. Allen Oliver (University of Notre Dame), and Jeffrey Rood (University of Notre Dame) for X-ray crystallography data. We acknowledge The University of Notre Dame and NIH (GM068012) for support of this work.

References

1. For recent reviews: (a) Araki S, Hirashita T, Crabtree RH, Mignos DMP. *Comprehensive Organometallic Chemistry III* 2007; Vol. 9 OxfordElsevier Chapter 9.14. (b) Podlech J, Maier TC. *Synthesis* 2003;5:633–655.655
2. Loh T-P, Tan K-T, Yang J-Y, Xiang C-L. *Tetrahedron Lett* 2001;42:8701–8703.
3. Loh T-P, Ho DS-C, Xu K-C, Sim K-Y. *Tetrahedron Lett* 1997;38:865–868.
4. Cook GR, Kargbo R, Maity B. *Org. Lett* 2004;6:1741–1743. [PubMed: 15151403]
5. Yadav JS, Anjaneyulu S, Ahmed M, Reddy BV. *Tetrahedron Lett* 2001;42:2557–2559.
6. Lee W, Kim KH, Surman MD, Miller MJ. *J. Org. Chem* 2003;68:139–149. [PubMed: 12515472]
7. (a) Mulvihill MJ, Surman MD, Miller MJ. *J. Org. Chem* 1998;63:4874–4875. (b) Surman MD, Mulvihill MJ, Miller MJ. *Org. Lett* 2002;4:139–141. [PubMed: 11772110]
8. Araki S, Kamei T, Hirashita T, Yamamura H, Kawai M. *Org. Lett* 2002;2:847–849. [PubMed: 10814439]
9. Ueda Y, Maynarda SC. *Tetrahedron Lett* 1985;26:6309–6312.
10. Kang K-U, Baik T-G, Jiao X-H, Lee K-J, Lee C-H. *Synlett* 1999;4:447–449.
11. Lee PH, Kim H, Lee K, Kim M, Noh K, Kim H, Seomoon D. *Angew. Chem. Int. Ed* 2005;44:1840–1843.
12. (a) Williams MA, Miller MJ. *Tetrahedron Lett* 1990;31:1807–1810. (b) Williams MA, Hsiao CH, Miller MJ. *J. Org. Chem* 1991;56:2688–2694.
13. (a) Di Modugno E, Erbeti I, Ferrari L, Galassi GL, Hammond SM, Xerri L. *Antimicrob. Agents Chemother* 1994;38:2362–2368. [PubMed: 7840571] (b) Rossi T, Biondi S, Contini S, Thomas RJ, Marchioro C. *J. Am. Chem. Soc* 1995;117:9604–9605. (c) Martel SR, Wisedale R, Gallagher T, Hall LD, Mahon MF, Bradbury RH, Hales NJ. *J. Am. Chem. Soc* 1997;119:2309–2310.
14. Hydroxamic acid **11** was purified on iron-free silica gel; refer to Supporting Information for preparation of iron-free silica gel.
15. See Surman MD, Mulvihill MJ, Miller MJ. *J. Org. Chem* 2002;67:4115–4121. [PubMed: 12054945] footnote 14.
16. Refer to Supporting Information for complete carbon and proton assignments for **11**.
17. Phenylacetyl amide **14** reacts with aldehydes in the presence of Pd(0) and InI to afford 1,4-*anti* products, ref. 6.

Figure 1.
Selected nOes in ROESY spectrum of 11.

Figure 2.
X-ray structure of **11** with thermal ellipsoids drawn at 50% probability.

Scheme 1.
Pd(0)/InI-Mediated Allylation with Cycloadduct ($\pm$)**1**

Scheme 2.
Pd(0)/InI-Mediated Allylation of Eschemoser's Salt

Scheme 3.
Pd(0)/InI-Mediated Allylation of 4-Acetoxy-2-Azetidinone

entry	substrate	R	R ¹	product	isolated yield (%)
1	13	CH_2Ph	OH	11	41
2	14	CH_2Ph	H	-	0
3	15	$\text{OC(CH}_3)_3$	H	-	0

Scheme 4.
Pd(0)/InI-Mediated Allylation with **13–15**