Published in final edited form as:

Cell Mol Life Sci. 2009 March ; 66(5): 852–875. doi:10.1007/s00018-008-8609-x.

Molecular Mechanisms of BK Channel Activation

Jianmin Cui, Huanghe Yang, and Urvi S. Lee

Department of Biomedical Engineering and Cardiac Bioelectricity and Arrhythmia Center,
Washington University, 1 Brookings Drive, St. Louis, MO 63130, USA

Abstract

Large conductance, Ca^{2+} -activated potassium (BK) channels are widely expressed throughout the animal kingdom and play important roles in many physiological processes, such as muscle contraction, neural transmission and hearing. These physiological roles derive from the ability of BK channels to be synergistically activated by membrane voltage, intracellular Ca^{2+} and other ligands. Similar to voltage-gated K^+ channels, BK channels possess a pore-gate domain (S5–S6 transmembrane segments) and a voltage sensor domain (S1–S4). In addition, BK channels contain a large cytoplasmic C-terminal domain that serves as the primary ligand sensor. The voltage sensor and the ligand sensor allosterically control K^+ flux through the pore-gate domain in response to various stimuli, thereby linking cellular metabolism and membrane excitability. This review summarizes the current understanding of these structural domains and their mutual interactions in voltage-, Ca^{2+} - and Mg^{2+} -dependent activation of the channel.

Keywords

channel gating; MaxiK; calcium activation; voltage activation; metal binding; allosteric

I. Introduction

BK channels have a large single channel conductance ($\sim 100\text{--}300$ pS), which is the basis for their designation as Big K^+ (or MaxiK) channels [1–3]. BK channels are found in neurons [4–8], chromaffin cells [9–13], inner hair cells of cochlea [14–18] and skeletal [19–21] and smooth muscles [22–29]. These channels are activated by membrane depolarization and elevation of intracellular Ca^{2+} concentration ($[\text{Ca}^{2+}]_i$) (Fig. 1A). Due to these properties, the activation of BK channels results in repolarization of the membrane and closing of voltage dependent Ca^{2+} channels to reduce Ca^{2+} entering the cell. Therefore, BK channels primarily serve as negative feedback regulators of membrane potential and $[\text{Ca}^{2+}]_i$, which are important in a number of physiological processes. These include controlling the interspike interval and spike frequency adaptation [5,6,30–34], modulating neurotransmitter release [17,32,35–38] and endocrine secretion [39–41], tuning hair cells firing frequencies in the auditory system [15,16,42–47], and regulating vascular [48–52], urinary bladder [53–56] and respiratory tone [26,57–59]. Consistent with their physiological importance, malfunction of BK channels can lead to epilepsy [60,61], motor impairment [62], noise-induced hearing loss [63], hypertension [49,64–70], urinary incontinence [25], overactive urinary bladder [25,71] and asthma [72].

In certain cells, BK channels are in close proximity to Ca^{2+} sources. For example, in neurons, BK channels are colocalized or physically associated with voltage dependent Ca^{2+} channels (VDCCs) [73–77], NMDA receptors [78] and ryanodine receptors (RyRs) [74]. Normally, the

global $[Ca^{2+}]_i$ is carefully regulated that small changes in the global $[Ca^{2+}]_i$ will only open few BK channels even at the peak of action potential. However, the $[Ca^{2+}]_i$ near the Ca^{2+} source, termed the microdomain $[Ca^{2+}]_i$ can be much higher [79] to ensure the opening of additional BK channels. The large K^+ efflux through BK channels efficiently repolarizes the membrane and halts Ca^{2+} entry.

Besides voltage and Ca^{2+} , intracellular Mg^{2+} also activates BK channels [80,81]. The activation of the BK channel by multiple stimuli is an important property that allows the channel to integrate various cellular signals in modulating membrane excitability and Ca^{2+} homeostasis. This property also makes the BK channel a fascinating molecule in studying the mechanism of ion channel gating. The purpose of this review is to summarize the current understanding of the molecular gating mechanisms of BK channels in response to voltage, Ca^{2+} and Mg^{2+} , as well as the major historic developments leading to these understandings. We will first sketch the structure of BK channels that reveals their homology to both voltage-gated and ligand-gated channels. Such a homology and functional similarities to both voltage and ligand gated channels suggest a similarity in the underlying molecular gating mechanisms. We will then dissect the gating mechanism for each stimulus: voltage, Ca^{2+} and Mg^{2+} . Channel gating involves two major steps: sensing of the stimulus by the sensor domain and conveying the conformational change from the sensor domain to open the activation gate, known as coupling. Structural information and functional studies of ion channels have shown that distinct structural domains underlie the activation gate and sensors for stimuli [82]. Accordingly, the gating mechanism of each of the stimuli for BK channels is divided into two parts, the identification of the sensor and the mechanism of coupling. We will then describe the effect of each stimulus on the mechanism of other stimuli in activating BK channels. Finally, we will describe the activation gate itself. Readers may obtain more insights from other excellent reviews about BK channel's role in physiology [2,33,51,83,84], gene family [85], modulation [86–89], auxiliary subunits [39,64,90], pharmacology [91–93] and gating mechanisms [94–101].

II. Structure of BK channels

BK channels are encoded by a single *slowpoke* gene termed *Slo1* [102–104]. These channels were first identified in mutant *Drosophila melanogaster*, which exhibited a lethargic phenotype due to a lack of Ca^{2+} dependent K^+ current [102,103]. Soon after, the mammalian orthologue of this gene was cloned from mouse and human [104,105]. The initial hydropathy analysis suggested that a Slo1 protein may contain 10 hydrophobic segments [104]. Subsequent electrophysiology characterization and immunocytochemistry experiments on epitope tagged BK channels revealed that the Slo1 protein is comprised of a membrane spanning domain with 7 transmembrane segments (S0–S6) and a large C-terminal cytoplasmic domain [106,107] (Fig. 2A).

The channel's primary sequence of the membrane spanning domain showed similarities to voltage-dependent K^+ (Kv) channels, including the voltage sensing (VSD, S1–S4) and pore-gate (PGD, S5–S6) domains (Fig. 2A). Similar to Kv channels, a functional BK channel is formed by tetramerization of Slo1 proteins [108,109]. However, unique to BK channel is an additional transmembrane segment, SO that confers the N-terminus to the extracellular side and a large cytoplasmic domain containing ~800 amino acids. The sequence and predicted secondary structure of the cytoplasmic domain are homologous to a regulatory domain for K^+ conductance (RCK domain) (Fig. 2) that is found in a number of K^+ channels and transporters [97,110–114]. Thus, a functional BK channel can be divided into three major structural domains: the VSD senses voltage, the C-terminal cytoplasmic domain senses various intracellular ligands, and the PGD controls ion permeation in response to different stimuli.

The 3-dimensional (3D) structure of the BK channel has not been solved. However, the X-ray crystallographic structure of Kv1.2 (Fig. 2B) [115] has been used as a model for the membrane-spanning domain of BK channels due to their sequence homology and functional similarity [116–119]. This model has not been systematically tested by experiments, but it is almost certain that the details of the BK channel structure would differ from that of Kv1.2, primarily because of the additional S0 transmembrane segment [117,120] and the long linker between S0 and S1 (~70 amino acids). By performing a set of disulfide crosslinking experiments in which double residues, one flanking S0 and the other flanking S1, S2, S5, S6 transmembrane segments, or in the S3–S4 loop, were mutated to cysteine, Liu et al. showed that the extracellular end of S0 is likely to reside close to S1, S2 and the S3–S4 loop, but not S5 or S6 [117]. In a model based on Kv1.2 structure, these results would place the extracellular end of S0 in the cleft among S1–S4 helices that insulate S0 from the pore domain. Koval et al. inspected the function of S0 by performing a tryptophan-scan and found that the mutants F25W, L26W and S29W, which are on the same side of the predicted S0 helix embedded in the membrane, altered channel function [120]. This result suggests that the middle of S0 may make direct contact with the VSD (S1–S4). The presence of S0 and its close contacts with S1–S4 would affect the conformation of BK channels differently in the open and the closed states [117,120].

Homology models for the cytoplasmic domain of BK channels have been proposed based on the RCK domain structures of the *Escherichia coli* 6-TM K⁺ channel and the Ca²⁺-gated K⁺ channel, MthK, from *Methanobacterium thermoautotrophicum* [94,110,111,121–126]. These prokaryotic RCK domains adopt a Rossman fold topology at its core with β strands (β A– β J) sandwiched by α helices (α A– α J) [110,111]. In the quaternary structure of the MthK channel, eight identical RCK domains assemble into an octameric ring-like shape that is known as the gating ring (Fig. 2C). Each RCK domain is associated with two neighboring RCK domains alternately with either a fixed (or assembly) or flexible interface. The cytoplasmic domain of Slo1 protein can be divided into two separate structures [127], both with sequence homology to RCK domain [110], known as RCK1 and RCK2. Therefore, it was proposed that BK channels would also contain a gating ring formed by eight RCK domains from the four Slo1 subunits [94,110,124–126].

Several laboratories have published similar sequence alignments between the RCK1 of BK channels and the prokaryotic RCK domains (Table 1). The predicted RCK1 structure has been supported by a number of experimental results. First, based on the structure of the RCK domain in the *E. coli* K⁺ channel, it was predicted that in the putative RCK1 domain, K448 at the C-terminus of α D and D481 in the loop between α E and β F would form a salt bridge [111]. The experimental data on the BK channel supported the existence of the salt bridge. Individual mutations, K448D and D481K both shifted the G–V relation of BK channels to more positive voltages, but the double mutation K448D:D481K restored the G–V close to that of the wild type channel [111]. Second, mutational studies suggested that E374 and E399 in RCK1 were part of the metal binding site for Mg²⁺ dependent activation [118,121,122,128,129]. When mapped onto the prokaryotic RCK structures, these two residues are located at the N-terminus of the parallel β B and β C, respectively [118,121], and are close to each other, supporting the mutational results that these two residues are part of the same metal binding site. Third, Hou et al. found that H365 and H394 in the RCK1 serve as the H⁺ sensor for BK channel modulation [130]. In a RCK1 homology model based on the MthK structure, the two histidine residues were found close to D367, which serves as a Ca²⁺ sensor [122]. Hou et al. found that the mutations on H365 and H394 affected Ca²⁺ sensitivity, and reciprocally, the mutations of D367 altered H⁺ sensitivity. These results are consistent with the model prediction that these residues affect both H⁺ and Ca²⁺ sensitivities with an electrostatic interaction [130]. The above experimental results all demonstrated that residues far apart in the sequence are located close in the 3D structure as predicted by the homology model of the RCK1 domain.

Unlike for RCK1, there has been little consensus in published sequence alignments for RCK2 (Table 1). Furthermore, based on a statistical calculation, Fodor and Aldrich suggested that the sequence homology between the BK channel and other RCK domains may not be sufficient to support the presence of the RCK-like structure in the putative RCK2 domain [112]. Nevertheless, using circular dichroism (CD) spectroscopy, Yusifov et al. measured the secondary structure composition of the C-terminus sequence of human Slo1 corresponding to 90% of the proposed RCK2 domain, and found that the α -helix and β -strand contents closely correlated with the predicted secondary structure based on their proposed structural alignments [124]. In another study, using the MthK gating ring as a guide, Kim et al. examined if BK channel residues arising from different RCK domains participated in a putative fixed interface. Using a mutant cycle strategy, they measured G-V relations of mutant channels at a fixed $[Ca^{2+}]_i$ of 5 μ M, and found that the coupling-energy of the mutant cycles was not zero. These results led Kim et al. to argue that residues from RCK1 and RCK2 interact, thereby participating in an interface [126]. On the other hand, the proposed RCK2 structure was predicted to contain a salt bridge between K898–D927, similar to that found in the RCK1 domain [111]. However, unlike the experimental results with the salt bridge in the RCK1 domain [111], mutations of K898 or D927 did not cause any significant functional change [126]. These mixed results make the homology model of RCK2 less certain than that of RCK1.

While the structures of Kv1.2 and MthK serve as a template for the homology model of the membrane spanning and the cytoplasmic domains, no structural template is available to show how the two domains of the BK channel assemble. Recent studies show that residues from both the membrane spanning VSD and the cytoplasmic RCK1 domain form the Mg^{2+} binding site [118], and the bound intracellular Mg^{2+} makes an electrostatic interaction with the voltage sensor [129]. These results indicate that the two domains are positioned close to each other and interact intimately. On the other hand, with a series of elegant experiments Zhang et al. showed that the BK channel has a central antechamber between the pore domain and the cytoplasmic domain that can accommodate the inactivation peptide from the accessory β 2 subunit and protect it from intracellular trypsin digestion [131]. Together, these results suggest that at some positions the membrane spanning and cytoplasmic domains come near each other, while at other positions they may be sufficiently separated to allow entry of bulky peptides. Besides the vertical distances between the membrane spanning and cytoplasmic domains, another question concerning domain assembly is how the VSD and the cytoplasmic domain align laterally with each other in BK channels. Figure 2D shows the combined structures of Kv1.2 [115] and MthK [110] by aligning the selectivity filter of the two channels. This figure shows that, if the VSD and the RCK1 domain of BK channels pack against the PGD similarly to that of Kv1.2 and MthK, the VSD would be located just above the RCK1 domain from the same subunit. However, by studying the formation of the Mg^{2+} binding site, Yang et al. recently suggested that the VSD of one subunit is located on top of the RCK1 domain from the neighboring subunit (See Section IIIC for more details) [118]. The contradiction between the experimental results and the homology model implies that the packing of VSD or the cytoplasmic domain relative to the PGD in BK channels may differ from that of Kv1.2 or MthK channels. Several unique features of BK channels may contribute to this discrepancy: 1) the VSD of BK channels contains an additional S0 segment and a long cytoplasmic loop connecting S0 and S1; 2) the interaction and packing between the RCK1 domain and the rest of the cytoplasmic domain of BK channels may differ from that of the RCK domains in the gating ring of the MthK channel; 3) other interactions may exist between the VSD and the cytoplasmic domain.

III. Voltage-, Ca^{2+} - and Mg^{2+} -Dependent Activation of BK Channels

A. Voltage-dependent activation of BK channels

a. Voltage sensors—The voltage dependence of BK channels changes in different $[\text{Ca}^{2+}]_i$ (Fig. 1A). In the early days, there were two different views on the origin of the voltage dependence. One suggested that the voltage dependent activation derived from the voltage dependent binding of Ca^{2+} ; while the other proposed that the voltage dependence was one of the intrinsic properties of BK channels, independent from the Ca^{2+} dependent activation. The latter view gained more support after the cloning of the BK channel, which predicted the presence of a VSD. Specifically, the S4 transmembrane segment contains multiple regularly-spaced Arg residues, a hallmark of VSD in Kv channels [102–104]. Macroscopic ionic current measurements in the absence of Ca^{2+} [132,133] and gating current recordings [134,135] support the notion that BK channels, similar to Kv channels, contain an intrinsic VSD. Mutation of the Arg residues in S4 to neutral amino acids reduced the steepness of the G-V relation [136,137], further suggesting that the S4 segment in BK channels also serves as a voltage sensor.

To identify the charged residues in the VSD (S1–S4) that contribute to voltage sensing, Ma et al. mutated each of these residues and measured the reduction in the effective gating charge [116]. The number of gating charges in the WT and mutant BK channels in their study was estimated by measuring the probability of channel opening (P_o : $10^{-6} \sim 10^{-1}$) over a broad voltage range and fitting the data to the allosteric HA model (Fig. 3A, see below). Four residues were thus identified and referred to as the voltage sensing residues: D153 and R167 in S2, D186 in S3, and R213 in S4 (Fig. 2A). Neutralization of each of these residues in four mSlo1 subunits reduced effective gating charge by 0.92, 0.48, 0.88, and 1.20, respectively, from 2.32 of the WT channel [116,135]. In an earlier study, Diaz et al. reported that, besides R213, R210 in S4 also contributed to gating charge based on limiting slope measurements [136]. However, Ma et al. found that with more stringent conditions (at $P_o \leq 10^{-6}$ vs. at $P_o \geq 10^{-3}$ in Diaz et al., 1998), limiting slope measurements indicate that R210 is not a voltage-sensing residue [116].

Compared with Kv channels such as Shaker, which has ~12–13 effective gating charges [138,139], the BK channel is less sensitive to voltage. Besides this difference, two unique features of the voltage sensor of BK channels are noteworthy: 1) Out of 3 Arg residues in S4, only R213 is a voltage sensing residue. Mutations of the other Arg residues (R207 and R210) do not affect the number of gating charge [116]. According to the alignment of Shaker and BK channels [116,140], R213 in S4 of BK channels corresponds to the 4th Arg in S4 of Kv channels. Each of the first 4 Arg residues (R1–R4) in S4 of Kv channels moves across partial or the entire transmembrane electric field, and accounts for either a half or a full gating charge [139]. However, in S4 of BK channels, only R213 accounts for a mere 0.3 gating charge [116]. This result suggests that either S4 in BK channels does not move a distance along the voltage gradient as large as that in Kv channels during activation or the voltage drop along the path of S4 movement is small. 2) Besides R213 in S4, other voltage sensing residues, D153, R167 and D186, are located in either S2 or S3. These residues are also conserved in Kv channels; none of them, however, serves as gating charge. Interestingly, in Shaker K^+ channels, the residues corresponding to D153 and D186 have been shown to form a network of electrostatic interaction with Arg residues in S4 at either the resting or active state [115,141,142]. If such a network of electrostatic interaction exists among the voltage sensing residues in the BK channel, it is possible that a mutation to any one of these residues could alter the movement of others, resulting in an indirect effect on the number of effective gating charges. This or other unspecified indirect effects must exist, because the summed decreases in the number of effective gating charge produced by neutralizing these residues (3.48) exceed the total charge of the WT channel (2.32) [116]. At present, it is not clear which of the 4 voltage sensing residues are affected by such indirect effects and how much of their reported contribution to the gating

charge is due to indirect effects. The contribution of specific amino acid residues to the effective gating charge in Kv channels has been estimated by a number of different approaches [138, 139], and the results, in general, agree with one another. For BK channels, more studies using different approaches would help to more firmly establish the identity and contribution of individual amino acid residues as voltage sensors.

b. Coupling between voltage sensors and the activation gate—In response to depolarization, the voltage sensor in BK channels moves from a resting state to an active state, resulting in transient gating currents (I_g) [134,135]. While it enhances P_O , voltage sensor activation is not obligatory for channel opening. At negative voltages (≤ -20 mV), where the voltage sensor of the BK channel is mostly at the resting state and the probability of channel opening due to voltage sensor movement is small ($P_O \leq 10^{-4}$), residual channel opening can be detected [143]. Channel opening at this voltage range has a much weaker voltage dependence than in positive voltages (30 – 80 mV), indicating that the activation gate can open with a weak voltage dependence while voltage sensors are in the resting state.

Analysis of the kinetics and steady state properties of BK currents suggests that, at positive voltages, the voltage sensor in each of the four subunits moves from a resting state to an active state independently, and the channel can open when 0, 1, 2, 3 or 4 voltage sensors are activated. Horrigan et al. proposed an allosteric model (the HCA model) to describe the coupling between voltage sensor and the activation gate (Fig. 3A) [143]. In this model, voltage sensors can move from the resting state to the active state at either closed or open conformation, and voltage sensor activation promotes channel opening (with an equilibrium constant L) by selectively destabilizing the closed conformation with an allosteric factor D . Similarly, the equilibrium constant for voltage sensor activation (J) increases D -fold in favor of the active state, when the channel opens (Fig. 3A). The model illustrates the idea that the BK channel undergoes two types of conformational change: voltage-sensor activation and channel opening; each type is not obligatory for the other to occur but they influence each other with an allosteric mechanism. This two-tiered allosteric model is consistent with the kinetic properties of the macroscopic K^+ current in response to voltage pulses; the activation and deactivation time courses of the current are exponential after a brief delay [133,143]. Single channel analyses have also revealed multiple open and closed states in 0 Ca^{2+} , supporting the two-tiered allosteric model for voltage dependent activation [144–146]. The voltage dependence of channel gating derives primarily from the voltage dependence of transitions among closed (C-C) or open states (O-O) [146], compatible with the idea that the voltage sensor activates at either closed or open states, while channel opening has a weak voltage dependence [143].

The properties of gating currents (I_g) are also consistent with key features of the HCA model. First, the kinetics of gating current relaxation is rapid [134,143]. The time course of the fast on-gating current accounts for the initial delay in channel opening after a depolarizing voltage pulse, and its relaxation is almost completed before channels begin to open [143]. This result reveals voltage sensor activation and channel opening as two types of conformational change, and the movement of voltage sensors promotes channel opening. Second, gating currents contain kinetically distinctive components, which can be assigned to voltage sensor movements during C-C, O-O, and C-O transitions [143]. These results clearly demonstrate that voltage sensors can move in both the closed and open conformations. Third, voltage sensor activation contains a slow component that increases with a time course in parallel with channel opening [143]. This result is consistent with the idea that voltage sensor activation is facilitated at the open conformation; as more channels open, the easier it is for the gating charges to move to the active state. In a study in which residues in the S3–S4 linker of hSlo1 were tagged with fluorescent molecules, Savalli et al. showed that the change of fluorescence had time constants similar to that of ionic currents (I_k), but voltage dependences similar to that of I_g , consistent with the view that the VSD moves as the BK channels open [147].

In contrast to the allosteric model for BK channel gating, so-called obligatory models are usually used to describe Kv channel gating. The simplest obligatory model is the Hodgkin-Huxley type sequential model, C-C-C-C-O, in which four identical voltage sensors are activated independently by depolarization, and the channel remains closed (C) until all four voltage-sensors are activated that opens the channel (O) [82]. This model assumes an obligatory coupling between voltage-sensor activation and channel opening: voltage-sensor must move to an active state in order for the channel to open, and channel opening is obligatory once all four voltage-sensors are activated. However, one prevalent model proposed for Shaker K⁺ channels includes a C-O transition after all four voltage-sensors are activated [148,149]. This model shares similarity with the allosteric model for BK channels based on the idea that the channels undergo two types of conformational change: voltage-sensor activation and channel opening. In this model, although voltage sensor activation is obligatory for the channel to open, channel opening is not obligatory after voltage sensors are activated. This concept was best illustrated by the ILT mutation of Shaker K⁺ channels [149], which includes three conservative amino acids substitutions in S4, V369I, I372L and S376T. The ILT mutation alters the rate constants of channel opening to make this conformational change rate limiting, resulting in a separation of voltage ranges for channel opening and for voltage sensor activation. Therefore, activation of voltage sensors saturates at a voltage ~60 mV below the threshold of activation of ionic currents [149]. In other words, within the 60 mV voltage-interval, the activation gate does not open even though voltage sensors have been activated. In Shaker channels, the voltage dependence of P_O does not change even when P_O is as low as 10⁻⁷ at negative voltages [148], which differs from the results of BK channels [143]. This result is consistent with the idea that voltage sensor activation is obligatory for the Kv channel to open [138]. However, in the case of BK channels, it was shown that, if the parameters of the HCA model are altered in certain ways, channel opening can also be difficult to observe when voltage sensors are at the resting state [135,143]. In other words, the possibility remains open that the coupling between voltage sensor activation and channel opening is actually allosteric even in Shaker channels, but the properties of the two types of conformational change happen to be such that it would not allow the observation of channel opening when voltage sensors are at the resting state. Consistent with this view, some mutations that disrupt physical connections between voltage sensors and the pore in Shaker channels make the channels open at negative voltages [150]. In fact, an allosteric model of Shaker channel gating was proposed to account for the block of 4-aminopyridine (4-AP) on both the ionic and gating currents [151]. Taken together, although different models have been developed to describe voltage dependent gating of BK and K_v channels, the coupling between voltage sensor activation and channel opening in these channels may be fundamentally similar.

The structural basis of the coupling between voltage sensors and the activation gate has not been explored in BK channels. In K_v channels, interactions between the S4-S5 linker and the C-terminus of S6 are important for the coupling between voltage sensor activation and channel opening [152]. Experimental evidence suggests that a direct interaction between S4 and S5 also contributes to the VSD-PGD coupling [149,153]. Based on above discussions and the structural homology between BK and K_v channels, it is likely that similar molecular mechanisms apply to BK channels. However, the unique structural features of BK channels may alter the molecular mechanism of the coupling between voltage sensors and the activation gate: 1) The additional S0 TM segment and the long intracellular S0-S1 linker increase molecular mass of the VSD and may interact with parts of the VSD (see Section II), thereby potentially altering the interactions of the VSD with the pore domain [118]. 2) The residues in the large cytoplasmic domain interact with those in the VSD to affect voltage sensor activation [118,129] (see Sections II and IIIC). On the other hand, the cytoplasmic domain may affect channel opening by pulling S6 via the peptide linker [154] (see Section IIIB). Thus, the cytoplasmic domain of the BK channel may affect the coupling between voltage sensor and the activation gate by interactions with both domains.

B. Ca^{2+} -dependent activation of BK channels

a. Ca^{2+} binding sites—Intracellular Ca^{2+} binds to the cytoplasmic domain of BK channels to promote channel opening [127]. Each Slo1 subunit of BK channels contains two high-affinity Ca^{2+} binding sites (K_d : 0.8 ~ 11 μM) [85,97,122,155–157]. One site is located in a motif in the C-terminus of the cytoplasmic domain that contains a series of Asp residues known as the “ Ca^{2+} bowl” [155], and the other is in the RCK1 domain [122].

The Ca^{2+} bowl was first proposed as a Ca^{2+} binding site because it may structurally resemble the Ca^{2+} binding loop of serine proteinases [158,159] and it contains a high density of acidic residues [155]. Mutations to the Asp residues in the Ca^{2+} bowl reduced Ca^{2+} sensitivity of channel activation [122,155,156,160,161]. By eliminating 1, 2, 3, and 4 functional Ca^{2+} bowls from four subunits, the apparent Ca^{2+} binding affinity and the Hill coefficient, which represents the allosteric interaction among multiple Ca^{2+} binding sites, diminish in a step-wise fashion [162]. By studying channels containing C-terminus chimeras between the Ca^{2+} -sensitive mSlo1 and the Ca^{2+} -insensitive mSlo3, Schreiber et al. revealed that the Ca^{2+} bowl region of mSlo1 is required for part of Ca^{2+} sensitivity [163]. In most studies, Ca^{2+} sensitivity of the channel is measured by the G-V shift caused by varying $[\text{Ca}^{2+}]_i$ (Fig. 1). The method to directly and quantitatively measure Ca^{2+} binding to BK channels is not available. However, employing a $^{45}\text{Ca}^{2+}$ -overlay technique, Bian et al. and Braun and Sy showed that purified C-terminal peptides of Slo1 containing the Ca^{2+} bowl region binds Ca^{2+} [160,161]. Furthermore, mutating Asp residues in Ca^{2+} bowl significantly reduced Ca^{2+} binding [160]. More recently, using circular dichroism (CD) spectroscopy, Yusifov et al. found that the secondary structure composition of a hSlo1's C-terminal peptide that includes the Ca^{2+} bowl changed with Ca^{2+} concentration in solution, and the mutation of five sequential Asp to Asn (5D5N) in the Ca^{2+} bowl significantly reduced the Ca^{2+} effect [124]. These experiments provide a strong support for Ca^{2+} binding to the Ca^{2+} bowl. In a comprehensive study of mSlo1, it was shown that the acidic residues were not equal in binding to Ca^{2+} [156]. Mutation to individual Asp residues showed various degrees of reduced Ca^{2+} sensitivity, as measured by G-V shifts, and Ca^{2+} binding, as measured by $^{45}\text{Ca}^{2+}$ -overlay [156]. Among these mutations, D898A and D900A eliminated the Ca^{2+} sensitivity contributed by the Ca^{2+} bowl and caused the largest reduction in Ca^{2+} binding. Therefore, it was proposed that D898 and D900 serve as ligands for Ca^{2+} coordination. In the same study, due to discrepancies between the effects of mutations in the intact subunit on Ca^{2+} -dependent gating and in the isolated peptides on Ca^{2+} binding, Bao and Cox suggested that in the biochemistry studies the purified peptides may have a somewhat distorted structure [156]. In addition, in the studies using a $^{45}\text{Ca}^{2+}$ -overlay technique or CD spectroscopy, the mutations in the Ca^{2+} bowl could not abolish Ca^{2+} effects entirely, suggesting that Ca^{2+} binding to the purified peptides may not be specific to the Ca^{2+} bowl [124,160,161].

Mutations in the Ca^{2+} bowl reduce, but do not eliminate Ca^{2+} sensitivity of BK channels [155,163]. Additionally, besides Ca^{2+} , Cd^{2+} activates BK channels, but Cd^{2+} activation is not modified by the mutations in the Ca^{2+} bowl [155]. Therefore, it was proposed that the channel contained a second high affinity Ca^{2+} binding site [155,163]. In 2002, mutations of two acidic residues (D362A/D367A) in the RCK1 domain were found to eliminate the rest of the Ca^{2+} sensitivity [122]. Subsequent characterization of this mutation indicated that it also abolished the Cd^{2+} -dependent activation [164], further supporting D362/D367 as another putative Ca^{2+} binding site. Interestingly, the human hereditary disease of generalized epilepsy and paroxysmal dyskinesia (GEPD) was found to be associated with a missense mutation of an Asp residue two amino acids downstream of D367 [60]. The Asp-to-Gly mutation increased BK channel's Ca^{2+} sensitivity. The identification of this mutation further highlights the importance of the Ca^{2+} binding site in RCK1 for Ca^{2+} sensing of BK channels. Consistent with this idea, recent studies on carbon monoxide (CO) and intracellular pH modulation of BK

channel function [130,165] indicate that these ligands may bind in the vicinity of the D362/D367 site, exerting direct or indirect effects on Ca^{2+} binding to this site.

Many molecular details about the putative Ca^{2+} binding site in RCK1 are still unclear. D362 seems to be less important than D367 in Ca^{2+} binding to the RCK1 site [121,122,166]. While D367A resembles the effect of D362A/D367A, D362 only causes a small reduction of Ca^{2+} sensitivity. More intriguingly, another mutation in the RCK1 domain, M513I, significantly reduces, but does not abolish, the Ca^{2+} sensitivity derived from the RCK1 domain [157]. Ca^{2+} prefers to bind to “hard” oxygen-containing ligands, such as carboxylates, carbonyls, water, and hydroxyl oxygen atoms. The “soft” sulfur in the side chain of Met residue usually is not considered to be a Ca^{2+} ligand [167,168]. Nevertheless, since the D362/D367 site can also bind to Cd^{2+} [164], which prefers “soft” ligands such as sulfur and nitrogen, the possibility that M513 also contributes to Ca^{2+} binding to the RCK1 site cannot be completely ruled out [169]. Because the loop containing D367 is not conserved in the MthK RCK1 domain, it is not known whether D362, D367 and M513 are indeed close to each other to participate in the same Ca^{2+} binding site. Further studies are needed to clarify their individual contribution to the Ca^{2+} binding to the RCK1 site.

The RCK1-related Ca^{2+} binding site has somewhat of a lower affinity for Ca^{2+} than the Ca^{2+} bowl, but is responsible for a larger portion of Ca^{2+} sensitivity, accounting for ~60% of total G-V shift due to Ca^{2+} binding [95,122,157]. Consistently, the Ca^{2+} bowl accelerated activation kinetics mainly at $[\text{Ca}^{2+}]_i$'s below 10 μM , while the RCK1-related site altered both activation and deactivation rate at $[\text{Ca}^{2+}]_i$'s between 10 and 300 μM [164]. The contribution to the Ca^{2+} sensitivity by each of the two sites was studied by eliminating the function of the other site. The results showed a roughly additive Ca^{2+} sensitivity, indicating that each site contributes to the channel gating independently [122,157]. However, by studying a combination of mixed mSlo1 WT and mutant subunits, in which the function of either the Ca^{2+} bowl or the RCK1-related site was eliminated, Qian et al. found that the two sites in the same Slo1 subunit cooperate positively; the two intact sensors on the same subunit are more effective in increasing channel opening than when they are on different subunits [170]. It is not clear how the cooperativity observed by Qian et al. affects the overall Ca^{2+} sensitivity quantitatively.

Piskorowski and Aldrich [171] reported that a truncated BK channel that lacks the entire cytoplasmic C-terminus, made by deleting the amino acids right after the transmembrane segment S6 (after residue 323 of mSlo1), still has a Ca^{2+} sensitivity similar to that of the wild type channels. Since neither the Ca^{2+} bowl nor the RCK1 is present in the truncated channel, this result suggested that the observed Ca^{2+} -induced activation was either due to a Ca^{2+} binding site within the membrane spanning domain or derived from a Ca^{2+} sensing molecule physically associated with the truncated channel. The truncation reduced the expression of the channel on the cell membrane of *Xenopus laevis* oocytes and mammalian cell lines [171–173] so that only single channel activities of the truncated channel could be recorded. However, the surface expression level of the truncated channel in *Xenopus* oocytes was apparently higher than that of the endogenous BK channels [171] [174]. Piskorowski and Aldrich were careful to make sure that they were indeed studying the truncated channels by including a secondary mutation T294V, which reduces the channel blockade by TEA [109] and thus can be distinguished from endogenous or contaminating WT BK channels. Consistent with these results, Qian and Magleby showed that the coexpression of the accessory $\beta 1$ subunit of BK channels with a triple mutant mSlo1, which eliminates the function of the Ca^{2+} bowl (5D5N), the RCK1-related Ca^{2+} binding site (D362A/D367A), and the Mg^{2+} binding site (E399A), partially restored Ca^{2+} sensitivity [175]. Since the $\beta 1$ subunit itself does not contain any Ca^{2+} binding site, the restored Ca^{2+} sensitivity should derive from a Ca^{2+} binding site other than the Ca^{2+} bowl or the RCK1-related site. On the other hand, there are controversies in regard to the results of

Piskorowski and Aldrich [171] and how to interpret these results in relation to the large body of studies on the Ca^{2+} bowl and the RCK1-related Ca^{2+} binding sites discussed above. In a direct comparison, Schmalhofer et al. found that the identically truncated Slo1 channels were trapped inside the TsA-201 cells and no single channel activities could be detected on the cell membrane under various recording conditions [173]. Using an iberiotoxin binding assay, they showed that the channels trapped in the intracellular membranes either were not tetramerized nor had an altered architecture in the outer vestibule of the channel. Even in *Xenopus* oocytes, the same truncated channel failed to express in different laboratories [176,177]. To examine whether the Ca^{2+} sensor resides in the membrane spanning domain or in the cytoplasmic domain, Xia et al. studied chimera channels between the Ca^{2+} sensitive mSlo1 and the pH sensitive mSlo3 channels [176]. The results showed that the cytoplasmic domain of mSlo1 conferred Ca^{2+} sensitivity when connected to either the mSlo1 or mSlo3 membrane spanning domain, while the cytoplasmic domain of mSlo3 conferred pH sensitivity. Therefore, the specific ligand sensors of the two channels are defined by the cytoplasmic domain. Another important unsettling aspect in regard to the results of Piskorowski and Aldrich is that no studies have identified any Ca^{2+} binding site in the membrane spanning domain, while a large body of data identified the Ca^{2+} bowl and the RCK1-related site in the cytoplasmic domain. Braun and Sy proposed an EF hand-like Ca^{2+} binding site in the S0–S1 linker, but mutations in this putative site only reduced Ca^{2+} sensitivity by less than 20% [161]. In a recent study, each of the oxygen-containing residues in the intracellular side of transmembrane segments and loops between them were mutated to Ala. The results showed that none of these mutations reduced Ca^{2+} sensitivity in 0–100 μM Ca^{2+} [178]. Since oxygen is the preferred ligand for Ca^{2+} coordination [168] and these residues are possibly exposed to the cytosol, this result would normally indicate that there is no Ca^{2+} binding site in the membrane spanning domain. However, both studies [176,178] were based on the entire mSlo channels containing both the membrane spanning and the cytoplasmic domains, which may use different Ca^{2+} binding sites for channel activation as compared to the truncated channel lacking the cytoplasmic domain. Therefore, taken together, the results from both studies suggest that if the membrane spanning domain contains a Ca^{2+} binding site, it does not contribute to Ca^{2+} sensing in the intact BK channel. Such a Ca^{2+} binding site may be exposed only after the removal of the cytoplasmic domain to give rise to Ca^{2+} sensitivity in the truncated channel. Alternatively, the truncated channel may associate with intracellular factors that provide a Ca^{2+} sensor.

b. Coupling between Ca^{2+} binding and channel activation—In the absence of Ca^{2+} , BK channel gating involves two types of conformational changes, the activation of voltage sensors and the opening of the activation gate (Section IIIA–b). Ca^{2+} binding primarily promotes gate opening, with a small facilitating effects on voltage sensor activation. The effects of Ca^{2+} on gate opening can be measured at negative voltages where voltage sensors are at the resting state; under such conditions, the P_O increased $\geq 2,000$ times between 0 and 100 μM $[\text{Ca}^{2+}]_i$ [137,179]. Thus, voltage and Ca^{2+} activate BK channels using two parallel mechanisms, with the voltage sensors and the Ca^{2+} binding sites coupling to the activation gate independently, except for a weak interaction between the two mechanisms. Here we will review the coupling between Ca^{2+} binding and gate opening. The effect of Ca^{2+} binding on voltage sensor activation will be discussed in section VI “Relationship between activation pathways”.

The relationship between BK channel open probability and Ca^{2+} concentration can be fitted by the Hill equation, with the Hill coefficient ranging from 1.5 to 6 [133,160,162,180–183]. The Hill equation is a semi-empirical relationship that describes cooperative association of ligands by multiple binding sites. The values of the Hill coefficient for Ca^{2+} dependent activation indicate that there are at least 2–6 Ca^{2+} binding sites in a BK channel molecule, which is consistent with our current knowledge that each Slo1 subunit contains two Ca^{2+} binding sites (see above), and that, perhaps more importantly, there is a positive cooperativity among these Ca^{2+} binding sites. Such a cooperative interaction among Ca^{2+} binding sites in

BK channels can be explained by the MWC model that was originally proposed by Monod, Wyman, and Changeux [184] to account for the cooperative binding of oxygen to hemoglobin (Fig. 3B) [133,179,185]. The central idea of the MWC model is that ligand binding to any of the multiple binding sites will promote a conformational change that alters all binding sites and thus the affinity of ligand binding. In the MWC model for BK channels, the conformational change of the activation gate in channel opening (C-O) is accompanied by a conformational change of the Ca^{2+} binding sites that would increase the affinity of Ca^{2+} binding ($K_{\text{dO}} < K_{\text{dC}}$), giving rise to the positive cooperativity. Because the open state has a higher Ca^{2+} affinity than the closed state, the Ca^{2+} -bound open state is energetically more favorable than the Ca^{2+} -bound closed state with the same number of Ca^{2+} ions. Therefore, Ca^{2+} binding promotes channel opening, and the C-O transition is altered by factor C ($C = K_{\text{dO}}/K_{\text{dC}}$) for the binding of each additional Ca^{2+} (Fig. 3B).

Properties of BK channels are consistent with the key features of the MWC model. First, BK channels can open without binding to Ca^{2+} . Although earlier studies had observed BK channel activities with low open probability at very low Ca^{2+} concentrations [20,40,41,186,187], this property was explicitly demonstrated only after the cloning of Slo1. Expression of cloned BK channels in *Xenopus* oocytes and mammalian cells made it possible to study macroscopic BK channel currents at extremely low Ca^{2+} concentrations (0.5 ~ 100 nM). It was shown that at 0.5 nM Ca^{2+} , mSlo1 channels open with a rate of at least 200 times faster than the diffusion limit for Ca^{2+} binding; thus the channels must open prior to Ca^{2+} binding to the channel [133]. In addition, the P_{O} or activation rate of the channel does not change with Ca^{2+} concentration in the range between 0.5–100 nM [132,133], consistent with the idea that the channel opens without sensing the low Ca^{2+} concentrations. Furthermore, in the low Ca^{2+} concentrations, the channel can be activated to its maximum P_{O} [133], demonstrating that channel opening is not limited by Ca^{2+} binding. Second, channel opening follows a two-state model (C-O) at different Ca^{2+} concentrations. This is supported by the activation and deactivation kinetics that are well fitted with an exponential time course, after a brief delay due to voltage sensor activation, over a wide range of voltages (–350 – + 300 mV) and Ca^{2+} concentrations (0 – 1000 μM) [133,143,179]. Single channel measurements revealed multiple open and closed states and the transitions among open or closed states are dependent on Ca^{2+} concentration [146,180,181,183], consistent with the two-tiered MWC model (Fig. 3B). Third, the Hill coefficient of Ca^{2+} dependent activation depends on voltage, increasing with membrane depolarization [133,160,180]. This indicates that the cooperativity among Ca^{2+} binding sites increases with voltage, even though Ca^{2+} binding itself may not be voltage dependent and the interaction between Ca^{2+} binding and voltage sensor activation is weak [133,179]. This result is consistent with the idea that, in conjunction with channel opening, the conformation of the Ca^{2+} binding site changes to increase Ca^{2+} affinity: at more depolarized voltages the open probability is higher and thus it is more likely for Ca^{2+} binding sites to change conformation. More discussion on allosteric mechanisms of Ca^{2+} dependent activation of BK channels may be found in an insightful review by Karl Magleby [95].

Studies on the molecular mechanism of the coupling between Ca^{2+} binding and gate opening in BK channels have been predominately influenced by the studies of MthK from the laboratories of MacKinnon and Jiang. A recent review by Christopher Lingle has provided an excellent account of these studies [97]. Here we will briefly summarize the key features of Ca^{2+} dependent activation of MthK channels and then compare these to the results obtained in BK channel studies. The gating ring of the MthK channel is comprised of eight identical RCK domains (Fig. 2C), 4 of which derive from the intact MthK subunit and covalently link to the tetrameric pore with short peptides, while the additional 4 RCK domains are generated from the *MthK* gene by a second start site downstream from the pore domain. The Ca^{2+} binding site in each RCK domain is strategically located adjacent to the hinge of the flexible interface between the neighboring subunits, allowing Ca^{2+} to influence its conformation [110]. Ca^{2+}

binding changes the flexible interface such that the two RCK domains open like a clamshell [188]. As the result of the changes in the flexible interface, the gating ring expands to increase in diameter [188]. It is proposed that such an expansion will directly pull the pore open through the peptide linker between the pore and the RCK domains [110,188].

In the following, we compare BK and MthK channels in terms of the location of Ca^{2+} binding sites, the location and role of the flexible interface, and the coupling between the cytoplasmic domain and the pore. Differing from the MthK channel, the two putative RCK domains in BK channels are not identical. Consequently, the Ca^{2+} binding sites and flexible interfaces differ in the RCK1 and RCK2 domains of BK channels.

1. Ca^{2+} binding sites. In MthK, the Ca^{2+} binding site is located at the C-terminal end of βD (D184) & βE (E210 & E212) that places the bound Ca^{2+} at the hinge of the flexible interface. In contrast, in the RCK1 of mSlo1, D362 (at the C-terminus of αA) and D367 (at the αA - βB loop) are located at the opposite side of the β -sheets, far away from the hinge of the putative flexible interface. M513 (C-terminus of αG), however, is located at the expected flexible interface. In the homology model, the side chain of D362 makes direct contact with that of M513 [177], suggesting that the binding of Ca^{2+} to the RCK1-related site may be able to affect the putative flexible interface. The location of the Ca^{2+} bowl in relation to the putative RCK2 domain differs in published alignments (Table 1), and there is no data to suggest whether Ca^{2+} binding to the Ca^{2+} bowl would affect the flexible interface.
2. The flexible interface. In the RCK1 of mSlo1, residues 487–498 and 501–511 correspond to helices αF and αG , respectively, which are part of the flexible interface in the MthK channel (Table 1). However, no experimental data have been published to show whether these residues are part of an interface between structural domains or if they are important for Ca^{2+} dependent gating. The residues corresponding to αF and αG in the RCK2 of mSlo1 differ among published alignments (Table 1), and similarly, no experimental data is available to verify the prediction of their involvement in any interactions with the RCK1 domain or in Ca^{2+} dependent activation. Kim et al. found a mutation in RCK2 that altered the gate opening in the absence of Ca^{2+} binding and voltage sensor activation (Table 1) [125]. This residue, located between the putative βA and αA , was proposed to be part of the flexible interface with RCK1 [125]. However, this mutation did not affect Ca^{2+} dependent activation, thus its role in Ca^{2+} dependent activation may not be compatible with that of the flexible interface in MthK.
3. The coupling between the cytoplasmic domain and the pore. The S6 transmembrane segment of BK channels is connected to the RCK1 domain with a peptide linker of 16 amino acids. Niu et al. found that changes to the linker length by either deleting or adding different number of amino acids alter channel open probability both in the absence or presence of Ca^{2+} [154,177]. Shortening the linker increases channel activity and lengthening the linker decreases channel activity. This result is consistent with the hypothesis proposed for the MthK channel [110] that the cytoplasmic domain tugs the linker during Ca^{2+} dependent activation.

While the mechanism of Ca^{2+} dependent activation in MthK is a prevalent model for the coupling between Ca^{2+} binding and gate opening in BK channels, some results have suggested features of Ca^{2+} -dependent regulation of BK gating that may be outside the framework of the MthK model. Krishnamoorthy et al. found that the dSlo1 BK channel has a higher Ca^{2+} sensitivity in Ca^{2+} dependent gating than mSlo1 [189]. By studying the chimeras between the two homologous channels, the N-terminal half of the RCK1 corresponding to βA - αC (thus named the AC region) was identified to be responsible for the phenotypical difference. A molecular dynamics simulation based on the structure of the MthK RCK domain suggested

that the AC region of dSlo1 has a more tightly packed structure and less flexible dynamics than that of mSlo1. These results suggest that, unlike a rigid-body motion of the entire RCK domain in the clamshell model of MthK channels, conformational changes of a subdomain in RCK may be important for Ca^{2+} dependent gating in BK channels. Interestingly, it has been shown that the AC region is located close to the membrane spanning domain, and the two domains make multiple physical contacts [118,129]. These results suggest that, in addition to the direct pulling through the peptide linker between S6 and RCK1, the interactions among residues in the membrane spanning domain and the AC region may also contribute to the coupling between Ca^{2+} binding and channel opening.

C. Mg^{2+} -dependent activation of BK channels

a. Mg^{2+} binding sites— Mg^{2+} effects on BK channel activation were discovered in the late 1980's [190,191]. However, until 2001, it was not clear whether Mg^{2+} activates the channel by modulating Ca^{2+} dependent activation or via a separate mechanism. In 2001, two independent studies of the cloned BK channel illustrated that millimolar Mg^{2+} activates the BK channel via a low-affinity metal binding site that is independent from Ca^{2+} dependent activation [80,81]. One of the studies [80] measured BK channel activities in a broad range of Ca^{2+} and Mg^{2+} concentrations and found that the results could be fitted by a model including a high-affinity Ca^{2+} binding site and a lower affinity Mg^{2+} binding site. The other study showed that at both zero and saturating Ca^{2+} concentration (110 μM), Mg^{2+} activated the channel similarly with a binding affinity in the millimolar range, indicating that Ca^{2+} binding was not involved in Mg^{2+} dependent activation [81]. In addition, by comparing the chimera channels between the Mg^{2+} -sensitive mSlo1 and Mg^{2+} -insensitive mSlo3, Shi and Cui found that the low-affinity Mg^{2+} binding site resided at a different location from the Ca^{2+} bowl [81], demonstrating that the low and high-affinity metal binding sites are different.

Characterization of more chimera channels between mSlo1 and mSlo3 further pinpointed the Mg^{2+} binding site to the N-terminus of the RCK1 domain [121]. Three residues in this region, E374, E399 and Q397, were proposed as putative Mg^{2+} coordination sites based on systematic mutagenesis and structural simulation [121] of a homology model based on the RCK domain of the *E. coli* K^{+} channel [111]. At the same time, the Lingle group independently identified E399 as an essential residue for millimolar $\text{Mg}^{2+}/\text{Ca}^{2+}$ binding [122], thereby clearly separating the contribution of this putative low affinity $\text{Mg}^{2+}/\text{Ca}^{2+}$ binding site from that of the other two putative high affinity Ca^{2+} binding sites, the Ca^{2+} bowl and the D362/D367 site, to channel activation [122]. Further mutagenesis studies suggest that the side chains of E374 and E399 may coordinate Mg^{2+} , while the side chain of Q397 does not. Instead, Q397 is located close to the binding site and affects Mg^{2+} binding [128]. This conclusion was largely based on two lines of evidence. First, the carboxylate groups of E374 and E399 are required for Mg^{2+} sensing. Other side chains on either of these two residues completely abolished Mg^{2+} sensing. Second, adding positive charges to residue 397 by mutagenesis or chemical modification reduced but did not abolish Mg^{2+} binding. On the other hand, adding negative charge to residue 397 increased Mg^{2+} sensitivity. The opposite charge effects on Mg^{2+} sensing indicated that Q397 is close to the Mg^{2+} binding site and charges at this residue affect Mg^{2+} sensitivity through electrostatic interaction with the bound Mg^{2+} .

Owing to its chemical properties, Mg^{2+} is predominantly coordinated by six oxygen atoms from the side chains of oxygen-containing residues, main chain carbonyl groups in proteins, or water molecules [168]. Besides E374 and E399, are there any other oxygen-containing residues that might participate in Mg^{2+} binding? A recent study [118] addressed this question. Based on the homology model of the BK channel RCK1 domain, E374 and E399 are located on the top surface of the RCK1 domain, spatially close to the VSD [118,128,129]. Since the octahedral geometry of Mg^{2+} binding site constrains the distances among its coordinates, the

authors postulated that other protein ligands, if there are any, can only come from the top surface of the RCK1 domain and/or the intracellular portion of the membrane spanning domain. Through a systematic mutagenesis scanning of the oxygen-containing residues in these potential regions, the side chain oxygens of two residues in the voltage sensor domain were found essential for Mg^{2+} binding. These residues are D99 at the C-terminus of the S0–S1 loop and N172 in the S2–S3 loop.

Attribution of D99 as another Mg^{2+} coordinate is based on the following evidence [118]. First, the mutations of D99 that remove side chain oxygen completely abolished Mg^{2+} sensitivity, while the mutations that preserve oxygen with carbonyl or carboxylate groups retained partial Mg^{2+} sensitivity. In addition, these mutations specifically affected Mg^{2+} sensing without changing voltage- or Ca^{2+} -dependence. Second, the D99A mutation abolished the Mg^{2+} effects, but did not abolish channel activation by a positive charge that was covalently added in the vicinity of the Mg^{2+} binding site to mimic Mg^{2+} in activating the channel. Thus, D99A only abolished Mg^{2+} binding but not the coupling mechanism that opens the channel. Third, the formation of a disulfide bond between two cysteine residues that replace D99 and Q397 indicated that D99 in the voltage sensor domain is located close to the Mg^{2+} binding site in the RCK1 domain and thus can be part of the binding site. Evidence also suggested the contribution of N172 to Mg^{2+} coordination. Mutations of N172 to positively charged residues (R or K) abolished Mg^{2+} sensitivity; while mutations to negatively charged residues (D or E) increased Mg^{2+} sensitivity. More importantly, mutation N172D rescued Mg^{2+} sensitivity that had been abolished by mutating other Mg^{2+} binding residues (D99A, E374A, or E399N), indicating that the carboxylate group on the side chain of residue 172 may directly contribute to Mg^{2+} coordination to compensate for the loss of coordination at other positions. These studies indicate that the Mg^{2+} binding site may be comprised of residues from two different domains: D99 and N172 from the VSD, and E374 and E399 from the cytoplasmic RCK1 domain.

Yang et al. further demonstrated that in each Mg^{2+} binding site D99/N172 may not come from the same subunit as E374/E399, but rather from a neighboring subunit [118]. While individual mutations D99R, N172R, E374R, or E399C abolished Mg^{2+} sensitivity, hybrid channels resulting from the co-expression of the single mutations from different domains (for example, D99R from VSD and E374R from RCK1) retained some Mg^{2+} sensitivity. The retention of Mg^{2+} sensitivity cannot be explained by an intra-subunit binding site model in which all four Mg^{2+} binding residues come from the same subunit because in this model, each Mg^{2+} binding site would contain a mutation that abolished Mg^{2+} sensitivity. Instead, this result suggests that D99/N172 in the VSD and E374/E399 in the RCK1 domains from neighboring subunits form an inter-subunit Mg^{2+} binding site. Thus, according to binomial distribution of the single mutations in the tetrameric channels, some of the Mg^{2+} binding sites may contain two mutations, leaving other Mg^{2+} binding sites intact. The retained Mg^{2+} sensitivity measured in experiments can be fitted well with the contribution from these intact sites. Therefore, the Mg^{2+} binding site of BK channels may be comprised of D99/N172 in the VSD of one subunit and E374/E399 in the RCK1 domain of the neighboring subunit.

Besides this low affinity divalent cation (Mg^{2+}/Ca^{2+}) binding site in the RCK1 domain, BK channels may contain divalent cation binding sites with even lower affinity ($K_d \sim 40$ mM), termed as very low affinity sites [166]. These very low affinity sites might be responsible for the additional shifts of the G–V relation induced by increasing $[Mg^{2+}]_i$ or $[Ca^{2+}]_i$ from 10 to 100 mM [80,122,166], partially because the shifts are too large to be solely attributed to surface charge screening [166]. In addition, these very low affinity sites are distinct from the two high affinity Ca^{2+} and the low affinity Mg^{2+} binding sites because mutations that eliminate the three binding sites do not affect the G–V shifts induced by high concentrations of Ca^{2+} or Mg^{2+} [166]. A recent study further illustrated that these very low affinity sites activated BK channels allosterically by affecting both the opening of the activation gate (L) and voltage sensor

movement (J) (Fig. 3A) [192]. At present, the location, number or physiological significance of the very low affinity sites is still not clear. However, importantly, at Mg^{2+} concentrations around 10 mM the contribution of these very low affinity sites to channel activation overlaps with that of the Mg^{2+} binding site in RCK1, which interferes with the study of Mg^{2+} dependent activation through the Mg^{2+} binding site in RCK1. Nevertheless, the separation of the activation effects by these distinct sites can be achieved by a simple subtraction [118,128,129,166] because studies show that these sites affect channel activation independently [166].

b. Coupling between Mg^{2+} binding and channel activation—Similar to Ca^{2+} dependent activation of BK channels, millimolar Mg^{2+} increases open probability and shifts the G - V relation to more negative voltages [80,81]. Likewise, these properties can also be described by an allosteric MWC gating model. However, unlike micromolar Ca^{2+} that accelerates activation rate and reduces deactivation rate of the channel, millimolar Mg^{2+} only reduces deactivation rate but has no effect on activation rate [80,81,164,166,192], implying that the underlying molecular mechanism of Mg^{2+} -dependent activation may differ from that of Ca^{2+} -dependent activation.

Several recent studies further demonstrate that Mg^{2+} and Ca^{2+} indeed activate BK channels through distinct molecular mechanisms [129,140,192]. While Ca^{2+} activates the channel largely independent of the voltage sensor [179], Mg^{2+} activates the channel by an electrostatic interaction with the voltage sensor [129,192]. The involvement of the voltage sensor in Mg^{2+} -dependent activation is suggested by two lines of evidence. First, Mg^{2+} has no measurable effect on channel activation at negative voltages where voltage sensors are in the resting state [129,192]; in contrast, 70 μM [Ca^{2+}]_i can increase the open probability >2,000-fold at similar voltages [179]. This result suggests that Mg^{2+} activates the channel only when the voltage sensor can be activated. Second, neutralization of R213, a voltage-sensing residue at the C-terminus of S4, specifically eliminated Mg^{2+} sensitivity, but did not affect Ca^{2+} sensing [140]. This result suggests that the voltage sensor, more specifically R213, is important for Mg^{2+} dependent activation. Recently, it was shown that Mg^{2+} activated the channel by an electrostatic repulsion of R213 [129] based on the following four lines of evidence. 1) Altering ionic strength of intracellular solutions specifically changed Mg^{2+} sensing but had little effect on Ca^{2+} sensing, indicating that Mg^{2+} -dependent activation is electrostatic by nature. 2) A positive charge at residue 397, introduced either by mutagenesis or chemical modification, mimicked Mg^{2+} effects on channel activation. Similar to Mg^{2+} , effects of this positive charge were also sensitive to ionic strength. Since residue 397 is in the vicinity of the RCK1 Mg^{2+} binding site, these results further support the involvement of electrostatic interaction in Mg^{2+} -dependent activation. 3) 10 mM [Mg^{2+}]_i reduced the amplitude and slowed the relaxation rate of off gating currents (I_{gOFF}), indicating that Mg^{2+} stabilizes the voltage sensor in its active state. 4) R213 was found to be the only charged residue in the VSD that sensed the electric field from Mg^{2+} or charges at position 397 that is close to the RCK1 Mg^{2+} binding site. Adding back a positive charge to R213C, a mutation that abolishes Mg^{2+} sensitivity, by chemical modification, rescued partial Mg^{2+} sensitivity. This result indicates that a positive charge at position 213 is necessary and sufficient for Mg^{2+} to activate the channel through an electrostatic repulsion.

Interestingly, while Mg^{2+} affects I_{gOFF} significantly, it has only a small effect on the on gating currents (I_{gON}) [129]. Since I_{gON} reflects voltage sensor activation when the channels are closed and I_{gOFF} reflects the return of the voltage sensor from the active state to the resting state when channels are open, this result suggests that the electrostatic interaction between Mg^{2+} and R213 is state-dependent, i.e., the interaction is much stronger when the channel is open than closed. A fitting of the HCA model to the experimental data showed that the equilibrium of voltage sensor movements at the open state (V_{hO}) was affected by Mg^{2+} much more than at the closed state (V_{hC}), which means that Mg^{2+} strengthens the allosteric coupling

between voltage sensor and the activation gate (described by the allosteric factor D , see Fig. 3B) [129]. Likewise, in a recent study, Horrigan and Ma evaluated the biophysical mechanism of the electrostatic interaction and attributed Mg^{2+} effects to the ability of Mg^{2+} to stabilize channels in states with both activated voltage sensors and open gates (open-activated states or OA states) [192]. The stabilization of the OA state by Mg^{2+} results in the enhancement of the coupling between voltage sensor and channel activation (the allosteric factor D), thereby facilitating channel activation. The state dependent interaction suggests that the distance between R213 and Mg^{2+} differs between the open and the closed states. In other words, the cytoplasmic domain and the membrane-spanning domain may undergo a shift in their relative positions during channel opening.

IV. Relationship between Activation Pathways

A. Ca^{2+} - and voltage-dependent activation

Although distinct mechanisms underlie Ca^{2+} - and voltage-dependent activation of BK channels, it is apparent that the G–V relation shifts to more negative voltages with increasing $[Ca^{2+}]_i$ (Fig. 1A), while the G– $[Ca^{2+}]_i$ dose response curve is altered by increasing voltage to decrease EC_{50} and increase the Hill coefficient [133]. These changes, however, are primarily mediated by channel opening, rather than a direct effect of Ca^{2+} on the voltage dependent mechanism or vice versa. First, both voltage and Ca^{2+} can activate the channel independently to increase P_o . Therefore, P_o at the same voltage is larger with higher $[Ca^{2+}]_i$, and at the same $[Ca^{2+}]_i$ is larger with a more depolarized voltage. Second, due to the allosteric coupling of the activation gate with the voltage sensor and the Ca^{2+} binding sites (see Section III), channel opening facilitates voltage sensor activation as well as Ca^{2+} binding (Fig. 3). Thus, the opening of the channel due to one stimulus (Ca^{2+} or voltage) will enhance the effect of the other via the allosteric coupling between the activation gate and the VSD or Ca^{2+} binding sites. These indirect interactions mediated by channel opening account for the majority of the Ca^{2+} -dependent G–V shift and the voltage dependent G– $[Ca^{2+}]_i$ shift [179,185]. In addition, there is a weak direct interaction between voltage sensor and Ca^{2+} that is not mediated by channel opening that is described by an allosteric factor E (Fig. 3B). This interaction was shown by Horrigan and Aldrich by recording the fast component of on gating currents (I_{gfast}) at 0 and 70 μM $[Ca^{2+}]_i$ [179]. During their recordings, the channels were all closed so that the Ca^{2+} dependence of I_{gfast} reflected a direct interaction between Ca^{2+} binding and voltage sensor movement that is independent of the ability of Ca^{2+} to alter channel opening. They found that the Q_{fast} -V relation shifted by -20 mV in 70 μM $[Ca^{2+}]_i$ as compared to that in 0 $[Ca^{2+}]_i$. This shift of the Q_{fast} -V relation was mainly due to the reduction in the rate of voltage sensor returning from the active state to the resting state [179].

B. Ca^{2+} - and Mg^{2+} - dependent activation

While the high-affinity Ca^{2+} binding sites in BK channels favor Ca^{2+} , other divalent cations at higher concentration can bind to these sites [164,191]. It was shown that Mg^{2+} competes with Ca^{2+} for the Ca^{2+} binding sites, with an affinity of ~ 5 mM [80,81,166]. However, the binding of Mg^{2+} to high affinity Ca^{2+} binding sites does not activate the channel; thus the net effect of such binding is to reduce Ca^{2+} dependent activation at low Ca^{2+} concentrations [80, 81]. The competition of Mg^{2+} to the Ca^{2+} binding sites was more clearly demonstrated in the study of mutant channels, in which the Mg^{2+} binding site was abolished so that Mg^{2+} no longer activated the channel. Thus, the G–V relation at low Ca^{2+} concentrations shifted to more positive voltage ranges upon the addition of 10 mM Mg^{2+} , opposite to the effect of Mg^{2+} in wild type BK channels (Fig. 1) [166]. It is not known whether Mg^{2+} selectively binds to the Ca^{2+} bowl or the RCK1-related Ca^{2+} binding site. On the other hand, the low-affinity Mg^{2+} binding site at the interface of the VSD and the RCK1 domain is not selective for Ca^{2+} or Mg^{2+} ions. Like Mg^{2+} , Ca^{2+} at millimolar concentrations binds to this low affinity site to

activate the channel [80,81,122]. This site is called the Mg^{2+} binding site because it binds to Mg^{2+} around the physiological concentration of intracellular Mg^{2+} [193].

The molecular mechanisms of Mg^{2+} coupling to the activation gate do not interfere with those of Ca^{2+} . The effect of millimolar $[Mg^{2+}]_i$ on channel activation is not altered regardless of the presence of Ca^{2+} [80,81], and the effects of saturating $[Ca^{2+}]_i$ and millimolar $[Mg^{2+}]_i$ on activation are largely additive [122]. Additionally, mutations that reduce or abolish Mg^{2+} sensitivity usually do not affect Ca^{2+} sensitivity [118,121,122,128,129,140,166]. Similarly, mutations that reduce Ca^{2+} sensitivity generally have little effect on Mg^{2+} dependent activation [121,122]. These results are remarkable since the putative Mg^{2+} and one of Ca^{2+} binding sites are both in the RCK1 domain and close to each other. In the homology model of the RCK1 domain in BK channels, E374 is located at the N-terminus of βB for Mg^{2+} binding, while D367, only 7 amino acids away in the primary sequence, is located in the loop connecting the N-terminus of βB and αA for Ca^{2+} dependent activation. These results illustrate an important feature in BK channel activation, i.e., each metal ion perturbs only a subset of amino acid residues to affect channel gating; these perturbations do not overlap perhaps because, as discussed above, Mg^{2+} activation is mediated via the interaction with the voltage sensor and Ca^{2+} activation is through forces applied to the linker between the gate and the cytoplasmic domain.

V. Activation Gate

Energy provided by voltage, Ca^{2+} and Mg^{2+} binding will propagate to the activation gate of BK channels to initiate ion conduction through the pore. In canonical K^+ channels, the selectivity filter with the signature “GYG” sequence determines ion selectivity and permeation [102,103,194], the hydrophobic residues at the C-terminus of S6 (or inner helix) form the inner mouth of the pore to restrict K^+ ion flux when channels are closed [195], and the Gly hinge in the middle of S6 [196] or the Pro kink(s) [197] at the C-terminus of S6 swings the gate to open upon stimulation. However, several unique features of the BK channel indicate that, while the selectivity filter of the BK channel may be highly conserved, its pore and activation gate may differ somewhat from this picture.

First, BK channels exhibit much larger single channel conductance (~100–300 pS) than other K^+ channels while still maintaining high selectivity to K^+ ions. Several elegant studies indicated that rings of negatively charges at the intracellular entrance of the S6 (composed by eight Glu residues) [198,199] and the extracellular outer pore [200,201] of BK channels partially determine the large conductance of BK channels. These negatively charged rings attract K^+ ions, resulting in several fold increase in local $[K^+]$ [199], thereby speeding up K^+ flux. In addition, a large inner vestibule and a wide intracellular entrance to the vestibule might also contribute to the large conductance. In a series of experiments conducted by the Magleby group [202] and the Aldrich group [203–205], chemicals with various sizes were used to probe the size of the inner vestibule and the cytoplasmic entrance of the pore. Based on the changes of the K^+ diffusion rate from bulk intracellular solution to the inner vestibule induced by sucrose interference, the inner mouth of BK channels was estimated to be twice (~20Å) as large as that of the Shaker K^+ channel [202]. Consistently, quaternary ammonium (QA) ions, such as tetrabutylammonium (TBA) and decyltriethylammonium (C_{10}), showed much faster block and unblock kinetics in BK channels than in canonical K_V channels. The time-independent blockade indicates that BK channels may have an enlarged inner vestibule and a widened intracellular entrance [203,205] so that large QA ions can access their binding site from the cytoplasmic solution with less restriction. All these features together may contribute to the large conductance of BK channels.

Second, different from canonical K^+ channels, the inner mouth of the closed BK channel pore may not obstruct small hydrated ions, such as K^+ and QA ions. Instead, the permeation gate may be near or in the selectivity filter, similar to several other ligand-gated ion channels such as CNG (cyclic nucleotide-gated) channels [206] and SK (small conductance, Ca^{2+} -activated K^+) channels [207]. The Aldrich group showed that C_{10} and TBA did not slow deactivation kinetics of the gate, indicating that the closure of the BK channel permeation gate was not hindered by the presence of these blockers. These experiments excluded the classic open-channel block mechanism (“foot in the door”), but left an open question on whether the blockers are trapped inside the inner vestibule when the gate closes (the trapping model), or can freely access their binding site inside the vestibule regardless of the state of the gate (the state-independent free access model). These two possibilities could not be distinguished using these QA ions due to their very fast block/unblock kinetics. Then a different strategy was devised by Willkens and Aldrich who used a TBA derivative containing a bulky benzoyl-benzoyl group (bbTBA) to study the steady-state and kinetic behavior [205]. Different from TBA and C_{10} , bbTBA blocks BK channels in a time-dependent fashion. The slow block kinetics allowed the investigators to study bbTBA blockade using a classic trapping experiment protocol. Their experiments showed no evidence that bbTBA was trapped inside the channel pore after gate closing. Thus, bbTBA blocks BK channels independent of gate opening and closing. Together with other careful characterizations, the authors concluded that BK channels might employ their selectivity filter as the permeation gate so that both K^+ ions and small QA ions can have relatively free access to the internal vestibule independent of channel states [205]. Consistent with the selectivity filter being the ion permeation gate, Piskorowski and Aldrich proposed that the interaction between the selectivity filter and permeating ions is critical for BK channel opening based on the effects of permeating thallium ion (Tl^+) on channel activation [208]. The authors suggested that permeating Tl^+ may stabilize a collapsed state of the selectivity filter; thereby shifting the activation curve to more positive voltages and increasing the flicker frequency at the single channel level [208].

Third, the movement of the pore-lining helix (S6) of BK channels might be different from canonical K^+ channels. Although BK channels may employ their selectivity filter as the permeation gate, the S6 helix may still move during channel gating. Li and Aldrich found that a Shaker ball peptide (ShBP) homologue, which has a larger size than QA ions, blocked BK channels in a state-dependent fashion, i.e., ShBP blockade depended on the activation kinetics and open probability of the channel [204]. In addition, ShBP blockade slowed the deactivation kinetics without affecting the onset of the activation kinetics [204]. Thus, the cytoplasmic entrance of the BK channel pore may indeed move from the open to the closed state to restrict the entry of large ShBP molecules but not smaller K^+ and QA ions. In another study, Guo et al. found that hydrophobicity of a residue (323 in mSlo1) at the inner mouth of the BK channel pore is important for channel gating [209]. All hydrophilic mutations of residue 323 exhibited increased subconductance levels, while the hydrophobic mutations showed prolonged open duration. The authors proposed that a reduced hydrophobicity at residue 323 would reduce the cooperativity among the four subunits of the channel in opening the activation gate, thus increasing the chance of partial channel openings that result in subconductance levels. These results are also consistent with the idea that the inner mouth of the BK channel is involved in the opening of the activation gate although the gate for ion permeation may reside in the selectivity filter.

Structural and functional studies of K^+ channels suggest that the highly conserved “Gly hinge” at the middle of the pore-lining helix [196] and/or the Pro-Val-Pro (PVP motif) at the C-terminus [197] are two important pivot points for the movement of the activation gate. In BK channels, the PVP motif is substituted by YVP and there are two adjacent Gly residues at the position of the “Gly hinge.” Single Ala mutations to either one of these Gly residues shifted the G-V relation similarly to more positive voltages for about +70 mV in 100 μM $[Ca^{2+}]_i$, while

the double Ala mutation shifted almost twice as much as the single mutations [210], suggesting that the flexibility around the “Gly hinge” is critical to the gating of BK channels. It is worth noting that in Shaker K⁺ channels, an Ala mutation of the “Gly hinge” completely abolished gate opening without affecting the membrane trafficking of the mutant channel [211]. This is different from the Ala mutations of the “Gly hinge” in BK channels, as mutant channels can still be opened by voltage and intracellular Ca²⁺. In addition, altering the side chain of a phenylalanine residue (F315 in mSlo1) four amino acids after the “Gly hinge” changes the gating properties of BK channels [212,213]. F315Y greatly facilitates BK channel opening, while F315I extensively decreases channel open probability. This phenylalanine residue is not conserved in K_V channels. All the above evidences suggest that the physical motions of the activation gate in BK channels might differ from those in K_V channels. Further studies are needed to elucidate their differences.

VI. Conclusion remarks

Keeping true to their name, BK channels have a large single channel conductance and a large molecular mass among K⁺ channels. The massive structure contains sensors to a variety of cellular signals including phosphorylation [86,89], oxidation [214–218], CO [165], H⁺ [130, 219], Heme [89,123,220], and PIP₂ [221] in addition to voltage, Ca²⁺ and Mg²⁺. BK channels respond to these stimuli by changing the P_O, thus integrating cellular signals to modulate excitability and Ca²⁺ homeostasis. The BK channel molecule is comprised of the membrane spanning pore-gate domain and the voltage sensing domain as well as a large cytoplasmic domain. The studies of channel activation in response to voltage, Ca²⁺ and Mg²⁺ reveal that the interactions among these structural domains are crucial in mediating the stimulation to channel opening. These findings provide principles that may also govern channel gating in response to other stimuli. In addition, the function of BK channels is modulated by its association with the accessory β subunits, which are important for the phenotypic differences of BK channels in various tissues [39,90]. Some of the β subunits modulate voltage, Ca²⁺ or Mg²⁺ sensitivity of BK channels [107,132,145,175,222–233] but the molecular mechanisms of such modulation remain unclear. The different mechanisms of BK channel gating reviewed here may provide a basis for further investigation in this area.

BK channels share structural and functional similarities with K_V channels and ligand activated K⁺ channels. Common principles of ion channel gating apply to these channels. However, the BK channel is unique in that it combines the structure and function of both K_V channels and ligand-gated K⁺ channels. Thus, the gating by one stimulus can be tuned by the change in another stimulus. Elucidation of the effects of voltage, Ca²⁺ and Mg²⁺ on BK channels has clearly demonstrated that allosteric mechanisms are fundamental to the regulation of BK gating. Although in other ion channels such allosteric couplings between distinct domains of the channel protein may not be readily observable, it would not be surprising if similar allosteric mechanisms may underlie gating in all ion channels.

Acknowledgments

We thank Chris Lingle (Washington University, St. Louis, MO) for critical discussions. This work was supported by National Institutes of Health Grant R01-HL70393 (to J.C.). J.C. is an associate professor of Biomedical Engineering on the Spencer T. Olin Endowment.

References

1. Latorre R, Miller C. Conduction and selectivity in potassium channels. *J Membr Biol* 1983;71:11–30. [PubMed: 6300405]
2. Latorre R, Oberhauser A, Labarca P, Alvarez O. Varieties of calcium-activated potassium channels. *Annu Rev Physiol* 1989;51:385–399. [PubMed: 2653189]

3. Marty A. Ca^{2+} -dependent K^{+} channels with large unitary conductance. *Trends Neurosci* 1983;6:262–265.
4. Adams PR, Constanti A, Brown DA, Clark RB. Intracellular Ca^{2+} activates a fast voltage-sensitive K^{+} current in vertebrate sympathetic neurones. *Nature* 1982;296:746–749. [PubMed: 6280066]
5. Faber ES, Sah P. Ca^{2+} -activated K^{+} (BK) channel inactivation contributes to spike broadening during repetitive firing in the rat lateral amygdala. *J Physiol (Lond)* 2003;552:483–497. [PubMed: 14561831]
6. Lancaster B, Nicoll RA. Properties of two calcium-activated hyperpolarizations in rat hippocampal neurones. *J Physiol (Lond)* 1987;389:187–203. [PubMed: 2445972]
7. Knaus HG, Schwarzer C, Koch RO, Eberhart A, Kaczorowski GJ, Glossmann H, Wunder F, Pongs O, Garcia ML, Sperk G. Distribution of high-conductance Ca^{2+} -activated K^{+} channels in rat brain: targeting to axons and nerve terminals. *J Neurosci* 1996;16:955–963. [PubMed: 8558264]
8. Maue RA, Dionne VE. Patch-clamp studies of isolated mouse olfactory receptor neurons. *J Gen Physiol* 1987;90:95–125. [PubMed: 2442298]
9. Neely A, Lingle CJ. Two components of calcium-activated potassium current in rat adrenal chromaffin cells. *J Physiol (Lond)* 1992;453:97–131. [PubMed: 1464851]
10. Neely A, Lingle CJ. Effects of muscarine on single rat adrenal chromaffin cells. *J Physiol (Lond)* 1992;453:133–166. [PubMed: 1464826]
11. Solaro CR, Prakriya M, Ding JP, Lingle CJ. Inactivating and noninactivating Ca^{2+} - and voltage-dependent K^{+} current in rat adrenal chromaffin cells. *J Neurosci* 1995;15:6110–6123. [PubMed: 7545225]
12. Marty A, Neher E. Potassium channels in cultured bovine adrenal chromaffin cells. *J Physiol (Lond)* 1985;367:117–141. [PubMed: 2414437]
13. Lingle CJ, Solaro CR, Prakriya M, Ding JP. Calcium-activated potassium channels in adrenal chromaffin cells. *Ion Channels* 1996;4:261–301. [PubMed: 8744211]
14. Ohmori H. Studies of ionic currents in the isolated vestibular hair cell of the chick. *J Physiol (Lond)* 1984;350:561–581. [PubMed: 6086899]
15. Hudspeth AJ, Lewis RS. Kinetic analysis of voltage- and ion-dependent conductances in saccular hair cells of the bull-frog, *Rana catesbeiana*. *J Physiol (Lond)* 1988;400:237–274. [PubMed: 2458454]
16. Fuchs PA, Nagai T, Evans MG. Electrical tuning in hair cells isolated from the chick cochlea. *J Neurosci* 1988;8:2460–2467. [PubMed: 3249237]
17. Roberts WM, Jacobs RA, Hudspeth AJ. Colocalization of ion channels involved in frequency selectivity and synaptic transmission at presynaptic active zones of hair cells. *J Neurosci* 1990;10:3664–3684. [PubMed: 1700083]
18. Issa NP, Hudspeth AJ. Clustering of Ca^{2+} channels and Ca^{2+} -activated K^{+} channels at fluorescently labeled presynaptic active zones of hair cells. *Proc Natl Acad Sci U S A* 1994;91:7578–7582. [PubMed: 8052623]
19. Latorre R, Vergara C, Hidalgo C. Reconstitution in planar lipid bilayers of a Ca^{2+} -dependent K^{+} channel from transverse tubule membranes isolated from rabbit skeletal muscle. *Proc Natl Acad Sci U S A* 1982;79:805–809. [PubMed: 6278496]
20. Barrett JN, Magleby KL, Pallotta BS. Properties of single calcium-activated potassium channels in cultured rat muscle. *J Physiol (Lond)* 1982;331:211–230. [PubMed: 6296366]
21. Pallotta BS, Magleby KL, Barrett JN. Single channel recordings of Ca^{2+} -activated K^{+} currents in rat muscle cell culture. *Nature* 1981;293:471–474. [PubMed: 6273730]
22. Inoue R, Kitamura K, Kuriyama H. Two Ca-dependent K-channels classified by the application of tetraethylammonium distribute to smooth muscle membranes of the rabbit portal vein. *Pflugers Arch* 1985;405:173–179. [PubMed: 2415914]
23. Toro L, Vaca L, Stefani E. Calcium-activated potassium channels from coronary smooth muscle reconstituted in lipid bilayers. *Am J Physiol* 1991;260:H1779–1789. [PubMed: 1711788]
24. McCobb DP, Fowler NL, Featherstone T, Lingle CJ, Saito M, Krause JE, Salkoff L. A human calcium-activated potassium channel gene expressed in vascular smooth muscle. *Am J Physiol* 1995;269:H767–777. [PubMed: 7573516]

25. Meredith AL, Thorneloe KS, Werner ME, Nelson MT, Aldrich RW. Overactive bladder and incontinence in the absence of the BK large conductance Ca^{2+} -activated K^{+} channel. *J Biol Chem* 2004;279:36746–36752. [PubMed: 15184377]
26. Kume H, Takai A, Tokuno H, Tomita T. Regulation of Ca^{2+} -dependent K^{+} -channel activity in tracheal myocytes by phosphorylation. *Nature* 1989;341:152–154. [PubMed: 2550823]
27. Kume H, Graziano MP, Kotlikoff MJ. Stimulatory and inhibitory regulation of calcium-activated potassium channels by guanine nucleotide-binding proteins. *Proc Natl Acad Sci U S A* 1992;89:11051–11055. [PubMed: 1438313]
28. Savaria D, Lanoue C, Cadieux A, Rousseau E. Large conducting potassium channel reconstituted from airway smooth muscle. *Am J Physiol* 1992;262:L327–336. [PubMed: 1372487]
29. Markwardt F, Isenberg G. Gating of maxi K^{+} channels studied by Ca^{2+} concentration jumps in excised inside-out multi-channel patches (myocytes from guinea pig urinary bladder). *J Gen Physiol* 1992;99:841–862. [PubMed: 1322449]
30. Storm JF. Action potential repolarization and a fast after-hyperpolarization in rat hippocampal pyramidal cells. *J Physiol (Lond)* 1987;385:733–759. [PubMed: 2443676]
31. Shao LR, Halvorsrud R, Borg-Graham L, Storm JF. The role of BK-type Ca^{2+} -dependent K^{+} channels in spike broadening during repetitive firing in rat hippocampal pyramidal cells. *J Physiol (Lond)* 1999;521(Pt 1):135–146. [PubMed: 10562340]
32. Hu H, Shao LR, Chavoshy S, Gu N, Trieb M, Behrens R, Laake P, Pongs O, Knaus HG, Ottersen OP, Storm JF. Presynaptic Ca^{2+} -activated K^{+} channels in glutamatergic hippocampal terminals and their role in spike repolarization and regulation of transmitter release. *J Neurosci* 2001;21:9585–9597. [PubMed: 11739569]
33. Faber ES, Sah P. Calcium-activated potassium channels: multiple contributions to neuronal function. *Neuroscientist* 2003;9:181–194. [PubMed: 15065814]
34. Gu N, Vervaeke K, Storm JF. BK potassium channels facilitate high-frequency firing and cause early spike frequency adaptation in rat CA1 hippocampal pyramidal cells. *J Physiol (Lond)* 2007;580:859–882. [PubMed: 17303637]
35. Robitaille R, Charlton MP. Presynaptic calcium signals and transmitter release are modulated by calcium-activated potassium channels. *J Neurosci* 1992;12:297–305. [PubMed: 1370323]
36. Robitaille R, Garcia ML, Kaczorowski GJ, Charlton MP. Functional colocalization of calcium and calcium-gated potassium channels in control of transmitter release. *Neuron* 1993;11:645–655. [PubMed: 7691106]
37. Ling S, Sheng JZ, Braun JE, Braun AP. Syntaxin 1A co-associates with native rat brain and cloned large conductance, calcium-activated potassium channels in situ. *J Physiol (Lond)* 2003;553:65–81. [PubMed: 12949219]
38. Cibulsky SM, Fei H, Levitan IB. Syntaxin-1A binds to and modulates the Slo calcium-activated potassium channel via an interaction that excludes syntaxin binding to calcium channels. *J Neurophysiol* 2005;93:1393–1405. [PubMed: 15496493]
39. Orio P, Rojas P, Ferreira G, Latorre R. New disguises for an old channel: MaxiK channel beta-subunits. *News Physiol Sci* 2002;17:156–161. [PubMed: 12136044]
40. Wong BS, Lecar H, Adler M. Single calcium-dependent potassium channels in clonal anterior pituitary cells. *Biophys J* 1982;39:313–317. [PubMed: 6291655]
41. Findlay I, Dunne MJ, Petersen OH. High-conductance K^{+} channel in pancreatic islet cells can be activated and inactivated by internal calcium. *J Membr Biol* 1985;83:169–175. [PubMed: 2582121]
42. Hudspeth AJ, Lewis RS. A model for electrical resonance and frequency tuning in saccular hair cells of the bull-frog, *Rana catesbeiana*. *J Physiol (Lond)* 1988;400:275–297. [PubMed: 2458455]
43. Art JJ, Wu YC, Fettiplace R. The calcium-activated potassium channels of turtle hair cells. *J Gen Physiol* 1995;105:49–72. [PubMed: 7730789]
44. Wu YC, Art JJ, Goodman MB, Fettiplace R. A kinetic description of the calcium-activated potassium channel and its application to electrical tuning of hair cells. *Prog Biophys Mol Biol* 1995;63:131–158. [PubMed: 7624477]
45. Art JJ, Fettiplace R. Variation of membrane properties in hair cells isolated from the turtle cochlea. *J Physiol (Lond)* 1987;385:207–242. [PubMed: 2443666]

46. Fuchs PA, Evans MG. Potassium currents in hair cells isolated from the cochlea of the chick. *J Physiol (Lond)* 1990;429:529–551. [PubMed: 2277357]
47. Mammano F, Bortolozzi M, Ortolano S, Anselmi F. Ca^{2+} signaling in the inner ear. *Physiology (Bethesda)* 2007;22:131–144. [PubMed: 17420304]
48. Knot HJ, Standen NB, Nelson MT. Ryanodine receptors regulate arterial diameter and wall $[\text{Ca}^{2+}]$ in cerebral arteries of rat via Ca^{2+} -dependent K^+ channels. *J Physiol (Lond)* 1998;508 (Pt 1):211–221. [PubMed: 9490841]
49. Brenner R, Perez GJ, Bonev AD, Eckman DM, Kosek JC, Wiler SW, Patterson AJ, Nelson MT, Aldrich RW. Vasoregulation by the beta1 subunit of the calcium-activated potassium channel. *Nature* 2000;407:870–876. [PubMed: 11057658]
50. Brayden JE, Nelson MT. Regulation of arterial tone by activation of calcium-dependent potassium channels. *Science* 1992;256:532–535. [PubMed: 1373909]
51. Ledoux J, Werner ME, Brayden JE, Nelson MT. Calcium-activated potassium channels and the regulation of vascular tone. *Physiology (Bethesda)* 2006;21:69–78. [PubMed: 16443824]
52. Nelson MT, Cheng H, Rubart M, Santana LF, Bonev AD, Knot HJ, Lederer WJ. Relaxation of arterial smooth muscle by calcium sparks. *Science* 1995;270:633–637. [PubMed: 7570021]
53. Herrera GM, Heppner TJ, Nelson MT. Regulation of urinary bladder smooth muscle contractions by ryanodine receptors and BK and SK channels. *Am J Physiol Regul Integr Comp Physiol* 2000;279:R60–68. [PubMed: 10896865]
54. Herrera GM, Etherton B, Nausch B, Nelson MT. Negative feedback regulation of nerve-mediated contractions by K_{Ca} channels in mouse urinary bladder smooth muscle. *Am J Physiol Regul Integr Comp Physiol* 2005;289:R402–R409. [PubMed: 15845880]
55. Petkov GV, Bonev AD, Heppner TJ, Brenner R, Aldrich RW, Nelson MT. Beta1-subunit of the Ca^{2+} -activated K^+ channel regulates contractile activity of mouse urinary bladder smooth muscle. *J Physiol (Lond)* 2001;537:443–452. [PubMed: 11731577]
56. Werner ME, Knorn AM, Meredith AL, Aldrich RW, Nelson MT. Frequency encoding of cholinergic- and purinergic-mediated signaling to mouse urinary bladder smooth muscle: modulation by BK channels. *Am J Physiol Regul Integr Comp Physiol* 2007;292:R616–624. [PubMed: 16931654]
57. Semenov I, Wang B, Herlihy JT, Brenner R. BK channel beta1-subunit regulation of calcium handling and constriction in tracheal smooth muscle. *Am J Physiol Lung Cell Mol Physiol* 2006;291:L802–810. [PubMed: 16632519]
58. Kotlikoff MI. Potassium channels in airway smooth muscle: a tale of two channels. *Pharmacol Ther* 1993;58:1–12. [PubMed: 8105495]
59. Jones TR, Charette L, Garcia ML, Kaczorowski GJ. Selective inhibition of relaxation of guinea-pig trachea by charybdotoxin, a potent Ca^{++} -activated K^+ channel inhibitor. *J Pharmacol Exp Ther* 1990;255:697–706. [PubMed: 1700817]
60. Du W, Bautista JF, Yang H, Diez-Sampedro A, You SA, Wang L, Kotagal P, Luders HO, Shi J, Cui J, Richerson GB, Wang QK. Calcium-sensitive potassium channelopathy in human epilepsy and paroxysmal movement disorder. *Nat Genet* 2005;37:733–738. [PubMed: 15937479]
61. Brenner R, Chen QH, Vilaythong A, Toney GM, Noebels JL, Aldrich RW. BK channel beta4 subunit reduces dentate gyrus excitability and protects against temporal lobe seizures. *Nat Neurosci* 2005;8:1752–1759. [PubMed: 16261134]
62. Sausbier M, Hu H, Arntz C, Feil S, Kamm S, Adelsberger H, Sausbier U, Sailer CA, Feil R, Hofmann F, Korth M, Shipston MJ, Knaus HG, Wolfer DP, Pedroarena CM, Storm JF, Ruth P. Cerebellar ataxia and Purkinje cell dysfunction caused by Ca^{2+} -activated K^+ channel deficiency. *Proc Natl Acad Sci USA* 2004;101:9474–9478. [PubMed: 15194823]
63. Pyott SJ, Meredith AL, Fodor AA, Vazquez AE, Yamoah EN, Aldrich RW. Cochlear function in mice lacking the BK channel alpha, beta1, or beta4 subunits. *J Biol Chem* 2007;282:3312–3324. [PubMed: 17135251]
64. Patterson AJ, Henrie-Olson J, Brenner R. Vasoregulation at the molecular level: a role for the beta1 subunit of the calcium-activated potassium (BK) channel. *Trends Cardiovasc Med* 2002;12:78–82. [PubMed: 11852255]

65. Amberg GC, Bonev AD, Rossow CF, Nelson MT, Santana LF. Modulation of the molecular composition of large conductance, Ca^{2+} activated K^{+} channels in vascular smooth muscle during hypertension. *J Clin Invest* 2003;112:717–724. [PubMed: 12952920]
66. Amberg GC, Santana LF. Downregulation of the BK channel beta1 subunit in genetic hypertension. *Circ Res* 2003;93:965–971. [PubMed: 14551242]
67. Fernandez-Fernandez JM, Tomas M, Vazquez E, Orio P, Latorre R, Senti M, Marrugat J, Valverde MA. Gain-of-function mutation in the KCNMB1 potassium channel subunit is associated with low prevalence of diastolic hypertension. *J Clin Invest* 2004;113:1032–1039. [PubMed: 15057310]
68. Senti M, Fernandez-Fernandez JM, Tomas M, Vazquez E, Elosua R, Marrugat J, Valverde MA. Protective effect of the KCNMB1 E65K genetic polymorphism against diastolic hypertension in aging women and its relevance to cardiovascular risk. *Circ Res* 2005;97:1360–1365. [PubMed: 16293791]
69. Chang T, Wu L, Wang R. Altered expression of BK channel beta1 subunit in vascular tissues from spontaneously hypertensive rats. *Am J Hypertens* 2006;19:678–685. [PubMed: 16814121]
70. Pluger S, Faulhaber J, Furstenau M, Lohn M, Waldschutz R, Gollasch M, Haller H, Luft FC, Ehmke H, Pongs O. Mice with disrupted BK channel beta1 subunit gene feature abnormal Ca^{2+} spark/STOC coupling and elevated blood pressure. *Circ Res* 2000;87:E53–60. [PubMed: 11090555]
71. Thorneloe KS, Meredith AL, Knorn AM, Aldrich RW, Nelson MT. Urodynamic properties and neurotransmitter dependence of urinary bladder contractility in the BK channel deletion model of overactive bladder. *Am J Physiol Renal Physiol* 2005;289:F604–610. [PubMed: 15827347]
72. Seibold MA, Wang B, Eng C, Kumar G, Beckman KB, Sen S, Choudhry S, Meade K, Lenoir M, Watson HG, Thyne S, Williams LK, Kumar R, Weiss KB, Grammer LC, Avila PC, Schleimer RP, Burchard EG, Brenner R. An african-specific functional polymorphism in KCNMB1 shows sex-specific association with asthma severity. *Hum Mol Genet* 2008;17:2681–2690. [PubMed: 18535015]
73. Womack MD, Chevez C, Khodakhah K. Calcium-activated potassium channels are selectively coupled to P/Q-type calcium channels in cerebellar Purkinje neurons. *J Neurosci* 2004;24:8818–8822. [PubMed: 15470147]
74. Chavis P, Ango F, Michel JM, Bockaert J, Fagni L. Modulation of big K^{+} channel activity by ryanodine receptors and L-type Ca^{2+} channels in neurons. *Eur J Neurosci* 1998;10:2322–2327. [PubMed: 9749760]
75. Marrion NV, Tavalin SJ. Selective activation of Ca^{2+} -activated K^{+} channels by co-localized Ca^{2+} channels in hippocampal neurons. *Nature* 1998;395:900–905. [PubMed: 9804423]
76. Gola M, Crest M. Colocalization of active K_{Ca} channels and Ca^{2+} channels within Ca^{2+} domains in helix neurons. *Neuron* 1993;10:689–699. [PubMed: 8386529]
77. Berkefeld H, Sailer CA, Bildl W, Rohde V, Thumfart JO, Eble S, Klugbauer N, Reisinger E, Bischofberger J, Oliver D, Knaus HG, Schulte U, Fakler B. BK_{Ca} -Cav channel complexes mediate rapid and localized Ca^{2+} -activated K^{+} signaling. *Science* 2006;314:615–620. [PubMed: 17068255]
78. Isaacson JS, Murphy GJ. Glutamate-mediated extrasynaptic inhibition: direct coupling of NMDA receptors to Ca^{2+} -activated K^{+} channels. *Neuron* 2001;31:1027–1034. [PubMed: 11580901]
79. Clapham DE. Calcium signaling. *Cell* 2007;131:1047–1058. [PubMed: 18083096]
80. Zhang X, Solaro CR, Lingle CJ. Allosteric regulation of BK channel gating by Ca^{2+} and Mg^{2+} through a nonselective, low affinity divalent cation site. *J Gen Physiol* 2001;118:607–636. [PubMed: 11696615]
81. Shi J, Cui J. Intracellular Mg^{2+} enhances the function of BK-type Ca^{2+} -activated K^{+} channels. *J Gen Physiol* 2001;118:589–606. [PubMed: 11696614]
82. Hille, B. Ion channels of excitable membranes. Sinauer; Sunderland, MA: 2001.
83. Toro L, Wallner M, Meera P, Tanaka Y. Maxi- K_{Ca} , a Unique Member of the Voltage-Gated K Channel Superfamily. *News Physiol Sci* 1998;13:112–117. [PubMed: 11390773]
84. Vergara C, Latorre R, Marrion NV, Adelman JP. Calcium-activated potassium channels. *Curr Opin Neurobiol* 1998;8:321–329. [PubMed: 9687354]
85. Salkoff L, Butler A, Ferreira G, Santi C, Wei A. High-conductance potassium channels of the SLO family. *Nat Rev Neurosci* 2006;7:921–931. [PubMed: 17115074]

86. Schubert R, Nelson MT. Protein kinases: tuners of the BK_{Ca} channel in smooth muscle. *Trends Pharmacol Sci* 2001;22:505–512. [PubMed: 11583807]
87. Weiger TM, Hermann A, Levitan IB. Modulation of calcium-activated potassium channels. *J Comp Physiol A Neuroethol Sens Neural Behav Physiol* 2002;188:79–87. [PubMed: 11919690]
88. Fury M, Marx SO, Marks AR. Molecular BKology: the study of splicing and dicing. *Sci STKE* 2002;2002:PE12.
89. Lu R, Alioua A, Kumar Y, Eghbali M, Stefani E, Toro L. MaxiK channel partners: physiological impact. *J Physiol (Lond)* 2006;570:65–72. [PubMed: 16239267]
90. Torres YP, Morera FJ, Carvacho I, Latorre R. A marriage of convenience: beta-subunits and voltage-dependent K⁺ channels. *J Biol Chem* 2007;282:24485–24489. [PubMed: 17606609]
91. Ghatta S, Nimmagadda D, Xu X, O'Rourke ST. Large-conductance, calcium-activated potassium channels: structural and functional implications. *Pharmacol Ther* 2006;110:103–116. [PubMed: 16356551]
92. Wu SN. Large-conductance Ca²⁺-activated K⁺ channels: physiological role and pharmacology. *Curr Med Chem* 2003;10:649–661. [PubMed: 12678784]
93. Calderone V. Large-conductance, Ca²⁺-activated K⁺ channels: function, pharmacology and drugs. *Curr Med Chem* 2002;9:1385–1395. [PubMed: 12132994]
94. Latorre R, Brauchi S. Large conductance Ca²⁺-activated K⁺ (BK) channel: activation by Ca²⁺ and voltage. *Biol Res* 2006;39:385–401. [PubMed: 17106573]
95. Magleby KL. Gating mechanism of BK (Slol) channels: so near, yet so far. *J Gen Physiol* 2003;121:81–96. [PubMed: 12566537]
96. Lingle CJ. Mg²⁺-dependent regulation of BK channels: importance of electrostatics. *J Gen Physiol* 2008;131:5–11. [PubMed: 18166623]
97. Lingle CJ. Gating rings formed by RCK domains: keys to gate opening. *J Gen Physiol* 2007;129:101–107. [PubMed: 17261839]
98. Rothberg BS. Allosteric modulation of ion channels: the case of maxi-K. *Sci STKE* 2004;2004:pe16.
99. Magleby KL. Kinetic gating mechanisms for BK channels: when complexity leads to simplicity. *J Gen Physiol* 2001;118:583–587. [PubMed: 11696613]
100. Lingle CJ. Setting the stage for molecular dissection of the regulatory components of BK channels. *J Gen Physiol* 2002;120:261–265. [PubMed: 12198086]
101. Cox, DH. BK_{Ca}-Channel Structure and Function. In: Chung, S-H.; Andersen, OS.; Krishnamurthy, V., editors. *Biological Membrane Ion Channels: Dynamics, Structure, and Applications*. Springer Science+Business Media, LLC; New York: 2007. p. 171-218.
102. Adelman JP, Shen KZ, Kavanaugh MP, Warren RA, Wu YN, Lagrutta A, Bond CT, North RA. Calcium-activated potassium channels expressed from cloned complementary DNAs. *Neuron* 1992;9:209–216. [PubMed: 1497890]
103. Atkinson NS, Robertson GA, Ganetzky B. A component of calcium-activated potassium channels encoded by the *Drosophila slo* locus. *Science* 1991;253:551–555. [PubMed: 1857984]
104. Butler A, Tsunoda S, McCobb DP, Wei A, Salkoff L. mSlo, a complex mouse gene encoding “maxi” calcium-activated potassium channels. *Science* 1993;261:221–224. [PubMed: 7687074]
105. Pallanck L, Ganetzky B. Cloning and characterization of human and mouse homologs of the *Drosophila* calcium-activated potassium channel gene, slowpoke. *Hum Mol Genet* 1994;3:1239–1243. [PubMed: 7987297]
106. Meera P, Wallner M, Song M, Toro L. Large conductance voltage- and calcium-dependent K⁺ channel, a distinct member of voltage-dependent ion channels with seven N-terminal transmembrane segments (S0–S6), an extracellular N terminus, and an intracellular (S9–S10) C terminus. *Proc Natl Acad Sci U S A* 1997;94:14066–14071. [PubMed: 9391153]
107. Wallner M, Meera P, Toro L. Determinant for beta-subunit regulation in high-conductance voltage-activated and Ca²⁺-sensitive K⁺ channels: an additional transmembrane region at the N terminus. *Proc Natl Acad Sci U S A* 1996;93:14922–14927. [PubMed: 8962157]
108. Quirk JC, Reinhart PH. Identification of a novel tetramerization domain in large conductance K_{Ca} channels. *Neuron* 2001;32:13–23. [PubMed: 11604135]

109. Shen KZ, Lagrutta A, Davies NW, Standen NB, Adelman JP, North RA. Tetraethylammonium block of Slowpoke calcium-activated potassium channels expressed in *Xenopus* oocytes: evidence for tetrameric channel formation. *Pflugers Arch* 1994;426:440–445. [PubMed: 7517033]
110. Jiang Y, Lee A, Chen J, Cadene M, Chait BT, MacKinnon R. Crystal structure and mechanism of a calcium-gated potassium channel. *Nature* 2002;417:515–522. [PubMed: 12037559]
111. Jiang Y, Pico A, Cadene M, Chait BT, MacKinnon R. Structure of the RCK domain from the *E. coli* K⁺ channel and demonstration of its presence in the human BK channel. *Neuron* 2001;29:593–601. [PubMed: 11301020]
112. Fodor AA, Aldrich RW. Statistical limits to the identification of ion channel domains by sequence similarity. *J Gen Physiol* 2006;127:755–766. [PubMed: 16735758]
113. Fodor AA, Aldrich RW. Convergent Evolution of Alternative Splices at Domain Boundaries of the BK Channel. *Annu Rev Physiol* 2009;71:1.1–1.18.
114. Albright RA, Ibar JL, Kim CU, Gruner SM, Morais-Cabral JH. The RCK domain of the KtrAB K⁺ transporter: multiple conformations of an octameric ring. *Cell* 2006;126:1147–1159. [PubMed: 16990138]
115. Long SB, Campbell EB, Mackinnon R. Crystal structure of a mammalian voltage-dependent Shaker family K⁺ channel. *Science* 2005;309:897–903. [PubMed: 16002581]
116. Ma Z, Lou XJ, Horrigan FT. Role of Charged Residues in the S1–S4 Voltage Sensor of BK Channels. *J Gen Physiol* 2006;127:309–328. [PubMed: 16505150]
117. Liu G, Zakharov SI, Yang L, Deng SX, Landry DW, Karlin A, Marx SO. Position and role of the BK channel alpha subunit S0 helix inferred from disulfide crosslinking. *J Gen Physiol* 2008;131:537–548. [PubMed: 18474637]
118. Yang H, Shi J, Zhang G, Yang J, Delaloye K, Cui J. Activation of Slol BK channels by Mg²⁺ coordinated between the voltage sensor and the RCK1 domains. *Nat Struct Mol Biol*. 2008;10.1038/nsmbl507
119. Liu G, Zakharov SI, Yang L, Wu RS, Deng SX, Landry DW, Karlin A, Marx SO. Locations of the betal transmembrane helices in the BK potassium channel. *Proc Natl Acad Sci U S A* 2008;105:10727–10732. [PubMed: 18669652]
120. Koval OM, Fan Y, Rothberg BS. A role for the S0 transmembrane segment in voltage-dependent gating of BK channels. *J Gen Physiol* 2007;129:209–220. [PubMed: 17296928]
121. Shi J, Krishnamoorthy G, Yang Y, Hu L, Chaturvedi N, Harilal D, Qin J, Cui J. Mechanism of magnesium activation of calcium-activated potassium channels. *Nature* 2002;418:876–880. [PubMed: 12192410]
122. Xia XM, Zeng X, Lingle CJ. Multiple regulatory sites in large-conductance calcium-activated potassium channels. *Nature* 2002;418:880–884. [PubMed: 12192411]
123. Tang XD, Xu R, Reynolds MF, Garcia ML, Heinemann SH, Hoshi T. Haem can bind to and inhibit mammalian calcium-dependent Slol BK channels. *Nature* 2003;425:531–535. [PubMed: 14523450]
124. Yusifov T, Savalli N, Gandhi CS, Ottolia M, Olcese R. The RCK2 domain of the human BK_{Ca} channel is a calcium sensor. *Proc Natl Acad Sci U S A* 2008;105:376–381. [PubMed: 18162557]
125. Kim HJ, Lim HH, Rho SH, Bao L, Lee JH, Cox DH, Kim do H, Park CS. Modulation of the conductance-voltage relationship of the BK_{Ca} channel by mutations at the putative flexible interface between two RCK domains. *Biophys J* 2008;94:446–456. [PubMed: 17890381]
126. Kim HJ, Lim HH, Rho SH, Eom SH, Park CS. Hydrophobic interface between two regulators of K⁺ conductance domains critical for calcium-dependent activation of large conductance Ca²⁺-activated K⁺ channels. *J Biol Chem* 2006;281:38573–38581. [PubMed: 17040919]
127. Wei A, Solaro C, Lingle C, Salkoff L. Calcium sensitivity of BK-type K_{Ca} channels determined by a separable domain. *Neuron* 1994;13:671–681. [PubMed: 7917297]
128. Yang H, Hu L, Shi J, Cui J. Tuning Magnesium Sensitivity of BK Channels by Mutations. *Biophys J* 2006;91:2892–2900. [PubMed: 16877509]
129. Yang H, Hu L, Shi J, Delaloye K, Horrigan FT, Cui J. Mg²⁺ mediates interaction between the voltage sensor and cytosolic domain to activate BK channels. *Proc Natl Acad Sci U S A* 2007;104:18270–18275. [PubMed: 17984060]

130. Hou S, Xu R, Heinemann SH, Hoshi T. Reciprocal regulation of the Ca^{2+} and H^{+} sensitivity in the SLO1 BK channel conferred by the RCK1 domain. *Nat Struct Mol Biol* 2008;15:403–410. [PubMed: 18345016]
131. Zhang Z, Zhou Y, Ding JP, Xia XM, Lingle CJ. A limited access compartment between the pore domain and cytosolic domain of the BK channel. *J Neurosci* 2006;26:11833–11843. [PubMed: 17108156]
132. Meera P, Wallner M, Jiang Z, Toro L. A calcium switch for the functional coupling between alpha (hslo) and beta subunits (KV, Ca beta) of maxi K channels. *FEBS Lett* 1996;382:84–88.
133. Cui J, Cox DH, Aldrich RW. Intrinsic voltage dependence and Ca^{2+} regulation of mslo large conductance Ca -activated K^{+} channels. *J Gen Physiol* 1997;109:647–673. [PubMed: 9154910]
134. Stefani E, Ottolia M, Noceti F, Olcese R, Wallner M, Latorre R, Toro L. Voltage-controlled gating in a large conductance Ca^{2+} -sensitive K^{+} channel (hslo). *Proc Natl Acad Sci U S A* 1997;94:5427–5431. [PubMed: 9144254]
135. Horrigan FT, Aldrich RW. Allosteric voltage gating of potassium channels II. Mslo channel gating charge movement in the absence of Ca^{2+} . *J Gen Physiol* 1999;114:305–336. [PubMed: 10436004]
136. Diaz L, Meera P, Amigo J, Stefani E, Alvarez O, Toro L, Latorre R. Role of the S4 segment in a voltage-dependent calcium-sensitive potassium (hSlo) channel. *J Biol Chem* 1998;273:32430–32436. [PubMed: 9829973]
137. Cui J, Aldrich RW. Allosteric linkage between voltage and Ca^{2+} -dependent activation of BK-type mslo1 K^{+} channels. *Biochemistry (Mosc)* 2000;39:15612–15619.
138. Bezanilla F. The voltage sensor in voltage-dependent ion channels. *Physiol Rev* 2000;80:555–592. [PubMed: 10747201]
139. Gandhi CS, Isacoff EY. Molecular Models of Voltage Sensing. *J Gen Physiol* 2002;120:455–463. [PubMed: 12356848]
140. Hu L, Shi J, Ma Z, Krishnamoorthy G, Sieling F, Zhang G, Horrigan FT, Cui J. Participation of the S4 voltage sensor in the Mg^{2+} -dependent activation of large conductance (BK) K^{+} channels. *Proc Natl Acad Sci U S A* 2003;100:10488–10493. [PubMed: 12925732]
141. Tiwari-Woodruff SK, Schulteis CT, Mock AF, Papazian DM. Electrostatic interactions between transmembrane segments mediate folding of Shaker K^{+} channel subunits. *Biophys J* 1997;72:1489–1500. [PubMed: 9083655]
142. Silverman WR, Roux B, Papazian DM. Structural basis of two-stage voltage-dependent activation in K^{+} channels. *Proc Natl Acad Sci U S A* 2003;100:2935–2940. [PubMed: 12606713]
143. Horrigan FT, Cui J, Aldrich RW. Allosteric voltage gating of potassium channels I. Mslo ionic currents in the absence of Ca^{2+} . *J Gen Physiol* 1999;114:277–304. [PubMed: 10436003]
144. Talukder G, Aldrich RW. Complex voltage-dependent behavior of single unliganded calcium-sensitive potassium channels. *Biophys J* 2000;78:761–772. [PubMed: 10653789]
145. Nimigean CM, Magleby KL. Functional coupling of the beta₁ subunit to the large conductance Ca^{2+} -activated K^{+} channel in the absence of Ca^{2+} . Increased Ca^{2+} sensitivity from a Ca^{2+} -independent mechanism. *J Gen Physiol* 2000;115:719–736. [PubMed: 10828246]
146. Rothberg BS, Magleby KL. Voltage and Ca^{2+} activation of single large-conductance Ca^{2+} -activated K^{+} channels described by a two-tiered allosteric gating mechanism. *J Gen Physiol* 2000;116:75–99. [PubMed: 10871641]
147. Savalli N, Kondratiev A, Toro L, Olcese R. Voltage-dependent conformational changes in human Ca^{2+} - and voltage-activated K^{+} channel, revealed by voltage-clamp fluorometry. *Proc Natl Acad Sci U S A* 2006;103:12619–12624. [PubMed: 16895996]
148. Schoppa NE, Sigworth FJ. Activation of shaker potassium channels. I Characterization of voltage-dependent transitions. *J Gen Physiol* 1998;111:271–294. [PubMed: 9450944]
149. Ledwell JL, Aldrich RW. Mutations in the S4 region isolate the final voltage-dependent cooperative step in potassium channel activation. *J Gen Physiol* 1999;113:389–414. [PubMed: 10051516]
150. Lu Z, Klem AM, Ramu Y. Coupling between Voltage Sensors and Activation Gate in Voltage-gated K^{+} Channels. *J Gen Physiol* 2002;120:663–676. [PubMed: 12407078]
151. McCormack K, Joiner WJ, Heinemann SH. A characterization of the activating structural rearrangements in voltage-dependent Shaker K^{+} channels. *Neuron* 1994;12:301–315. [PubMed: 8110460]

152. Lu Z, Klem AM, Ramu Y. Ion conduction pore is conserved among potassium channels. *Nature* 2001;413:809–813. [PubMed: 11677598]
153. Soler-Llavina GJ, Chang TH, Swartz KJ. Functional interactions at the interface between voltage-sensing and pore domains in the Shaker K_v channel. *Neuron* 2006;52:623–634. [PubMed: 17114047]
154. Niu X, Qian X, Magleby KL. Linker-gating ring complex as passive spring and Ca²⁺-dependent machine for a voltage- and Ca²⁺-activated potassium channel. *Neuron* 2004;42:745–756. [PubMed: 15182715]
155. Schreiber M, Salkoff L. A novel calcium-sensing domain in the BK channel. *Biophys J* 1997;73:1355–1363. [PubMed: 9284303]
156. Bao L, Kaldany C, Holmstrand BC, Cox DH. Mapping the BK_{Ca} channel's "Ca²⁺ bowl": side-chains essential for Ca²⁺ sensing. *J Gen Physiol* 2004;123:475–489. [PubMed: 15111643]
157. Bao L, Rapin AM, Holmstrand BC, Cox DH. Elimination of the BK_{Ca} channel's high-affinity Ca²⁺ sensitivity. *J Gen Physiol* 2002;120:173–189. [PubMed: 12149279]
158. Moss GW, Marshall J, Moczydlowski E. Hypothesis for a serine proteinase-like domain at the COOH terminus of Slowpoke calcium-activated potassium channels. *J Gen Physiol* 1996;108:473–484. [PubMed: 8972386]
159. Moss GW, Marshall J, Morabito M, Howe JR, Moczydlowski E. An evolutionarily conserved binding site for serine proteinase inhibitors in large conductance calcium-activated potassium channels. *Biochemistry (Mosc)* 1996;35:16024–16035.
160. Bian S, Favre I, Moczydlowski E. Ca²⁺-binding activity of a COOH-terminal fragment of the *Drosophila* BK channel involved in Ca²⁺-dependent activation. *Proc Natl Acad Sci U S A* 2001;98:4776–4781. [PubMed: 11274367]
161. Braun A, Sy L. Contribution of potential EF hand motifs to the calcium-dependent gating of a mouse brain large conductance, calcium-sensitive K⁺ channel. *J Physiol (Lond)* 2001;533:681–695. [PubMed: 11410626]
162. Niu X, Magleby KL. Stepwise contribution of each subunit to the cooperative activation of BK channels by Ca²⁺. *Proc Natl Acad Sci U S A* 2002;99:11441–11446. [PubMed: 12161564]
163. Schreiber M, Yuan A, Salkoff L. Transplantable sites confer calcium sensitivity to BK channels. *Nat Neurosci* 1999;2:416–421. [PubMed: 10321244]
164. Zeng XH, Xia XM, Lingle CJ. Divalent cation sensitivity of BK channel activation supports the existence of three distinct binding sites. *J Gen Physiol* 2005;125:273–286. [PubMed: 15738049]
165. Hou S, Xu R, Heinemann SH, Hoshi T. The RCK1 high-affinity Ca²⁺ sensor confers carbon monoxide sensitivity to Slo1 BK channels. *Proc Natl Acad Sci U S A* 2008;105:4039–4043. [PubMed: 18316727]
166. Hu L, Yang H, Shi J, Cui J. Effects of Multiple Metal Binding Sites on Calcium and Magnesium-dependent Activation of BK Channels. *J Gen Physiol* 2006;127:35–50. [PubMed: 16344323]
167. Harding MM. The architecture of metal coordination groups in proteins. *Acta Crystallogr D Biol Crystallogr* 2004;60:849–859. [PubMed: 15103130]
168. Dudev T, Lim C. Principles governing Mg, Ca, and Zn binding and selectivity in proteins. *Chem Rev* 2003;103:773–788. [PubMed: 12630852]
169. Cox DH. The BK_{Ca} channel's Ca²⁺-binding sites, multiple sites, multiple ions. *J Gen Physiol* 2005;125:253–255. [PubMed: 15738047]
170. Qian X, Niu X, Magleby KL. Intra- and Intersubunit Cooperativity in Activation of BK Channels by Ca²⁺. *J Gen Physiol* 2006;128:389–404. [PubMed: 17001085]
171. Piskorowski R, Aldrich RW. Calcium activation of BK_{Ca} potassium channels lacking the calcium bowl and RCK domains. *Nature* 2002;420:499–502. [PubMed: 12466841]
172. Bravo-Zehnder M, Orio P, Norambuena A, Wallner M, Meera P, Toro L, Latorre R, Gonzalez A. Apical sorting of a voltage- and Ca²⁺-activated K⁺ channel alpha-subunit in Madin-Darby canine kidney cells is independent of N-glycosylation. *Proc Natl Acad Sci U S A* 2000;97:13114–13119. [PubMed: 11069304]
173. Schmalhofer WA, Sanchez M, Dai G, Dewan A, Secades L, Hanner M, Knaus HG, McManus OB, Kohler M, Kaczorowski GJ, Garcia ML. Role of the C-terminus of the high-conductance calcium-

activated potassium channel in channel structure and function. *Biochemistry (Mosc)* 2005;44:10135–10144.

174. Krause JD, Foster CD, Reinhart PH. *Xenopus laevis* oocytes contain endogenous large conductance Ca^{2+} -activated K^{+} channels. *Neuropharmacology* 1996;35:1017–1022. [PubMed: 8938732]
175. Qian X, Magleby KL. Beta subunits facilitate gating of BK channels by acting through the Ca^{2+} , but not the Mg^{2+} activating mechanisms. *Proc Natl Acad Sci U S A* 2003;100:10061–10066. [PubMed: 12893878]
176. Xia XM, Zhang X, Lingle CJ. Ligand-dependent activation of Slo family channels is defined by interchangeable cytosolic domains. *J Neurosci* 2004;24:5585–5591. [PubMed: 15201331]
177. Pico, AR. PhD Thesis. 2003. RCK Domain Model of Calcium Activation in BK channels.
178. Yang, H. PhD Thesis. 2008. Molecular Gating Mechanism of BK-type Large Conductance Potassium Channels.
179. Horrigan FT, Aldrich RW. Coupling between voltage sensor activation, Ca^{2+} binding and channel opening in large conductance (BK) potassium channels. *J Gen Physiol* 2002;120:267–305. [PubMed: 12198087]
180. Moczydlowski E, Latorre R. Gating kinetics of Ca^{2+} -activated K^{+} channels from rat muscle incorporated into planar lipid bilayers. Evidence for two voltage-dependent Ca^{2+} binding reactions. *J Gen Physiol* 1983;82:511–542. [PubMed: 6315857]
181. McManus OB, Magleby KL. Accounting for the Ca^{2+} -dependent kinetics of single large-conductance Ca^{2+} -activated K^{+} channels in rat skeletal muscle. *J Physiol (Lond)* 1991;443:739–777. [PubMed: 1822543]
182. DiChiara TJ, Reinhart PH. Distinct effects of Ca^{2+} and voltage on the activation and deactivation of cloned Ca^{2+} -activated K^{+} channels. *J Physiol (Lond)* 1995;489:403–418. [PubMed: 8847636]
183. Nimigeon CM, Magleby KL. The beta subunit increases the Ca^{2+} sensitivity of large conductance Ca^{2+} -activated potassium channels by retaining the gating in the bursting states. *J Gen Physiol* 1999;113:425–440. [PubMed: 10051518]
184. Monod J, Wyman J, Changeux JP. On the nature of allosteric transitions: a plausible model. *J molec Biol* 1965;12:88–118. [PubMed: 14343300]
185. Cox DH, Cui J, Aldrich RW. Allosteric gating of a large conductance Ca^{2+} -activated K^{+} channel. *J Gen Physiol* 1997;110:257–281. [PubMed: 9276753]
186. Methfessel C, Boheim G. The gating of single calcium-dependent potassium channels is described by an activation/blockade mechanism. *Biophys Struct Mech* 1982;9:35–60. [PubMed: 6293598]
187. Pallotta BS. N-bromoacetamide removes a calcium-dependent component of channel opening from calcium-activated potassium channels in rat skeletal muscle. *J Gen Physiol* 1985;86:601–611. [PubMed: 2415669]
188. Ye S, Li Y, Chen L, Jiang Y. Crystal structures of a ligand-free MthK gating ring: insights into the ligand gating mechanism of K^{+} channels. *Cell* 2006;126:1161–1173. [PubMed: 16990139]
189. Krishnamoorthy G, Shi J, Sept D, Cui J. The NH2 terminus of RCK1 domain regulates Ca^{2+} -dependent BK_{Ca} channel gating. *J Gen Physiol* 2005;126:227–241. [PubMed: 16103277]
190. Golowasch J, Kirkwood A, Miller C. Allosteric effects of Mg^{2+} on the gating of Ca^{2+} -activated K^{+} channels from mammalian skeletal muscle. *J Exp Biol* 1986;124:5–13. [PubMed: 2428908]
191. Oberhauser A, Alvarez O, Latorre R. Activation by divalent cations of a Ca^{2+} -activated K^{+} channel from skeletal muscle membrane. *J Gen Physiol* 1988;92:67–86. [PubMed: 3171535]
192. Horrigan FT, Ma Z. Mg^{2+} enhances voltage sensor/gate coupling in BK channels. *J Gen Physiol* 2008;131:13–32. [PubMed: 18166624]
193. Flatman PW. Mechanisms of magnesium transport. *Annu Rev Physiol* 1991;53:259–271. [PubMed: 2042962]
194. Gouaux E, Mackinnon R. Principles of selective ion transport in channels and pumps. *Science* 2005;310:1461–1465. [PubMed: 16322449]
195. Hille B, Armstrong CM, MacKinnon R. Ion channels: from idea to reality. *Nat Med* 1999;5:1105–1109. [PubMed: 10502800]
196. Jiang Y, Lee A, Chen J, Cadene M, Chait BT, MacKinnon R. The open pore conformation of potassium channels. *Nature* 2002;417:523–526. [PubMed: 12037560]

197. Webster SM, Del Camino D, Dekker JP, Yellen G. Intracellular gate opening in Shaker K⁺ channels defined by high-affinity metal bridges. *Nature* 2004;428:864–868. [PubMed: 15103379]
198. Nimigean CM, Chappie JS, Miller C. Electrostatic tuning of ion conductance in potassium channels. *Biochemistry (Mosc)* 2003;42:9263–9268.
199. Brelidze TI, Niu X, Magleby KL. A ring of eight conserved negatively charged amino acids doubles the conductance of BK channels and prevents inward rectification. *Proc Natl Acad Sci U S A* 2003;100:9017–9022. [PubMed: 12843404]
200. Haug T, Olcese R, Toro L, Stefani E. Regulation of K⁺ flow by a ring of negative charges in the outer pore of BK_{ca} channels. Part II: Neutralization of aspartate 292 reduces long channel openings and gating current slow component. *J Gen Physiol* 2004;124:185–197. [PubMed: 15277579]
201. Haug T, Sigg D, Ciani S, Toro L, Stefani E, Olcese R. Regulation of K⁺ flow by a ring of negative charges in the outer pore of BK_{ca} channels. Part I: Aspartate 292 modulates K⁺ conduction by external surface charge effect. *J Gen Physiol* 2004;124:173–184. [PubMed: 15277578]
202. Brelidze TI, Magleby KL. Probing the geometry of the inner vestibule of BK channels with sugars. *J Gen Physiol* 2005;126:105–121. [PubMed: 16043773]
203. Li W, Aldrich RW. Unique inner pore properties of BK channels revealed by quaternary ammonium block. *J Gen Physiol* 2004;124:43–57. [PubMed: 15197222]
204. Li W, Aldrich RW. State-dependent Block of BK Channels by Synthesized Shaker Ball Peptides. *J Gen Physiol* 2006;128:423–441. [PubMed: 16966472]
205. Wilkens CM, Aldrich RW. State-independent block of BK channels by an intracellular quaternary ammonium. *J Gen Physiol* 2006;128:347–364. [PubMed: 16940557]
206. Flynn GE, Zagotta WN. Conformational changes in S6 coupled to the opening of cyclic nucleotide-gated channels. *Neuron* 2001;30:689–698. [PubMed: 11430803]
207. Bruening-Wright A, Lee WS, Adelman JP, Maylie J. Evidence for a Deep Pore Activation Gate in Small Conductance Ca²⁺-activated K⁺ Channels. *J Gen Physiol* 2007;130:601–610. [PubMed: 17998394]
208. Piskorowski RA, Aldrich RW. Relationship between pore occupancy and gating in BK potassium channels. *J Gen Physiol* 2006;127:557–576. [PubMed: 16636204]
209. Guo Z, Lv C, Yi H, Xiong Y, Wu Y, Li W, Xu T, Ding J. A residue at the cytoplasmic entrance of BK-type channels regulating single-channel opening by its hydrophobicity. *Biophys J* 2008;94:3714–3725. [PubMed: 18400952]
210. Magidovich E, Yifrach O. Conserved gating hinge in ligand- and voltage-dependent K⁺ channels. *Biochemistry (Mosc)* 2004;43:13242–13247.
211. Ding S, Ingleby L, Ahern CA, Horn R. Investigating the putative glycine hinge in Shaker potassium channel. *J Gen Physiol* 2005;126:213–226. [PubMed: 16103276]
212. Wang B, Brenner R. An S6 Mutation in BK Channels Reveals {beta}1 Subunit Effects on Intrinsic and Voltage-dependent Gating. *J Gen Physiol* 2006;128:731–744. [PubMed: 17130522]
213. Lippiat JD, Standen NB, Davies NW. A residue in the intracellular vestibule of the pore is critical for gating and permeation in Ca²⁺-activated K⁺ (BK_{ca}) channels. *J Physiol (Lond)* 2000;529:131–138. [PubMed: 11080257]
214. Tang XD, Garcia ML, Heinemann SH, Hoshi T. Reactive oxygen species impair Slol BK channel function by altering cysteine-mediated calcium sensing. *Nat Struct Mol Biol* 2004;11:171–178. [PubMed: 14745441]
215. Santarelli LC, Chen J, Heinemann SH, Hoshi T. The beta 1 subunit enhances oxidative regulation of large-conductance calcium-activated K⁺ channels. *J Gen Physiol* 2004;124:357–370. [PubMed: 15452197]
216. Santarelli LC, Wassef R, Heinemann SH, Hoshi T. Three methionine residues located within the regulator of conductance for K⁺ (RCK) domains confer oxidative sensitivity to large-conductance Ca²⁺-activated K⁺ channels. *J Physiol (Lond)* 2006;571:329–348. [PubMed: 16396928]
217. Tang XD, Daggett H, Manner M, Garcia ML, McManus OB, Brot N, Weissbach H, Heinemann SH, Hoshi T. Oxidative regulation of large conductance calcium-activated potassium channels. *J Gen Physiol* 2001;117:253–274. [PubMed: 11222629]

218. Lunin VV, Dobrovetsky E, Khutoreskaya G, Zhang R, Joachimiak A, Doyle DA, Bochkarev A, Maguire ME, Edwards AM, Koth CM. Crystal structure of the CorA Mg^{2+} transporter. *Nature* 2006;440:833–837. [PubMed: 16598263]
219. Avdonin V, Tang XD, Hoshi T. Stimulatory action of internal protons on Slol BK channels. *Biophys J* 2003;84:2969–2980. [PubMed: 12719228]
220. Horrigan FT, Heinemann SH, Hoshi T. Heme Regulates Allosteric Activation of the Slol BK Channel. *J Gen Physiol* 2005;126:7–21. [PubMed: 15955873]
221. Vaithianathan T, Bukiya A, Liu J, Liu P, Asuncion-Chin M, Fan Z, Dopico A. Direct regulation of BK channels by phosphatidylinositol 4,5-bisphosphate as a novel signaling pathway. *J Gen Physiol* 2008;132:13–28. [PubMed: 18562499]
222. Orio P, Latorre R. Differential effects of beta 1 and beta 2 subunits on BK channel activity. *J Gen Physiol* 2005;125:395–411. [PubMed: 15767297]
223. Orio P, Torres Y, Rojas P, Carvacho I, Garcia ML, Toro L, Valverde MA, Latorre R. Structural determinants for functional coupling between the beta and alpha subunits in the Ca^{2+} -activated K^+ (BK) channel. *J Gen Physiol* 2006;127:191–204. [PubMed: 16446507]
224. Bao L, Cox DH. Gating and ionic currents reveal how the BK_{Ca} channel's Ca^{2+} sensitivity is enhanced by its beta 1 subunit. *J Gen Physiol* 2005;126:393–412. [PubMed: 16186565]
225. Cox DH, Aldrich RW. Role of the beta 1 subunit in large-conductance Ca^{2+} -activated K^+ channel gating energetics. Mechanisms of enhanced Ca^{2+} sensitivity. *J Gen Physiol* 2000;116:411–432. [PubMed: 10962017]
226. Yang H, Zhang G, Shi J, Lee US, Delaloye K, Cui J. Subunit-specific effect of the voltage sensor domain on Ca^{2+} sensitivity of BK channels. *Biophys J* 2008;94:4678–4687. [PubMed: 18339745]
227. Morrow JP, Zakharov SI, Liu G, Yang L, Sok AJ, Marx SO. Defining the BK channel domains required for beta 1-subunit modulation. *Proc Natl Acad Sci U S A* 2006;103:5096–5101. [PubMed: 16549765]
228. Brenner R, Jegla TJ, Wickenden A, Liu Y, Aldrich RW. Cloning and functional characterization of novel large conductance calcium-activated potassium channel beta subunits, hKCNMB3 and hKCNMB4. *J Biol Chem* 2000;275:6453–6461. [PubMed: 10692449]
229. Xia XM, Ding JP, Lingle CJ. Inactivation of BK channels by the NH2 terminus of the beta2 auxiliary subunit: an essential role of a terminal peptide segment of three hydrophobic residues. *J Gen Physiol* 2003;121:125–148. [PubMed: 12566540]
230. Lingle CJ, Zeng XH, Ding JP, Xia XM. Inactivation of BK channels mediated by the NH2 terminus of the beta3b auxiliary subunit involves a two-step mechanism: possible separation of binding and blockade. *J Gen Physiol* 2001;117:583–606. [PubMed: 11382808]
231. McManus OB, Helms LM, Pallanck L, Ganetzky B, Swanson R, Leonard RJ. Functional role of the beta subunit of high conductance calcium-activated potassium channels. *Neuron* 1995;14:645–650. [PubMed: 7695911]
232. Xia XM, Ding JP, Lingle CJ. Molecular basis for the inactivation of Ca^{2+} - and voltage-dependent BK channels in adrenal chromaffin cells and rat insulinoma tumor cells. *J Neurosci* 1999;19:5255–5264. [PubMed: 10377337]
233. Wallner M, Meera P, Toro L. Molecular basis of fast inactivation in voltage and Ca^{2+} -activated K^+ channels: a transmembrane beta-subunit homolog. *Proc Natl Acad Sci U S A* 1999;96:4137–4142. [PubMed: 10097176]
234. Roosild TP, Le KT, Choe S. Cytoplasmic gatekeepers of K^+ -channel flux: a structural perspective. *Trends Biochem Sci* 2004;29:39–45. [PubMed: 14729331]

Figure 1.

BK channel activation by voltage, Ca^{2+} and Mg^{2+} . **A.** The conductance-voltage (G-V) relationship of mSlo1 BK channels in 0 (black), 2 (blue) and 100 μM Ca^{2+} (green). **B.** The G-V relationship of BK channels in 0 (black) and 10 mM Mg^{2+} (red). With an increasing of $[\text{Ca}^{2+}]_i$ (**A**) or $[\text{Mg}^{2+}]_i$ (**B**), the BK channels open at less positive voltages.

Figure 2.

BK channel structure. **A.** Membrane topology of the Slo1 subunit. VSD: voltage sensor domain; P: pore loop. **B.** Kv1.2 structure (PDB ID: 2A79) [115]. Four subunits are represented in different colors. Only the membrane-spanning domain of the structure is shown. **C.** The gating ring structure of MthK (PDB ID: 1LNQ) [110]. Eight identical RCK domains are represented in different colors. **D.** BK channel homology model based on the superposition of the Kv1.2 structure and the MthK structure by aligning their selectivity filter regions. Four different colors represent four subunits. Only the membrane spanning domain of Kv1.2 (ribbons) and the four RCK domains of MthK close to the membrane (surface) are shown. Left: side view. Right: top

view from the membrane. Blue arrows indicate that the relative arrangement of the VSD and the cytoplasmic domain of BK channels might be different from the homology model [118].

Figure 3.

Allosteric gating mechanisms of BK channels. **A.** Mechanism of voltage dependent gating [143]. Channels can open when 0–4 voltage sensors activate in the four subunits, resulting in 5 closed (C_0 – C_4) and 5 open (O_0 – O_4) states. J represents the equilibrium constant for voltage sensor activation in each subunit. $J = \exp((V - V_{hC})(z_J e/kT))$, where V is voltage, $V_{hC} = +150$ mV is the half activation voltage at the closed state, $z_J = 0.58$ is the number of charge movement during a voltage sensor activation, e is the charge of an electron, k is Boltzmann's constant, and T is absolute temperature. L is the intrinsic equilibrium constant for channel opening when all voltage sensors are in the resting state. $L = L_0 \exp(V)(z_L e/kTy)$, where $L_0 = 9.8 \times 10^{-7}$ is the equilibrium constant at 0 mV and $z_L = 0.3$ is the number of equivalent charge movement during gate opening. $D = 25$ is the allosteric factor coupling voltage sensor activation to channel opening. **B.** A general allosteric mechanism including voltage and Ca^{2+} dependent activation and the interaction between Ca^{2+} binding and voltage sensor activation [179]. The allosteric coupling (with allosteric factor D) between voltage sensor activation $[R-A]_4$ and channel opening $[C-O]$ can be expanded as the 10-state scheme in A. Likewise, the allosteric coupling (with allosteric factor C) between Ca^{2+} binding $[X-XCa^{2+}]_4$ and channel opening $[C-O]$ can be expanded into a similar 10-state scheme. Here it is assumed that the channel contains four Ca^{2+} binding sites although the model can be easily expanded to contain eight Ca^{2+} binding sites. Channels can open when 0–4 Ca^{2+} are bound, resulting in 5 closed (C_0 – C_4) and 5 open (O_0 – O_4) states. K_{Ca} is the equilibrium constant for Ca^{2+} binding. $K_{Ca} = [Ca^{2+}]_i/K_{dC}$ at closed states and $[Ca^{2+}]_i/K_{dO}$ at open states, where $K_{dC} = 11 \mu M$ and $K_{dO} = 1.4 \mu M$ are the elementary Ca^{2+} dissociation constants. $C = K_{dC}/K_{dO} = 8$. The allosteric factor E describes the interaction

between Ca^{2+} binding and voltage sensor activation. At saturating Ca^{2+} binding, J increases by the factor $E = 2.4$.

Table 1
Localization and identification of important structural features in BK channels. The residue numbers refer to the position on *mbr5* splice variant of mSlol [104]. The locations of the fixed or flexible interface in RCK1 or RCK2 are identified by comparing the sequence alignment to that of MthK [110,111] without any experimental evidence, except for the residues denoted by * that were experimentally identified by Kim et al [125,126].

Ref (structure used for alignment)	Domains		Interfaces				Ca ²⁺ Binding Sites		
	RCK1	RCK2	Fixed		Flexible		D362	D367	Ca ²⁺ bowl
			RCK1	RCK2	RCK1	RCK2			
Jiang et al., 2001 [111] (E.coli)	344–514 (βA-αG)		αD (440–448)		αF (489–497) αG (507–514)		middle-end of αA	between αA and βB	
Jiang et al., 2002 [110] (MthK)	344–611 (βA-αJ)		αD (437–450) αE (465–473)		αF (487–496) αG (502–512)		middle-end of αA	between αA and βB	
A. Pico's Thesis, 2003 [177] (MthK)	344–612 (βA-αJ)	721–1055 (βA-βJ)	αD (437–450)	αD (820–833)	αF (487–498) αG (501–511)		middle-end of αA	between αA and βB	between βE and αE
Roosild et al., 2004 [234] (MthK)	345–498 (βA-αF)	722–871 (βA-αF)	αD (435–447)	αD (818–830)	αF (487–98)		middle of αA	end of αA	after RCK2 domain
Kim et al., 2006 & 2008 [126,125] (MthK)	344–619 (βA-αD)	721–879 (βA-βF)	αD (437–456) αD (441)* αD (442)* α (445)*	αD (820–830) αD (824)* αD (825)* αD (828)*	αF (487–496) αG (502–512) αF (490)*		middle of αA	end of αA	after RCK2 domain
Latorre and Brauchi, 2006 [94] (E.coli)	344–512 (βA-αD - normal β-α-β structure; after that all alpha helices - αX, αE, αF and αG)		αD (437–450)		αF (482–496) αG (504–512)		middle of αA	between αA and βB	
Yusifov et al., 2008 [124] (MthK)	344–612 (βA-αJ)	721–987 (βA-αJ)	αD (437–450)	αD (820–833)	αF (486–494) αG (503–512)		end of αA	between αA and βB	between αG and βG (891–902)