CORRESPONDENCE

Pertussis: Not Only a Disease of Childhood

by Dr. med. Dipl. Biol. Marion Riffelmann, Dr. med. Martina Littmann,
Dr. med. Wiebke Hellenbrand, Prof. Dr. med. Christel Hülbe, Prof. Dr. med.
Carl Heinz Wirsing von König in volume 37/2008

At a Loss

The review article leaves me somewhat at a loss. To grow a *Bordetella* culture or perform polymerase chain reaction (PCR) in suspected pertussis would not have occurred to me, a doctor who treats members of statutory medical insurance schemes. I think this is extremely uneconomical. The authors do not discuss measuring pertussis IgM antibodies, which is common and relevant in everyday clinical practice.

The suggested treatments lack practical application. The acute phase is not the problem. Any antibiotic would help here, provided the patient consults the doctor during this phase. Mostly, patients attend too late. Any discussion of antibiotics comes too late for a patient with persisting cough.

DOI: 10.3238/arztebl.2009.0057b

REFERENCES

1. Riffelmann M, Littmann M, Hellenbrand W, Hülbe C, Wirsing von König CH: Pertussis: Not Only a Disease of Childhood. *Dtsch Arztebl Int* 2008; 105(37): 623–8.

Norbert K. Mülleneisen

Königsberger Platz 5
51371 Leverkusen, Germany

Conflict of interest statement

The author receives research funding from Novartis, speaker's honoraria from AstraZeneca, Bayer, GlaxoSmithKline, Boehringer Ingelheim, and Dr Beckmann-Roxall. He is the clinical lead of the North-Rhine federation of doctors treating members of statutory medical insurance schemes and until 2008 was deputy chairman of North Rhine Westphalia's professional specialty association of pneumologists.

Cautious Advice

An experienced paediatrician, who healed patients long before the antibiotic era, advised to treat my daughter's croup coughing fits during colds with carefully dosed promethazine and dihydrocodein, a combination that worked every time.

Three years ago I managed to successfully treat coughing fits in an adult woman who suffered fits of suffocation during (probably) existing whooping cough.

DOI: 10.3238/arztebl.2009.0057b

REFERENCES

1. Riffelmann M, Littmann M, Hellenbrand W, Hülbe C, Wirsing von König CH: Pertussis: Not Only a Disease of Childhood. *Dtsch Arztebl Int* 2008; 105(37): 623–8.

Dr. med. Lieselotte Windhövel

Coubertinstr. 16
95615 Marktreidwitz, Germany

Conflict of interest statement

The author declares that no conflict of interest exists according to the guidelines of the International Committee of Medical Journal Editors.

Selection of Pneumococci

Are there any data from Germany about the sensitivity of pertussis strains to macrolides, trimethoprim-sulfamethoxazole, and tetracyclines, which are recommended for therapy, respectively prophylaxis (presumably post-exposure prophylaxis)? From the United States, erythromycin resistant strains of *Bordetella pertussis* have been reported (1). In vitro, gyrase inhibitors fluoroquinolones have been found to be highly active, even for macrolide resistant strains; they are therefore an alternative treatment option at least in adults (2). However, a word of warning is in order against the uncritical use of azithromycin because of possible selection of resistant pneumococci. This has occurred after only a single dose of azithromycin (3). Since resistant strains exist, cultures and sensitivity testing should be the way forward.

DOI: 10.3238/arztebl.2009.0057c

REFERENCES

1. Bartkus JM, Juni BA, Ehresmann K, Miller CA, Sanden GN, Cassiday PK et al.: Identification of a mutation associated with erythromycin resistance in *Bordetella pertussis*: implications for surveillance of antimicrobial resistance. *J Clin Microbiol* 2003; 41: 1167–72.
2. Bourgeois N, Ghnassia JC, Doucet-Populaire F: In vitro activity of fluoroquinolones against erythromycin-susceptible and -resistant *Bordetella pertussis*. *J Antimicrob Chemother* 2003; 51: 742–3.
3. Leach AJ, Shelby-James TM, Mayo M, Gratten M, Laming AC, Currie BJ et al.: A prospective study of the impact of community-based Azithromycin treatment of trachoma on carriage and resistance of *Streptococcus pneumoniae*. *Clin Infect Dis* 1997; 24: 356–62.
4. Riffelmann M, Littmann M, Hellenbrand W, Hülbe C, Wirsing von König CH: Pertussis: Not Only a Disease of Childhood. *Dtsch Arztebl Int* 2008; 105(37): 623–8.

Prof. Dr. med. Pramod M. Shah

Gutzkowstr. 69
60594 Frankfurt am Main, Germany
infektiologie@online.de

Conflict of interest statement

The author has received honoraria as a speaker and adviser from Sanofi-Aventis, Janssen-Cilag, and Pohl-Boskamp.

In Reply:

We are delighted that our article has sparked such a lively debate. Dr Mülleneisen reminds us of problems in everyday medical practice: Serological tests for *Bordetella pertussis* often entail IgM antibody measurements. In contrast to other infections, however, the importance of IgM antibodies is questionable, since these antibodies react mainly with the lipo-oligosaccharide and other components of the cellular wall and are therefore often false positives. As we explained in our article, measuring IgG and IgA antibodies to pertussis toxin (PT) is the best validated method and is least prone to non-specificities.

For treatment with antibiotics, we did mention the problem described by Mülleneisen. Antibiotics are not effective for symptom relief in patients who have had a cough for a while, but they may reduce transmission of the infection.

Dr Windhövel describes that administering promethazine and dihydrocodeine yielded success in an adult patient. Promethazine has been mentioned in several earlier case reports, although its efficacy cannot be assessed; Dihydrocodeine has mostly been classified as ineffective. Other possible cough relieving drugs such as diphenhydramine, salbutamol, dexamethasone, and pertussis immunoglobulin have been investigated in controlled studies in children, but no proof of efficacy has been found (1).

Controlled studies of treatment in adults are lacking. This does not exclude that medications may be effective in individual cases; but no therapeutic recommendations can be concluded from such cases.

The frustration about the lack of drug treatment to relieve the painful symptoms resulted in so called "whooping cough flights" in the middle of the 20th century, during which patients were exposed to a period of lower oxygen pressure in an airplane.

Professor Shah worries that using azithromycin in pertussis may result in increased resistance of *Streptococcus pneumoniae* to this macrolide. The current S3 guideline recommends macrolides for respiratory tract infections as the primary alternative to aminopenicillins if antibiotic treatment is indicated (2).

A recently published study found more macrolide resistant alpha haemolytic streptococci immediately after azithromycin had been administered, but this effect had disappeared after 12 weeks (3). Erythromycin resistant strains of *B. pertussis* have been described in the US; these are caused by a mutation to the binding site in the bacterial 23S ribosomal RNA gene. This mutation was found as early as in the 1980s in individual strains; but all strains tested in Europe so far have been sensitive to

erythromycin. The resistance situation vis-à-vis macrolides should be monitored in future by the national reference laboratories; however, general testing of all isolates still does not seem indicated. Fluoroquinolones are effective against *B. pertussis* in vitro, but clinical studies of their use in adults are lacking.

In sum, the discussion shows again that the only effective measure against pertussis is preventive immunization/vaccination of all susceptible persons.

DOI: 10.3238/arztebl.2009.0058

REFERENCES

1. Pillay V, Swingle G: Symptomatic treatment of the cough in whooping cough. *Cochrane Database Syst Rev* 2004; CD003257.
2. S3-Leitlinie: Epidemiologie, Diagnostik, antimikrobielle Therapie und Management von erwachsenen Patienten mit ambulant erworbenen tiefen Atemwegsinfektionen (akute Bronchitis, akute Exazerbation einer chronischen Bronchitis, Influenza und andere respiratorische Virusinfektionen sowie ambulant erworbener Pneumonie). www.uni-duesseldorf.de/AWMF
3. Eisenblätter M, Klaus C, Pletz MWR, Orawa H, Hahn H, Wagner J, Lode H: Influence of azithromycin and clarithromycin on macrolide susceptibility of viridans streptococci from the oral cavity of healthy volunteers. *Eur J Clin Microb Inf Dis* 2008; 27: 1087–92.
4. Riffelmann M, Littmann M, Hellenbrand W, Hülbe C, Wirsing von König CH: Pertussis: Not Only a Disease of Childhood. *Dtsch Arztebl Int* 2008; 105(37): 623–8.

Prof. Dr. med. Carl Heinz Wirsing von König

Institut für Hygiene und Labormedizin
HELIOS Klinikum Krefeld
Lutherplatz 40
47805 Krefeld, Germany
carlheinz.wirsingvonkoenig@helios-kliniken.de

Conflict of interest statement

The author declares that no conflict of interest exists according to the guidelines of the International Committee of Medical Journal Editors.