CORRESPONDENCE

Management of Patients With Aortic Dissection

by PD Dr. med. Ernst Weigang, Prof. Dr. med. Christoph A. Nienaber, Dr. med. Tim C. Rehders, Prof. Dr. med. Hüseyin Ince, Prof. Dr. med. Christian-Friedrich Vahl, Prof. Dr. med. Friedhelm Beyersdorf in volume 38/2008

Etiology Was Incomplete

The explanation of the etiology of aortic dissection surprised me, because more than a year ago I admitted a very young patient with mucoid media degeneration—Erdheim-Gsell syndrome—for aortic surgery to Mainz University Hospital, after I had found extensive widening of the aortic root on the echocardiogram. In addition to Marfan syndrome and Ehlers-Danlos syndrome, this should have been included as a possible etiology.

DOI: 10.3238/arztebl.2009.0171a

REFERENCES

1. Weigang E, Nienaber CA, Rehders TC, Ince H, Vahl C-F, Beyersdorf F: Management of patients with aortic dissection. Dtsch Arztebl Int 2008; 105(38): 639–45.

Dr. med. Achim Küppers
Heidelberger Str. 41
64285 Darmstadt, Germany
AXXDOC2@aol.com

New Insights

We think that some aspects of the article should be rectified, and that some additions are required.

Vascular Ehlers-Danlos syndrome (EDS) is not the most common form of EDS since 80–90% of all cases of EDS are “classic” EDS. The categorization of EDS cited by the authors, into 11 forms has been obsolete for 10 years. The authors do not mention Loeys-Dietz syndrome in their review article, which has a rather more aggressive clinical course than Marfan syndrome. Children with this syndrome often have a split uvula and hypertelorism. In patients with such findings, echocardiography should be considered.

We also missed the modalities combining surgery and catheterization (“frozen elephant trunk”) from the spectrum of surgical therapeutic options, which are used to replace the ascending aorta, the aortic arch, and the descending aorta in one session and which usually save the patient from having to undergo a second operation.

Unfortunately the authors do not discuss the importance of the angiotensin II receptor blocker losartan, the most important advance in the long term management of patients with aortic disorders in years. Mouse models have shown that losartan can prevent the development of aortic aneurysms (1), and recently, the New England Journal of Medicine reported positive outcomes for losartan in patients with Marfan syndrome (2).

In the light of these data it seems increasingly

questionable whether beta blocker monotherapy reflects the current state of knowledge. With regard to the wide spectrum of readers of the Deutsches Ärzteblatt, we wish to point out these new insights.

DOI: 10.3238/arztebl.2009.0171b

REFERENCES

1. Habashi JP, Judge DP, Holm TM et al.: Losartan, an AT1 antagonist, prevents aortic aneurysm in a mouse model of Marfan syndrome. Science 2006; 312: 117–21.
2. Brooke BS, Habashi JP, Judge DP, Patel N, Loeys B, Dietz HC 3rd: Angiotensin II blockade and aortic-root dilation in Marfan's syndrome. N Engl J Med 2008; 358: 2787–95.
3. Weigang E, Nienaber CA, Rehders TC, Ince H, Vahl C-F, Beyersdorf F: Management of patients with aortic dissection. Dtsch Arztebl Int 2008; 105(38): 639–45.

Dr. med. Florian S. Schoenhoff
Proteomics Center
Johns Hopkins University, Baltimore
21224 Maryland USA

PD Dr. sc.nat. Gabor Matyas
Institut für Humangenetik
Universität Zürich
8603 Schwerzenbach, Switzerland

Prof. Dr. med. Thierry Carrel
Klinik für Herz- und Gefäßchirurgie
Universitätsspital Bern
3010 Bern, Switzerland
thierry.carrel@insel.ch

In Reply:

Our article does not claim to be an all inclusive, comprehensive textbook chapter that includes all aspects of aortic dissections. Its objective was to provide the wide readership of the Deutsches Ärzteblatt with an overview, especially of the treatment of aortic dissection. We thank our correspondents for their comments and additions to our article. Our comments follow.

Etiology of aortic dissection—In mucoid media degeneration, excess mucopolysaccharides occur in the connective tissue of the aortic wall (in histopathological specimens after aortic surgery). This histopathological finding applies to Marfan syndrome (MFS), for example, but does not confirm the diagnosis.

Incidence of vascular Ehlers-Danlos syndrome (EDS)—1:10 000–25 000 people develop EDS. According to the Villefranche classification, six clinical types of EDS can be distinguished. The classic type of EDS is the most common form. Vascular type IV EDS affects 5–10% of patients (1).

Loeys-Dietz syndrome (LDS)—LDS is a rare etiology of aortic dissection with phenotypical overlaps with the vascular type of EDS (2). After mutation, a complex pathology develops which is mainly accompanied by aortic aneurysms at risk of rupturing. If patients with LDS are not operated on prophylactically by a vascular surgeon, they tend to die early (average age 26 years) from vascular complications (2).

Surgical therapeutic option “frozen elephant trunk”—In expert circles, combined therapy (open surgery in the area of the ascending aorta and arch, and endovascular surgery by stent graft in the region of the descending aorta) is being discussed as a one-time procedure, which is already being conducted in some specialized centers. Which stent graft prosthesis should be used and whether only a certain hybrid stent (stent graft with integrated vascular Dacron prosthesis) should be used is not clear at this time. Since the procedure is in its clinical testing stages and long term results are not yet available, we decided not to include this therapeutic option in our article.

Medical long term therapy—Treatment with beta blockers has for many years been the standard long term treatment of patients after aortic dissection and patients with Marfan syndrome. Animal model have shown a positive effect of angiotensin II receptor blockers (losartan) in inhibiting the growth of aortic aneurysms (3). This effect was also seen in 18 children with MFS in a non-randomized observational study (4). A general therapeutic recommendation for patients after aortic dissection or patients with MFS cannot be concluded from this study situation. A prospective randomized study, which compares losartan with atenolol in MFS patients, has been running for about a year. After this study has

been evaluated it may be possible to deduce a new therapeutic recommendation for patients with Marfan syndrome.
DOI: 10.3238/arztebl.2009.0172

REFERENCES

1. Germain DP: Ehlers-Danlos syndrome type IV. *Orphanet J Rare Dis* 2007; 2: 32.
2. Loeys BL, Schwarze U, Holm Tet al.: Aneurysm syndromes caused by mutations in the TGF-beta receptor. *N Engl J Med* 2006; 355: 788–98.
3. Habashi JP, Judge DP, Holm TM et al.: Losartan, an AT1 antagonist, prevents aortic aneurysm in a mouse model of Marfan syndrome. *Science* 2006; 312: 117–21.
4. Brooke BS, Habashi JP, Judge DP, Patel N, Loeys B, Dietz HC 3rd.: Angiotensin II blockade and aortic-root dilation in Marfan's syndrome. *N Engl J Med* 2008; 358: 2787–95.
5. Weigang E, Nienaber CA, Rehders TC, Ince H, Vahl C-F, Beyersdorf F: Management of patients with aortic dissection. *Dtsch Arztebl Int* 2008; 105(38): 639–45.

PD Dr. med. Ernst Weigang

Klinik und Poliklinik für Herz-, Thorax- und Gefäßchirurgie
Klinikum der Johannes Gutenberg-Universität Mainz
Langenbeckstr. 1, 55131 Mainz, Germany
weigang@uni-mainz.de

Conflict of interest statement

The authors of all letters and of the reply declare that no conflict of interest exists according to the guidelines of the International Committee of Medical Journal Editors.