Published in final edited form as:

Epilepsy Behav. 2009 March ; 14(3): 540–544. doi:10.1016/j.yebeh.2009.01.005.

Seizures in the elderly: Impact on mental status, mood and sleep

Sheryl R. Haut, M.D.^{1,2}, Mindy Katz, M.P.H.³, Jonathan Masur^{1,2}, and Richard B. Lipton, M.D.^{2,3}

¹ *Comprehensive Epilepsy Management Center, Montefiore Medical Center and the Albert Einstein College of Medicine Bronx, New York*

² *Department of Neurology, Montefiore Medical Center and the Albert Einstein College of Medicine Bronx, New York*

³ *Department of Epidemiology and Population Health, Montefiore Medical Center and the Albert Einstein College of Medicine Bronx, New York*

Abstract

Co-morbidities of epilepsy have not been well explored in the elderly. Herein, we examined mental status, mood, and sleep in elderly patients with epilepsy, compared to age and gender matched community controls without epilepsy from the Einstein Aging Study. Testing included a mental status test, the Blessed Information Memory and Concentration (BIMC) test; Prime-MD Patient Health Questionnaire (PHQ) Depression and Anxiety Modules; and Medical Outcomes Study Sleep Scale. Persons with epilepsy (n=31) had higher mean BIMC scores than controls (n=31, BIMC 6.3 vs. 1.2; $p<0.0001$). Mean PHQ Depression scores were higher for cases than controls indicating more depressive symptoms (4.2 vs. 0.8; $p=0.006$); six cases (18%) and no controls met screening criteria for depression. Mean PHQ Anxiety scores were also higher for cases than controls (3.7 vs. 0.0; $p=0.001$). Cases demonstrated poorer sleep scores in the categories of somnolence ($p=0.009$) and shortness of breath/headache ($p=0.021$). Thus, co-morbidities of epilepsy in this elderly population included decreased mental status, a higher prevalence of depression and anxiety, and poorer sleep health when compared to age mates without epilepsy. Mental status impairment was not related to anti-epileptic medication or mood disturbance. Further investigation will explore these associations prospectively.

Keywords

Epilepsy; elderly; epilepsy co-morbidity; cognitive decline; depression; sleep disturbance

INTRODUCTION

Elderly persons experience epilepsy at age-specific rates higher than any other age group [1–2]. They may be especially vulnerable to the sequelae of seizures and their treatment, and may experience worsened health-related quality of life as compared to their age-mates [3–5].

Furthermore, well-recognized co-morbidities of epilepsy such as depression, anxiety, and sleep

Corresponding Author: Sheryl R. Haut MD, Epilepsy Management Center, Montefiore Medical Center, 111 East 210th Street, Bronx, New York, 10467-2490, Phone: (718) 920-4898, Fax: (718) 882-0216, e-mail: E-mail: haut@aeccmsu.edu.

Presented in part at the American Epilepsy Society Annual Meeting, San Diego CA 2006

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

disorders [6–9] occur with increasing frequency with advancing age [10–12] and merit examination in this special population [13]. Few studies have examined these comorbidities of epilepsy in the elderly. Furthermore, while late stage dementia is an identified risk factor for seizures and epilepsy [14–15], the association in the opposite direction is less clear, and the issue of cognitive and brain health in the elderly with epilepsy has recently been described as neglected and poorly understood [16].

Herein we explored the association between epilepsy, mental status impairment and other selected co-morbidities in our elderly, community dwelling population, also addressing the potential contribution of anti-epileptic medication. We hypothesized that the prevalence of mental status dysfunction, sleep disturbance and mood disorders would be increased in persons with epilepsy in comparison with their age-mates free of seizures. Confirmation of these comorbidities in elderly persons with epilepsy would have important implications in clinical practice. It would imply that clinicians who treat the elderly with epilepsy should be vigilant to the presence of these other important conditions; particularly as sleep and mood disorders are treatable. Furthermore, if mental status impairment in epilepsy is a presenting feature of dementia, earlier detection may contribute to improved outcome. Finally, demonstration of comorbidities creates opportunities to explore the mechanisms that link these disorders.

METHODS

Subjects

Subjects were recruited from two sources serving the same community in the Bronx: the Epilepsy Management Center at Montefiore Medical Center, and the Einstein Aging Study (NIH PO1 AG03949 (RBL)). The Epilepsy Management Center includes the Faculty Practice and the Seizure clinic. The Einstein Aging Study (EAS) is a longitudinal study of systematically recruited community dwelling elderly adults. Subjects are followed annually with medical and epidemiologic histories, neuropsychological testing as well as general medical and neurological examinations. This study was performed with the approval of the Montefiore Medical Center (MMC) Institutional Review Board and the Albert Einstein College of Medicine (AECOM) Committee on Clinical Investigations (CCI). All subjects signed informed consent.

Persons with epilepsy (cases) were recruited both from the Epilepsy Management Center and the Einstein Aging Study. All case were age 65 or older and had a diagnosis of epilepsy confirmed by an epileptologist (SRH), by a review of clinical records applying standard criteria [17].

Controls were recruited from the EAS, did not have a history of seizures and were matched by gender and age (within 5 years) to the cases. Epilepsy was the only basis for excluding individuals from the control group. The EAS controls chosen for the present study are representative of the overall EAS study population. EAS subjects overall take 2.75 ± 2.50 medications; the EAS controls in this study take $3.03 (\pm 1.90)$ medications. Rates of the 5 most common medical conditions among the EAS control group (Hypertension, diabetes, cancer, myocardial infarction, claudication) are similar to rates in the overall EAS study. Both cases and controls were free of DSM-IV defined dementia [18], and had scores on the Blessed Information Memory and Concentration (BIMC) test of less than 8 [19]. The data were prospectively collected, with the current analysis as a planned post-hoc analysis.

Definition of outcomes

Mental Status—The BIMC test was used to measure mental status [20]. Items missed or answered incorrectly are scored so that higher scores indicate increasing mental status

impairment. A score of 4 or greater on the BIMC suggests cognitive dysfunction and scores of 8 or greater indicate dementia or delirium [19]. The BIMC was used because it has long been utilized in the Einstein Aging Study. The instrument was originally selected because of its demonstrated correlation with pathologic markers of dementia. The BIMC correlates very highly with the more widely used Mini Mental Status Exam (MMSE) [21].

Depression and Anxiety—Modules from the Patient Health Questionnaire were used to assess depression and anxiety [22–23]. These self-report questionnaires are widely used and specifically validated in the elderly and in person with a broad range of medical comorbidities [23–24]. These modules assess the frequency of specific symptoms of depression and anxiety; higher scores indicate greater symptomology. The 9-item PHQ depression screener has an empirical cut-score of 15 or greater, which identifies DSM-IV defined Major Depressive Disorder of at least moderate severity. The Prime-MD screener for anxiety is a 3-item assessment questionnaire. A positive endorsement to any one of the three items is defined as a positive screen for anxiety. While disease specific psychiatric screening tools are available for epilepsy [25] they are designed for use in persons with epilepsy and not suitable for comparing epilepsy cases and controls. We chose instruments that were validated in the elderly that would be more likely to operate similarly in both the case and control samples. The PHQ tools are validated in the elderly and robust in face of a number of concomitant illnesses [23–24].

Sleep Disturbances—The Medical Outcomes Study (MOS) Sleep Scale was used to examine various aspects of sleep [26]. The MOS measures sleep patterns and disturbances on 7 different scales. Answers are based on a retrospective assessment over the previous 4 weeks, and each domain is computed on a range of 0–100, with a higher score more indicative of the aspect being measured.

Epilepsy variables—In the intake interview, seizure frequencies in the prior month and the prior six months were ascertained. Those who experienced no seizures during the prior 6 months were considered “well controlled”. Current anti-epileptic drugs (AEDs) were recorded. Epilepsy was classified according to International League Against Epilepsy Guidelines [17].

Statistical Analysis

Scores on the BIMC, Prime MD PHQ Depression and Anxiety Scales, and the (MOS) Sleep Scale were treated as continuous variables. Epilepsy control was treated as a dichotomous variable (well controlled vs. not-well controlled) as was anti-epileptic medications (0–1 vs. >1). Differences in mean scores for continuous variables were tested for significance using Student’s t-test for paired samples (case-control comparison), independent samples (case comparison) if assumptions were met, or Mann Whitney U-test for nonparametric evaluation [27]. All analyses were computed using SPSS [28].

RESULTS

Data were analyzed for 62 subjects, 31 epilepsy cases and 31 controls. Demographic data is presented in Table 1. Groups did not differ in age or gender distribution. The mean age for subjects overall was 74.6 $\pm$ 6.4. Sixty seven percent of the subjects were female. Degree of independent living for cases and controls is indicated. Cases were significantly more likely than controls to live with a child rather than alone or with a spouse ($p < 0.005$).

Epilepsy etiology, classification and control

Epilepsy was localization-related in all cases; none of the cases had a primary generalized epilepsy syndrome. Thirteen subjects (42%) had remote symptomatic epilepsy, including five

subjects with stroke, and four subjects each with post-traumatic epilepsy, and tumor/vascular anomalies. Eighteen subjects (58%) had no epilepsy etiology identified and were considered idiopathic/cryptogenic. Three of these subjects have been diagnosed with mild cognitive impairment (MCI) subsequent to this study.

Age of epilepsy onset ranged from 6–85 years, and duration ranged from 2–63 years. Mean/median age of epilepsy onset was 51 years/58 years $\pm$ 20 years. Thirteen percent of subjects developed epilepsy prior to age 21. Twenty five percent of subjects experienced epilepsy onset by age 32; fifty percent experienced epilepsy onset by age 58. Over sixty percent of the cases had a duration of epilepsy for >10 years.

Forty six percent of cases with epilepsy were currently well controlled, while fifty four % had experienced at least one seizure during the prior six months. Mean number of seizures during the prior month prior to testing was 0.5, and the median was 0.

Anti-epileptic medication use is indicated (Table 1). Seventy five % of cases were receiving monotherapy with one AED; nearly half of these were being treated with a second-generation agent (lamotrigine or levetiracetam). Control subjects were not receiving AED therapy for other indications.

Mental status—Mean BIMC for the group was 3.82 ($\pm$ 5.1), median 2.0 (interquartile range 2.0 to 5.3). Median BIMC for cases with epilepsy was 5.0, as compared to 1.0 for controls (Table 2). Cases with epilepsy had significantly worse BIMC scores than controls (Mann Whitney U-test $p < 0.0001$). When controlling for presence of depression or anxiety, BIMC scores remained significantly worse for persons with epilepsy than for controls. Among persons with epilepsy, BIMC scores were not significantly associated with number of antiepileptic medications (0–1 vs. 2), or degree of seizure control (Mann Whitney U-test n.s.).

Depression and Anxiety—Mean Prime-MD Depression scores were significantly higher for cases (4.2 $\pm$ 5.4) than controls (0.8 $\pm$ 1.8; $p = 0.006$) with six of 31 case subjects (18%) and no control subjects meeting screening criteria for depression (Table 3). Among cases, subjects with poor seizure control had a significantly higher mean depression score (5.57 $\pm$ 5.4) than subjects with well-controlled seizures (0.38 $\pm$ 1.0) ($p < 0.0001$).

Mean Prime-MD Anxiety Scores were also significantly higher for cases (3.73) than controls (0.0) ($p < 0.0001$), although no subject met screening criteria for anxiety disorder (Table 3). Within cases, mean scores were higher but did not significantly differ for poorly controlled (5.75 $\pm$ 8.9) vs. well-controlled (1.8 $\pm$ 3.1) epilepsy ($p = 0.116$).

Sleep disorders—Of the 7 domains measured by the MOS Sleep Scales, (Adequacy, Disturbances-4 item, Somnolence, Snoring, Shortness of Breath/Headache, Problems-6 item, Problems 9-item) cases demonstrated significantly poorer sleep scores than controls in the categories of somnolence ($p = 0.009$) and shortness of breath/headache ($p = 0.021$) scales (Table 3). Among cases, poor seizure control was associated with shortness of breath/headache ($p = 0.006$) and general disturbances ($p = 0.047$). Mental status was not significantly different for subjects with and without significant somnolence or shortness of breath ($p = 0.34$).

DISCUSSION

There is an increasing recognition that elderly patients with epilepsy represent a unique population with particular needs [16,29–30]. Most studies in epilepsy have not focused on this age group, and few have examined multiple co-morbidities. Herein, we evaluated several well-known comorbidities of epilepsy whose prevalence also increases with advancing age.

In our sample of community dwelling elderly, mental status as measured by the BIMC was significantly impaired in cases with localization-related epilepsy as compared to controls without epilepsy. As a cross-sectional finding, these results are consistent with previous cross-sectional studies specifically addressing cognition in the elderly population with epilepsy [29,31–33]. The relationship of cognitive dysfunction to duration of epilepsy remains unclear. A number of prospective studies have demonstrated cognitive decline in association with chronic epilepsy [34–35], although in a review of all available longitudinal studies performed prior to 2004 [36], none had focused on the elderly, and 12 of 20 studies supported the finding of cognitive decline while 8 did not. In one recent longitudinal study of cognitive functioning in the elderly persons with partial epilepsies, overall cognitive deficit appeared to remain fairly stable over time [37], although executive dysfunction progressed.

In our study, cognitive deficits in the elderly with epilepsy were not associated with seizure burden. In contrast, studies of the general epilepsy population suggest that a longer duration of epilepsy or poor seizure control is significantly related to cognitive decline [38–41]. Seizure control over the last 6 months may not adequately measure lifetime seizure burden particularly in the elderly; we are therefore not willing to exclude the possibility that seizures cause cognitive impairment. Better measures of seizure burden and cognitive status, particularly in a longitudinal context might reveal associations not detected here.

Polypharmacy has been implicated as a risk factor for cognitive impairment; in prior studies, the number of AEDs was negatively associated with cognition, both in comparison to healthy controls [31] and even in comparison to older adults with mild cognitive impairment taking anti-cholinesterase inhibitors [32]. In these studies, the percent of patients taking monotherapy ranged from 52–57%. In our population, the number of AEDs was not significantly associated with poorer performance on the BIMC. This may in part reflect the high proportion (75%) of subjects taking monotherapy, and the high prevalence of newer anti-epileptic agents, which may produce less cognitive impairment than older agents [42]; these agents are better tolerated, especially in the elderly [43]. However, as the control group was not taking anti-epileptic medication, the effects of AEDs may have contributed to the difference between the groups. The presence of cognitive impairment in cases may explain the differences in living arrangements between cases and controls. Although all subjects were community dwellers, a significantly higher percentage of cases with epilepsy lived less independently than controls.

Psychiatric comorbidities, particularly mood disturbances such as depression and anxiety disorders are associated with epilepsy [7,44–48] and increase in prevalence with advancing age [11–12]. In our study, elderly persons with seizure disorders suffered from elevated rates of depression and anxiety as compared to controls. Several measures have found that depression or other psychological stress may be the strongest predictors of health-related quality of life for persons with epilepsy [49], highlighting the importance of these co-morbidities in the elderly population with epilepsy. Furthermore, antiepileptic medications have the potential to produce or worsen mood disorders [50–51], and in our study, it is difficult to separate the effect of medication from the effect of epilepsy. However, the elderly may be particularly sensitive to these medication effects, underscoring the need to evaluate mood in this population.

The overall rate of depression and anxiety in our study appears lower than other reports of depression in epilepsy; this may be related to the relatively long duration of epilepsy in our cases, as later age of onset of epilepsy appears to be a risk factor for mood disorders [52].

Sleep health is particularly important to address in the elderly with epilepsy, as seniors in the general population have a higher degree of sleep disturbance than younger individuals [53–54]. Sleep disturbance is an increasingly noted co-morbidity of epilepsy [8]. Sleep disorders such as insomnia and excessive daytime sleepiness lower quality of life measures for persons

with epilepsy [55], and sleep deprivation is considered a seizure precipitant in numerous reports [57–60]. Obstructive sleep apnea has recently been linked to increased seizure frequency in the elderly [61]. However, to our knowledge, general sleep disturbance has not been previously investigated in the elderly with epilepsy. We found a strong correlation between reported sleep health, the presence of epilepsy, and current degree of seizure control. Our elderly subjects with epilepsy had significantly elevated scales measuring sleep somnolence and shortness of breath as compared to cases. Although previously reported [54,56], sleep disturbance in our study was not accounted for by depressed mood. Poor sleep may also contribute to cognitive decline [62–63], although in our study, sleep disturbance was not associated with cognitive impairment. It is possible that improving sleep health in this age group will have a positive effect on both epilepsy control and cognition.

Our study has a number of limitations. It is a cross sectional study, and so cannot determine the temporal relationships between epilepsy and the examined co-morbidities. Planned prospective studies will address these concerns. In addition, though our epilepsy cases and controls are matched for age and sex and drawn from the same community it is possible that selection bias influences study results. Our cognitive assessment was limited to an excellent test of mental status, which correlates very highly with the better known Minimental State Exam [21]. It is possible that a battery of more robust neuropsychological measures would reveal differences not apparent here. We could not fully control for medication effects, including the potential effects of antiepileptic medication on cognition and mood. Finally, more than half of the cases with epilepsy were not well controlled; this sample may be more refractory to treatment than the general epilepsy population. This finding is likely accounted for by the selection of cases from an epilepsy tertiary care center. The applicability of our results to the elderly with epilepsy followed in general neurology practice should be explored.

The elderly with epilepsy are not a homogeneous population. Prior studies of epilepsy have demonstrated that significant differences exist between community dwellers and nursing home residents [64]. While epilepsy co-morbidities such as cognitive impairment may be expected to be high in an assisted living population, the subjects in our study represent a community-based sample; as such, the findings are perhaps even more significant. Future planned studies will utilize a more extensive cognitive assessment targeted to specific cortical regions (i.e. frontal lobe executive function) to further enhance our understanding of the relationship between cognition and epilepsy.

Thus, our results confirm that the elderly with epilepsy suffer from numerous clinically significant co-morbidities, including mental status impairment, mood disturbance, and poorer sleep health. These co-morbidities may impact on independent living and quality of life. In treating elderly patients with epilepsy, maintaining functional independence is an important goal. It is therefore important for treating clinicians to be aware of these common co-morbidities and address them. It is also important to determine if the prominent mental status impairment is a consequence of epilepsy or of its treatment so that management strategies that optimize cognitive function can be developed.

Acknowledgments

The authors acknowledge funding support by NIH P01 AG03949 (PI: Dr. Lipton), NIH K23 NS02192 (PI: Dr. Haut), and an unrestricted grant from Pfizer (PI: Dr. Haut). Dr. S.R. Haut serves on the speaker's bureau of UCB Pharma and the advisory boards of UCB Pharma and Abbott. She has served as a paid consultant for King, Jazz, and Genactis. Dr. R.B. Lipton has served as a paid consultant for, and received grant support from GSK. None of the other authors have any conflicts of interest to disclose.

The authors confirm that we have read the Journal's position on issues involved in ethical publication, and affirm that this report is consistent with those guidelines.

Funding supported by NIH P01 AG03949 (PI: Dr. Lipton), NIH K23 NS02192 (PI: Dr. Haut), and an unrestricted grant from Pfizer (PI: Dr. Haut).

References

1. Hauser WA, Annegers JF, Kurland LT. Incidence of epilepsy and unprovoked seizures in Rochester, Minnesota: 1935–1984. *Epilepsia* 1993;34(3):453–468. [PubMed: 8504780]
2. Hussain SA, Haut SR, Lipton RB, Derby C, Markowitz SY, Shinnar S. Incidence of epilepsy in a racially diverse, community-dwelling, elderly cohort: results from the Einstein aging study. *Epilepsy Res* 2006;71:195–205. [PubMed: 16870396]
3. Brodie MJ, Kwan P. Epilepsy in elderly people. *BMJ* 2005;331(7528):1317–1322. [PubMed: 16322020]
4. Cloyd J, Hauser W, Towne A, Ramsay R, Mattson R, Gilliam F, Walczak T. Epidemiological and medical aspects of epilepsy in the elderly. *Epilepsy Res* 2006;68:S39–S48. [PubMed: 16384689]
5. Laccheo I, Ablah E, Heinrichs R, Sadler T, Baade L, Liow K. Assessment of quality of life among the elderly with epilepsy. *Epilepsy Behav* 2008;12(2):257–261. [PubMed: 17988950]
6. Piazzini A, Paola Canevini M, Maggiori G, Canger R. Depression and anxiety in patients with epilepsy. *Epilepsy Behav* 2001;2(5):481–489. [PubMed: 12609287]
7. Jones JE, Hermann BP, Barry JJ, Gilliam F, Kanner AM, Meador KJ. Clinical assessment of Axis I psychiatric morbidity in chronic epilepsy: a multicenter investigation. *J Neuropsychiatry Clin Neurosci* 2005;17(2):172–9. [PubMed: 15939970]
8. Malow BA. The interaction between sleep and epilepsy. *Epilepsia* 2007;48 (Suppl 9):S36–38.
9. Kanner AM. Epilepsy and mood disorders. *Epilepsia* 2007;48(Suppl 9):S20–22.
10. Foley DJ, Monjan AA, Brown SL, Simonsick EM, Wallace RB, Blazer DG. Sleep complaints among elderly persons: an epidemiologic study of three communities. *Sleep* 1995;18(6):425–432. [PubMed: 7481413]
11. Steffens DC, Skoog I, Norton MC, Hart AD, Tschanz JT, Plassman BL, Wyse BW, Welsh-Bohmer KA, Breitner JC. Prevalence of depression and its treatment in an elderly population: the Cache County study. *Arch Gen Psychiatry* 2000;57:601–607. [PubMed: 10839339]
12. Schuurmans J, Comijs HC, Beekman AT, de Beurs E, Deeg DJ, Emmelkamp PM, van Dyck R. The outcome of anxiety disorders in older people at 6-year follow-up: results from the Longitudinal Aging Study Amsterdam. *Acta Psychiatr Scand* 2005;111:401–402. [PubMed: 15877705]
13. Leppik IE. Epilepsy in the elderly. *Epilepsia* 2006;47(Suppl 1):S65–70.
14. Hesdorffer DC, Hauser WA, Annegers JF, Kokmen E, Rocca WA. Dementia and adult-onset unprovoked seizures. *Neurology* 1996;46:727–730. [PubMed: 8618673]
15. Amatniek JC, Hauser WA, DelCastillo-Castaneda C, Jacobs DM, Marder K, Bell K, Albert M, Brandt J, Stern Y. Incidence and predictors of seizures in patients with Alzheimer's disease. *Epilepsia* 2006;47:867–872. [PubMed: 16686651]
16. Hermann B, Seidenberg M, Sager M, Carlsson C, Gidal B, Sheth R, Rutecki P, Asthana S. Growing old with epilepsy: the neglected issue of cognitive and brain health in aging and elder persons with chronic epilepsy. *Epilepsia* 2008;49(5):731–40. [PubMed: 18031544]
17. Commission on classification and terminology of the International League Against Epilepsy. Proposal for revised classification of epilepsies and epileptic syndromes. *Epilepsia* 1989;30:389–399. [PubMed: 2502382]
18. Diagnostic and Statistical Manual of Mental Disorders, *DSM-IV*. American Psychiatric Association; Washington, DC: 1994. p. 133
19. Katzman R, Brown T, Thal LJ, Fuld PA, Aronson M, Butters N, Klauber MR, Wiederholt W, Pay M, Xiong RB, Ooi WL, Hofstetter R, Terry RD. Comparison of rate of annual change of mental status score in four independent studies of patients with Alzheimer's disease. *Ann Neurol* 1988;24:38389.
20. Blessed G, Tomlinson BE, Roth M. The association between quantitative measures of dementia and of senile change in the cerebral gray matter of elderly subjects. *Brit J Psychiatry* 1968;114:79811.

21. Thal LJ, Grundman M, Golden R. Alzheimer's disease: a correlational analysis of the Blessed Information-Memory-Concentration Test and the Mini-Mental State Exam. *Neurology* 1986;36(2): 262–4. [PubMed: 3945395]
22. Spitzer RL, Kroenke K, Williams JB. Validation and utility of a self-report version of PRIME-MD: the PHQ primary care study. *Primary Care Evaluation of Mental Disorders. Patient Health Questionnaire*. *JAMA* 1999;282:1737–1744. [PubMed: 10568646]
23. Kroenke K, Spitzer RL, Williams JB. The PHQ-9: validity of a brief depression severity measure. *J Gen Intern Med* 2001;16:606–613. [PubMed: 11556941]
24. Lamers F, Jonkers CC, Bosma H, Penninx BW, Knottnerus JA, van Eijk JT. Summed score of the Patient Health Questionnaire-9 was a reliable and valid method for depression screening in chronically ill elderly patients. *J Clin Epidemiol* 2008;61(7):679–87. [PubMed: 18538262]
25. Devinsky O, Vickrey BG, Cramer J, et al. Development of the Quality of Life in Epilepsy Inventory. *Epilepsia* 1995;36(11):1089–1104. [PubMed: 7588453]
26. Hays RD, Martin SA, Sesti AM, Spritzer KL. Psychometric properties of the Medical Outcomes Study Sleep measure. *Sleep Med* 2005;6:41–44. [PubMed: 15680294]
27. Lehmann, EL. *Nonparametrics: Statistical Methods Based on Ranks*. McGraw Hill International Book Company; New York: 1975.
28. SPSS Base 10.0 Applications Guide. SPSS, Inc.; Chicago, IL: 1999.
29. Ramsay RE, Rowan AJ, Pryor FM. Special considerations in treating the elderly patient with epilepsy. *Neurology* 2004;62(5 Suppl2):S24–9. [PubMed: 15007161]
30. Leppik I. Epilepsy in the elderly: Scope of the problem. *Int Rev Neurobio* 2007;81:1–14.
31. Martin RC, Griffith HR, Faught E, Gilliam F, Mackey M, Vogtle L. Cognitive functioning in community dwelling older adults with chronic partial epilepsy. *Epilepsia* 2005;46:298–303. [PubMed: 15679511]
32. Griffith HR, Martin RC, Bambara JK, Marson DC, Faught E. Older adults with epilepsy demonstrate cognitive impairments compared with patients with amnesic mild cognitive impairment. *Epilepsy Behav* 2006;8:161–168. [PubMed: 16266829]
33. Piazzini A, Canevini MP, Turner K, Chifari R, Canger R. Elderly people and epilepsy: cognitive function. *Epilepsia* 2006;47(Suppl5):S82–4.
34. Helmstaedter C, Kurthen M, Lux S, Reuber M, Elger CE. Chronic epilepsy and cognition: a longitudinal study in temporal lobe epilepsy. *Ann Neurol* 2003;54(4):425–32. [PubMed: 14520652]
35. Hermann BP, Seidenberg M, Dow C, Jones J, Rutecki P, Bhattacharya A, Bell B. Cognitive prognosis in chronic temporal lobe epilepsy. *Ann Neurol* 2006;60(1):80–7. [PubMed: 16802302]
36. Dodrill CB. Neuropsychological effects of seizures. *Epilepsy Behav* 2004;(5 Suppl 1):S21–4. [PubMed: 14725843]
37. Griffith HR, Martin RC, Bambara JK, Faught E, Vogtle LK, Marson DC. Cognitive functioning over 3 years in community dwelling older adults with chronic partial epilepsy. *Epilepsy Res* 2007;74(2–3):91–96. [PubMed: 17368001]
38. Jokeit H, Ebner A. Long term effects of refractory temporal lobe epilepsy on cognitive abilities: a cross sectional study. *J Neurol Neurosurg Psychiatry* 1999;67:44–50. [PubMed: 10369821]
39. Jokeit H, Ebner A. Effects of chronic epilepsy on intellectual functions. *Prog Brain Res* 2002;135:455–63. [PubMed: 12143364]
40. Thompson PJ, Duncan JS. Cognitive decline in severe intractable epilepsy. *Epilepsia* 2005;46:1780–1787. [PubMed: 16302858]
41. Marques CM, Caboclo LO, da Silva TI, Noffs MH, Carrete H Jr, Lin K, Lin J, Sakamoto AC, Yacubian EM. Cognitive decline in temporal lobe epilepsy due to unilateral hippocampal sclerosis. *Epilepsy Behav* 2007;10:477–485. [PubMed: 17368105]
42. Aldenkamp AP, De Krom M, Reijs R. Newer antiepileptic drugs and cognitive issues. *Epilepsia* 2003;44(Suppl 4):S21–29.
43. Rowan AJ, Ramsay RE, Collins JF, Pryor F, Boardman KD, Uthman BM, Spitz M, Frederick T, Towne A, Carter GS, Marks W, Felicetta J, Tomyanovich MLVA. Cooperative Study 428 Group. New onset geriatric epilepsy: a randomized study of gabapentin, lamotrigine, and carbamazepine. *Neurology* 2005;64(4):1868–1873. [PubMed: 15955935]

44. Kanner AM, Nieto JC. Depressive disorders in epilepsy. *Neurology* 1999;53(5 Suppl 2):S26–32. [PubMed: 10496231]
45. Hermann BP, Seidenberg M, Bell B. Psychiatric Co-Morbidity in Chronic Epilepsy: Identification, Consequences and Treatment of Major Depression. *Epilepsia* 2000;41(Suppl 2):S31–41. [PubMed: 10885738]
46. Gupta, AK.; Ettinger, AB.; Weisbrot, DM. Psychiatric comorbidities in epilepsy. In: Ettinger, AB.; Devinsky, O., editors. *Managing epilepsy and co-existing disorders*. Butterworth-Heinemann; Boston: 2002. p. 343-387.
47. Johnson EK, Jones JE, Seidenberg M, Hermann BP. The relative impact of anxiety, depression, and clinical seizure features on health-related quality of life in epilepsy. *Epilepsia* 2004;45(5):544–550. [PubMed: 15101836]
48. Kobau R, Gilliam F, Thurman DJ. Prevalence of self-reported epilepsy or seizure disorder and its associations with self-reported depression and anxiety: results from the 2004 Healthstyles Survey. *Epilepsia* 2006;47(11):1915–1921. [PubMed: 17116032]
49. Gilliam F, Hecimovic H, Sheline Y. Psychiatric comorbidity, health, and function in epilepsy. *Epilepsy Behav* 2003;4(Suppl 4):S26–S30. [PubMed: 14654425]
50. Meador KJ, Gilliam FG, Kanner AM, Pellock JM. Cognitive and Behavioral Effects of Antiepileptic Drugs. *Epilepsy Behav* 2001;2(4):SS1–SS17. [PubMed: 12609218]
51. French JA, Kanner AM, Bautista J, Abou-Khalil B, Browne T, et al. Efficacy and Tolerability of the New Antiepileptic Drugs, II: Treatment of Refractory Epilepsy: Report of the TTA and QSS Subcommittees of the American Academy of Neurology and the American Epilepsy Society. *Epilepsia* 2004;45(5):410–423.
52. Jagadheesan K, Garg AK, Haque Nizamie S. Risk factors and outcome of mood disorders in epilepsy: a case-control study. *Seizure* 2005;12:121–125. [PubMed: 12566237]
53. Ganguli M, Reynolds CF, Gilby JE. Prevalence and persistence of sleep complaints in a rural older community sample: the MoVIES project. *J Am Geriatr Soc* 1996;44:778–784. [PubMed: 8675924]
54. Newman AB, Enright PL, Manolio TA, Haponik EF, Wahl PW. Sleep disturbance, psychosocial correlates, and cardiovascular disease in 5201 older adults: the Cardiovascular Health Study. *J Am Geriatr Soc* 1997;45:1–7. [PubMed: 8994480]
55. Piperidou C, Karlovasitou A, Triantafyllou N, Terzoudi A, Constantinidis T, Vadikolias K, Heliopoulos I, Vassilopoulos D, Balogiannis S. Influence of sleep disturbance on quality of life of patients with epilepsy. *Seizure* 2008;17(7):588–94. [PubMed: 18396419]
56. Foley DJ, Monjan A, Simonsick EM, Wallace RB, Blazer DG. Incidence and remission of insomnia among elderly adults: an epidemiologic study of 6,800 persons over three years. *Sleep* 1999;22(Suppl 2):S366–S372. [PubMed: 10394609]
57. Fountain NB, Kim JS, Lee SI. Sleep deprivation activates epileptiform discharges independent of the activating effects of sleep. *J Clin Neurophysiol* 1988;15:69–75. [PubMed: 9502515]
58. Rajna P, Veres J. Correlations between night sleep duration and seizure frequency in temporal lobe epilepsy. *Epilepsia* 1993;34:574–579. [PubMed: 8504789]
59. Badawy RAB, Curatolo JM, Newton M, Berkovic MD, Macdonell RAL. Sleep deprivation increases cortical excitability in epilepsy. *Neurology* 2006;67:1018–1022. [PubMed: 17000971]
60. Haut SR, Hall CB, Masur J, Lipton RB. Seizure occurrence: precipitants and prediction. *Neurology* 2007;69(20):1905–1910. [PubMed: 17998482]
61. Chihorek AM, Abou-Khalil B, Malow BA. Obstructive sleep apnea is associated with seizure occurrence in older adults with epilepsy. *Neurology* 2007;69(19):1823–7. [PubMed: 17984449]
62. Yaffe K, Blackwell T, Barnes DE, Ancoli-Israel S, Stone KL. Preclinical cognitive decline and subsequent sleep disturbance in older women. *Neurology* 2007;69:237–242. [PubMed: 17636060]
63. Schmutte T, Harris S, Levin R, Zweig R, Katz M, Lipton R. The relation between cognitive functioning and self-reported sleep complaints in nondemented older adults: results from the Bronx aging study. *Behav Sleep Med* 2007;5:39–56. [PubMed: 17313323]
64. Leppik I. Epilepsy in the elderly: Scope of the problem. *Int Rev Neurobio* 2007;81:1–14.

Table 1

Characteristics of the Study Population

	Cases	Controls	Significance
Mean Age in Years	72.8 (+/- 7.1)	76.6 (+/- 5.2)	NS
Gender Distribution (M/F)	34%/66%	34%/66%	NS
Living arrangement			
Lives alone	43%	53%	
Lives with spouse	36%	43%	
Lives with children	18%	4%	p<0.005
Home health aide	3%	0	
Current anti-epileptic medications		N/A	
0	3%		
1	75%		
2	22%		
>2	--		
Anti-epileptic medications in subjects receiving monotherapy:		N/A	
Phenytoin	32%		
Carbamazepine	18%		
Levetiracetam	27%		
Lamotrigine	18%		
Phenobarbital	5%		

Table 2

Cognitive function

	Cases	Controls	Significance
Mean BIMC Score	6.25 (+/- 6.0)	1.23 (+/- 1.3)	p<0.0001*
Median BIMC Score	5.0	1.0	
Cases only:			
Mean BIMC Score by number of AEDs:			
0-1	6.23 (+/- 5.0)		
2	8.5 (+/- 9.8)**		
Mean BIMC Score by seizure control			
Well-controlled	6.9 +/- 6.2		
Poorly-controlled	7.2 +/- 7.4**		

* By Mann Whitney U-test for nonparametric data

** Within column comparisons not significant

Table 3

Mood and Sleep Disturbance

	Cases	Controls	Significance
Prime MD Depression	4.2 (+/- 5.4)	0.8 (+/- 1.8)	p=0.006
Prime MD Anxiety	3.7 (+/- 4.1)	0	P < 0.0001
MOS Sleep Scales:			
Shortness of Breath	13.1 (+/- 23.6)	3.3 (+/- 7.6)	p=0.034
Somnolence	38.4 (+/- 31.0)	22.7 (+/- 13.9)	p=0.013
Sleep Disturbance	22.8 (+/- 26.7)	26.7 (+/- 12.9)	p=0.386
Sleep Adequacy	75.0 (+/- 32.2)	63.5 (+/- 25.0)	p=0.139
Sleep Disturbance	22.8 (+/- 26.6)	26.7 (+/- 12.9)	p=0.386
Sleep Problems-6 item	25.1 (+/- 24.5)	24.4 (+/- 12.4)	p=0.885
Sleep Problems-9 item	25.7 (+/- 22.8)	25.0 (+/- 10.1)	p=0.879