CORRESPONDENCE

Imaging Studies for the Early Detection of Breast Cancer

by Prof. Dr. med. Sylvia Helen Heywang-Köbrunner,
Prof. Dr. med. Ingrid Schreer, Prof. Dr. med. Walter Heindel,
PD Dr. med. Alexander Katalinic in volume 31–32/2008

Imaging Is Not Sufficient

Recently, quality assured mammography screening has been implemented in Germany, which is based on the guidelines of international experts but has important structural differences. Imaging is not the only factor that contributes to successful early detection of breast cancer. Women's confidence is undermined by diverging articles in the press, political campaigns, and different methods such as digital mammography, magnetic resonance tomography, or ultrasonography. The result of all this is that participation rates are still low.

How might the situation in Germany be improved? Early detection (screening) programs cannot be merely transferred from other countries to Germany.

The treatment and often the diagnosis of breast cancer in Germany are in the hands of gynecologists, who are not integrated into mammography screening projects. The gynecologists are not informed about the results of the mammograms which is an unnecessary loss of information and confidence in the system.

In Germany, screening units have not been affiliated to certified breast cancer centers in the same way as in other countries.

Information programs for women have not been conducted to a sufficient degree. Providing information and by doing so eliminating fears—this is an essential improvement for early detection programs.

Known risk groups should be screened differently, not only by means of mammography. The radiation density (>50%) affecting the breast is currently the biggest risk factor. Physical examination by trained personnel is not being undertaken either. How else is it possible that in breast cancer centers, women whose tumors are easily palpable are operated on an almost daily basis?

About half of all women (younger than 50 and older than 69) are not included in mammography screening.

We are in the position of being able to conduct controlled randomized studies of the validity of breast cancer screening. An initial, promising contribution has been made in Schleswig-Holstein. What has been lacking thus far is only the will to jointly succeed.

DOI: 10.3238/arztebl.2009.0145a

Prof. Dr. med. Dr. h. c. Manfred Kaufmann
Frauenklinik der J.-W. Goethe-Universität Frankfurt
Theodor-Stern-Kai 7
60590 Frankfurt/Main, Germany

REFERENCES

- Heywang-Köbrunner SH, Schreer I, Heindel W, Katalinic A: Imaging Studies for the Early Detection of Breast Cancer. Dtsch Arztebl Int 2008; 105(31–32): 541–7.

Bonn Study Was Interpreted Incorrectly

The results from the Bonn study (1) were summarized in a misleading manner. The authors say that the study reported that "high grade ductal carcinoma in situ (DCIS) is often more easily detected by MRI than low grade DCIS". It has been common knowledge for many years that mammography screening obviously helps to identify many high grade DCIS. Rather, the Bonn study showed that half of the high grade DCIS remained undetected on mammography. I do not understand why the authors cite the study listed in reference 20 as contradicting the Bonn study when the results of that study are clearly consistent with those of the Bonn study: MRI shows up many additional, mammographically inapparent DCIS.

Table 1 and the authors' arguments regarding complementary ultrasonography are de facto correct but still misleading. How are readers meant to rate the information that ultrasonography enabled the detection of "4.2/1000 cancers" when it is not explained how many cancers are detected by mammography alone? Table 1 (with regard to the ACRIN 6666 study) (2) remarks that "8.8% of positive findings" were detected by ultrasonography only. Readers may gain the impression that the benefit thus derived is only marginal. In fact, of 41 cancers, only half—20—were detected by mammography alone whereas 12 additional cancers were detected by ultrasonography. This is certainly not a marginal benefit. Nine further cancers were neither detected by mammography nor by ultrasonography but were palpated as interval cancers—which underlines the "reservoir" of cancers that remain to be diagnosed by using MRI.

DOI: 10.3238/arztebl.2009.0145b

Prof. Dr. med. Hans H. Schild
Radiologische Klinik
Friedrich-Wilhelms-Universität Bonn
Sigmund-Freud-Str. 25
53105 Bonn, Germany

REFERENCES

- Kuhl CK, Schrading S, Bieling HB et al.: MRI for diagnosis of pure ductal carcinoma in situ: a prospective observational cohort study. Lancet 2007; 370: 485–92.
- Berg WA, Blume JD, Cormack JB et al.: ACRIN 6666 Investigators. Combined screening with ultrasound and mammography vs. mammography alone in women at elevated risk for cancer. JAMA 2008; 299(18): 2151–63.
- Heywang-Köbrunner SH, Schreer I, Heindel W, Katalinic A: Imaging Studies for the Early Detection of Breast Cancer. Dtsch Arztebl Int 2008; 105(31–32): 541–7.