CORRESPONDENCE

Attention Deficit and Hyperactivity Disorder in Adulthood

by Dr. med. Alexandra Philipsen, Priv.-Doz. Dr. med. Bernd Heßlinger,
Prof. Dr. med. Ludger Tebartz van Elst in volume 17/2008

Exclusion Diagnosis

The phenomenon of underdiagnosed ADHD in adulthood is balanced by the number of (often) misdiagnosed cases of ADHD in children and adolescents. ADHD should always be an exclusion diagnosis—that is, all alternative physical or psychiatric differential diagnoses and psychodynamic treatment approaches should be excluded. Diagnosis and treatment of ADHD should in my opinion be the remit of child and adolescent psychiatrists and psychiatrists. Further to the article, I would like to encourage in addition to symptoms evaluation forms—for example, according to ICD—and their adaptation for adults, "double blind trials" using methylphenidate with the patients' consent, in order to make a diagnosis (placebo versus methylphenidate), so as to have an additional indication of the diagnosis in patients in whom the drug shows positive effects—which is easy to spot owing to its prompt efficacy. Writing samples after taking methylphenidate may also be indicative of a diagnosis if the [patient's handwriting is clearly improved, as well as concentration tests with and without methylphenidate. Thorough diagnostic testing (for depression, intelligence, personality, concentration) is the standard in diagnosing ADHD in children and adolescents and should in my opinion also be considered in adults (for example, by using Beck's depression inventory {BDI}, the Hamburg-Wechsler intelligence test for adults, personality questionnaires, concentration tests). As child and adolescent psychiatrists, some of us are very familiar with the problem of "off label" use for most psychopharmacological preparations, for example in the field of atypical neuroleptics. It is incumbent on the pharmaceutical industry and health policymakers to abolish unnecessary obstacles. I hope that the diagnosis and treatment of ADHD can become established beyond adolescence.

DOI: 10.3238/arztebl.2008.0764a

Giulio Calia
LWL-Klinik Hamm
Heithofer Allee 64
59071 Hamm, Germany
g.calia@wkp-lwl.org

Conflict of interest statement

The author declares that no conflict of interest exists according to the guidelines of the International Committee of Medical Journal Editors.

Causal Treatment for ADHD is Possible

To illustrate the average rate of 2% of ASDHD in adults, I would like to refer to a group of women who

show the clinical lead symptoms more often than might be expected: reduced attention span, inner unrest, disrupted affective regulation, depressive moods, and sometimes resultant disorganization. This group comprises women with severe menopausal symptoms aged around 50, who can be treated with medication without going "off label". Extreme symptoms of extreme estrogen deficiency (which also occurs after castration in the fertile years) are not associated with the clinical diagnosis of ADHD as often as they should. This physical correlation requires increased interdisciplinary exchanges.

From aromatase research (more than 40 years with the enzyme letrozole), estrogen deficiency is known to induced attention deficits, memory deficits, and cognitive deficits. In oncology, the same is known in patients treated with adjuvant aromatase inhibitor therapy. After only one year out of five years' planned therapy, patient compliance reduces to one third. Hence it is not surprising that the brain, which is well furnished with estrogen receptors, is not able to perform enough tasks to a satisfactory degree. Before the extent of ADHD is reached, with impaired quality of life, women decide on non-compliance.

The authors suggest exclusion diagnostics using internal medical and neurological parameters. In women, ovarian function should be taken into consideration. A notably raised prevalence of ADHD has been shown in prisoners, of some 25%. In women, this stress situation often results in amenorrhea—the ovarian function ceases and estrogen deficiency starts to affect the brain.

In men, an analogous reaction develops, for example in prisoners of war—less testosterone is being converted into estrogen. The authors refer to the disrupted dopaminergic system and disrupted synaptic function.

If estrogen treatment is attempted, the increased risk of breast cancer (in the per mille range) has to be highlighted. In the Women's Health Initiative cohort, new cases of breast cancer per 10 000 women were found in 38 women having gestagen therapy and in 30 women taking placebo (3). This difference of 8 women per 10 000 women/year, or 0.8 per mille, has been confirmed by a preceding, large study (1). In the Women's Health Initiative cohort with estrogen monotherapy, 7 fewer cases of breast cancer were detected in 10 000 women compared with placebo, which also means in the per mille range (2). These data will sway the decision in favour of estrogen substitution in ADHD.

DOI: 10.3238/arztebl.2008.0764b

REFERENCES

1. Beral V: Breast cancer and hormone replacement therapy: collaborative reanalysis of data from 51 epidemiological studies of 52,705 women with breast cancer and 108,411 women without breast cancer. *Lancet* 1997; 350:1047–59.

2. WHI Investigators: Risks and benefits of estrogen plus progestin in healthy postmenopausal women. Principal results from the WHI randomized controlled trial JAMA 2002; 288:321–33.
3. WHI Investigators: Effects of conjugated equine estrogen in postmenopausal women with hysterectomy. JAMA 2004; 291: 1701–12.

Prof. Dr. J. M. Wenderlein
Universität Ulm
Eythstr. 14, 89075 Ulm, Germany
wenderlein@gmx.de

Conflict of interest statement

The author declares that no conflict of interest exists according to the guidelines of the International Committee of Medical Journal Editors.

Primary and Secondary ADHD

The authors are to be congratulated for their clear and structured description of the pathology in adulthood. My only criticism is the categorization into primary and secondary ADHD. I think the term "secondary ADHD" is unfortunate and confusing because the problem in question is mostly organic cerebral psychiatric syndromes in the wider sense, which may of course resemble the symptoms of mainly genetically caused ADHD. In a patient with frontal lobe syndrome, one would not talk about "secondary ADHD", even though the patient may meet the criteria in terms of attention deficit, impaired concentration, and impulsiveness (or labile affect). I therefore think the term secondary ADHD is more of a hindrance to the difficult differential diagnostics and the often present comorbidities, for diagnostic as well as therapeutic and prognostic reasons. DOI: 10.3238/arztebl.2008.0765a

Dr. med Andreas Vogel
Lavendelweg 1
66424 Homburg/Saar, Germany
andreasvogel1101@web.de

Conflict of interest statement

The author declares that no conflict of interest exists according to the guidelines of the International Committee of Medical Journal Editors.

ADHD and Consumption of THC

In my everyday activities as an assessor, I often have dealings with young adults who have ADHD and consume tetrahydrocannabinol (THC). I therefore always test for drugs or cannabis in ADHD. I firmly believe that in those who regularly consume THC, their ADHD cannot be improved; and questioning or testing patients makes sense. Some young adults have been prescribed methylphenidate and consume THC on top of that. I would be delighted if you emphasized more strongly the contraindication to prescribing methylphenidate in patients with drug addiction-prescribing physicians should perform regular drug screening tests before issuing the prescription and during treatment with methylphenidate.

DOI: 10.3238/arztebl.2008.0765b

REFERENCES

1. Rösler M, Retz W: Sozialmedizinische Aspekte der ADHS – Über die Ursachen und Folgen komorbider Störungen. PsychoNeuro 2007, 33: 390–4.

Gisela Uhlig
Sudetenring 127
63303 Dreieich, Germany

Conflict of interest statement

The author declares that no conflict of interest exists according to the guidelines of the International Committee of Medical Journal Editors.

In Reply:

Progress assessment under methylphenidate

Much like children, many adults in clinical practice show improvement in their handwriting. Single or double blinded studies (the latter being more difficult to conduct in clinical practice) could be undertaken to prove the necessity of continuing medication. The diagnostic predictive power of the efficacy of methylphenidate, for example in single or double blinded trials is limited, however, owing to the response rate of 75% (1). Even patients who do not respond may have ADHD. Questionnaires and neuropsychological tests can complement diagnostic tests and assessment of progress.

Addiction (THC) and ADHD

As described in the article, addiction disorders are a common comorbidity and differential diagnosis in ADHD in adulthood. Tight control of the drug consumption is indicated. In patients with comorbid addiction, treatment with stimulants should be undertaken only under controlled conditions (drug screening).

ADHD and consumption of THC

Dr Vogel is right to point out that the categorization into primary and secondary ADHD is not generally established and no consensus exists. However, we think that this categorization is valid. The subgroup of patients that we want to draw attention to is not the one with clearly detectable organic psychiatric syndromes in the sense of the frontal lobe syndromes (to whom Dr Vogel alludes and in whom ADHD represents only a fraction of a wider psychopathology). We want to draw attention to the fact that there are many patient subgroups in whom the psychopathology is identical to primary, mainly genetically related ADHD. Additional findings such as irregularities on the EEG, complications during birth, a medical history of encephalitis, or clinically remitting pyknopsy may indicate a secondary and not a primary, ie genetic, pathogenesis, although the clinical picture does not exceed that of classic ADHD. The categorization into primary and secondary ADHD can help explain why no positive family history exists in a large subgroup of patients. Further, this categorization can be clinically relevant-for example, when considering whether substances such as carbamazepine (for which several double blinded studies have been done with respect to their use in ADHD) or whether valproate should be tried out in the context of off label medication.

Differential diagnosis estrogen deficiency syndrome

To make a diagnosis of ADHD in adulthood, corresponding symptoms in childhood are essential. It is not an episodic disorder or one that manifests only in

adulthood. These criteria should be considered when differentiating ADHD from estrogen deficiency syndrome—whether relevant as a differential diagnosis or as a comorbidity (for example during the menopausal years).

DOI: 10.3238/arztebl.2008.0766

REFERENCES

1. Faraone SV, Spencer T, Aleardi M, Pagano C, Biederman J: Meta-analysis of the efficacy of methylphenidate for treating adult attention-deficit/hyperactivity disorder. *J Clin Psychopharmacol* 2004; 24: 24 –9.

Dr. med. Alexandra Philipsen

Abteilung für Psychiatrie und Psychotherapie
Universitätsklinikum Freiburg
Hauptstr. 5
79104 Freiburg, Germany
alexandra.philipsen@uniklinik-freiburg.de

Conflict of interest statement

Dr Philipsen has received honoraria for speaking and advisory tasks as well as travel expenses from Medice, Janssen-Cilag, Novartis, Lilly. She has also received research support from Medice and Janssen-Cilag.