Published in final edited form as:

J Am Chem Soc. 2008 May 28; 130(21): 6712–6713. doi:10.1021/ja801981h.

Mössbauer Evidence for an Exchange-Coupled $\{[\text{Fe}_4\text{S}_4]^{1+}:\text{Ni}_p^{1+}\}$ A-cluster in Isolated Alpha Subunits of Acetyl-Coenzyme A Synthase/Carbon Monoxide Dehydrogenase

Xiangshi Tan¹, Marlène Martinho², Audria Stubna², Paul A. Lindahl^{1,3,*}, and Eckard Münck^{2,*}

¹ Department of Chemistry, Texas A&M University, College Station, Texas 77843, USA

² Department of Chemistry, Carnegie Mellon University, Pittsburgh, PA 15213, USA

³ Department of Biochemistry and Biophysics, Texas A&M University, College Station, Texas 77843

Abstract

The active site A-cluster in the α subunit of the title enzyme consists of an Fe_4S_4 cluster coordinated to a $[\text{Ni}_p \text{Ni}_d]$ subcomponent. The cluster must be activated for catalysis using low-potential reductants such as Ti(III) citrate. Relative to the inactive $\{[\text{Fe}_4\text{S}_4]^{2+} \text{Ni}_p^{2+} \text{Ni}_d^{2+}\}$ state, the activated state appears to be 2-electrons more reduced, but the location of these electrons within the A-cluster is uncertain, with $\{[\text{Fe}_4\text{S}_4]^{2+} \text{Ni}_p^0 \text{Ni}_d^{2+}\}$ and $\{[\text{Fe}_4\text{S}_4]^{1+} \text{Ni}_p^{1+} \text{Ni}_d^{2+}\}$ configurations proposed. Recombinant apo- α subunits oligomerize after activation with NiCl_2 . The dimer fraction, upon reduction with excess Ti(III)citrate, exhibited Mössbauer spectra consisting of 2 quadrupole doublets representing 51% and 21% of the Fe, with parameters indicating $[\text{Fe}_4\text{S}_4]^{1+}$ states. Spectra recorded in strong magnetic fields were typical of diamagnetic systems, indicating an exchange-coupled $S = 0$ $\{[\text{Fe}_4\text{S}_4]^{1+} \text{Ni}_p^{1+}\}$ state. Additional treatment with CO altered the doublet Mössbauer parameters, suggesting an interaction with CO, but maintaining the cluster in the $\{[\text{Fe}_4\text{S}_4]^{1+} \text{Ni}_p^{1+}\}$ state. Reduction with sub-stoichiometric equivalents of Ti(III) citrate afforded an EPR signal typical of Ni^{1+} ions, with $g_{\parallel} = 2.10$ and $g_{\perp} = 2.02$. Addition of more Ti caused the signal intensity to decline, suggesting that it arises from the semi-reduced $\{[\text{Fe}_4\text{S}_4]^{2+} \text{Ni}_p^{1+}\}$ state.

Acetyl-coenzyme A synthase/carbon monoxide dehydrogenase is an $\alpha_2\beta_2$ tetramer, found in anaerobic archaea and bacteria that grow chemoautotrophically on CO or CO_2 .¹ The active site A-cluster in the synthase α subunit consists of an Fe_4S_4 cluster coordinated through a bridging cysteinate to a $[\text{Ni}_p \text{Ni}_d]$ subcomponent (Chart 1).² The oxidized state of the cluster assembly, A_{ox} , can be formulated as $\{[\text{Fe}_4\text{S}_4]^{2+} \text{Ni}_p^{2+} \text{Ni}_d^{2+}\}$.^{3–5} A_{ox} does not accept, in the presumed first catalytic step, a methyl cation from the corrinoid-iron-sulfur protein ($\text{CH}_3\text{-Co}^{3+}\text{FeSP}$).⁶ However, methyl transfer occurs when A_{ox} is treated with low-potential reductants such as Ti(III) citrate. The electronic configuration for the reductively activated $\text{A}_{\text{red-act}}$ has not been established but the state is probably 2 electrons more reduced than A_{ox} .⁴ Reduction of the square-planar distal Ni_d^{2+} with amide N coordination is unlikely,⁷ and stopped-flow evidence disfavors the reduction of the $[\text{Fe}_4\text{S}_4]^{2+}$ cluster,⁶ leaving the proximal Ni_p^{2+} as the only site that might accept electrons. Attainment of a Ni_p^0 state has been suggested,⁵ and this idea is supported by a report that a Ni^0 phosphine complex can accept a methyl cation from a $\text{CH}_3\text{-Co}^{3+}$ complex.⁸ However, independent DFT studies by Brunold and Field argue against this suggestion and support a $\{[\text{Fe}_4\text{S}_4]^{1+} \text{Ni}_p^{1+}\}$ configuration for $\text{A}_{\text{red-act}}$.^{9, 10} This

hypothetical state was not observed in our previous Mössbauer study where it was found that Ti(III) citrate-reduced Ni-activated recombinant α subunits exhibited two major species: ~ 70 % of spectral intensity originated from $\{[\text{Fe}_4\text{S}_4]^{1+} \text{Ni}_p^{2+}\}$ centers while ~ 30 % reflected a diamagnetic state containing an $[\text{Fe}_4\text{S}_4]^{2+}$ moiety.⁴ Given that the enzyme is known to be heterogeneous, with ~ 30 % functional and ~ 70 % nonfunctional A-centers,¹¹ this result was consistent with a $\{[\text{Fe}_4\text{S}_4]^{2+} \text{Ni}_p^0\}$ state in functional subunits.

We recently altered our procedure for preparing recombinant α subunit by adding NiCl_2 to purified apo- α subunit (containing the Fe_4S_4 cluster but lacking the Ni subcomponent) rather than including it in the growth medium.¹¹ Gel filtration FPLC of Ni-activated α subunits indicated Ni-dependent oligomerization into dimers and other forms.¹² Here we report Mössbauer evidence that α dimer samples prepared in this manner, when reduced by Ti(III) citrate, contain A-clusters in the $\{[\text{Fe}_4\text{S}_4]^{1+} \text{Ni}_p^{1+}\}$ state. We also report a new EPR signal that appears to arise from the semi-reduced $\{[\text{Fe}_4\text{S}_4]^{2+} \text{Ni}_p^{1+}\}$ state.

The 4.2 K Mössbauer spectrum of Ni-activated α dimers reduced with Ti(III) citrate (Figure 1A) exhibits three components. A paramagnetic component (≈ 20 -30 % of Fe), extending from -2 mm/s to $+2$ mm/s Doppler velocity, most probably belongs to $\{[\text{Fe}_4\text{S}_4]^{1+} \text{Ni}_p^{2+}\}$ clusters. The central portion of the spectrum exhibits two doublets (**1** and **2**) with $\Delta E_{Q1} = 1.21(4)$ mm/s, $\delta_1 = 0.56(2)$ mm/s, 51(3) % of Fe, and $\Delta E_{Q2} = 0.47(2)$ mm/s, $\delta_2 = 0.55(2)$ mm/s, 21(2) %. The relative intensities of **1** and **2** are $\approx 2.4:1$ rather than 2:2 or 3:1 as one might expect if doublets **1** and **2** would represent subsites in one $[\text{Fe}_4\text{S}_4]$ cluster. The $[\text{Fe}_4\text{S}_4]^{2+}$ cubane of the A-cluster has $\delta = 0.45(2)$ mm/s in all states examined,³ while $[\text{Fe}_4\text{S}_4]^{1+}$ clusters typically have an average $\delta \sim 0.54$ -0.59 mm/s.^{13, 14} The $S_c = \frac{1}{2}$ $[\text{Fe}_4\text{S}_4]^{1+}$ cluster of apo- α has $\delta = 0.55$ mm/s and $\Delta E_Q = 0.92$ mm/s (Figure S1). Thus, the values for δ_1 and δ_2 indicate the $[\text{Fe}_4\text{S}_4]^{1+}$ state.

Magnetically isolated $S_c = \frac{1}{2}$ $[\text{Fe}_4\text{S}_4]^{1+}$ clusters generally exhibit paramagnetic hyperfine structure at 4.2 K (e.g. Figure S1), reflecting hyperfine fields up to ≈ 12 T. The observation of doublets **1** and **2** at 4.2 K suggests A-clusters with integer or zero electronic cluster spin. This argument is supported by spectra recorded in applied fields of 0.1 T, 8 T and 7.0 T (Figure 1D), which show that **1** and **2** belong to diamagnetic A-clusters, strongly suggesting exchange coupling between an $S_c = \frac{1}{2}$ $[\text{Fe}_4\text{S}_4]^{1+}$ cluster and an $S_{\text{Ni}} = \frac{1}{2}$ Ni_p^{1+} site.

Figure 1C shows a 4.2 K spectrum of Ti (III) citrate-reduced α dimers after incubation with CO. This treatment produces the $S = \frac{1}{2}$ NiFeC signal, the signature of the exchange coupled, $\{[\text{Fe}_4\text{S}_4]^{2+} \text{Ni}_p^{1+}\text{-CO}\}$, $A_{\text{red-CO}}$ state.^{3, 15} In one preparation, we obtained by EPR 0.39(4) spins/ α together with the corresponding Mössbauer component, representing 34(3) % of the Fe. For a second preparation (Figure 1C) 31(2) % of the A-clusters were in the $A_{\text{red-CO}}$ state (dashed line drawn above Figure 1C). The remainder of the sample exhibits two doublets, **3** and **4**, again reflecting $[\text{Fe}_4\text{S}_4]^{1+}$ clusters albeit with parameters that differ from the Ti(III) citrate-reduced sample: $\Delta E_{Q3} = 0.96$ mm/s, $\delta_3 = 0.51$ mm/s, 42(3) % of Fe, and $\Delta E_{Q4} = 1.88$ mm/s, $\delta_4 = 0.60$ mm/s, 21(2) %. Interestingly, addition of CO changed all spectral components. We speculate that **3** and **4** reflect a $\{[\text{Fe}_4\text{S}_4]^{1+} \text{Ni}_p^{1+}\text{-CO}\}$ cluster.¹⁶ The data of Figure 1C also imply that all α subunits in the sample have the proximal site occupied by Ni. Studies at 8.0 T applied field (Figure S3) show that **3** (and probably **4**) originates from a diamagnetic site.

The fractions quoted for **1,2** and **3,4** do not reveal whether doublets **1,2** (and **3,4**) belong to the same cluster; we defer this problem to future studies. Here we discuss aspects of the electronic structure of the A-cluster implied by the observation of diamagnetic species with $\delta \approx 0.55$ -0.60 mm/s, using **1** as an example.

$[\text{Fe}_4\text{S}_4]^{1+}$ clusters occur in various spin states, mostly with $S_c = \frac{1}{2}$ and $3/2$. We ignore $S_c = 3/2$ configurations here as they do not yield a diamagnetic ground state for $\{[\text{Fe}_4\text{S}_4]^{1+} \text{Ni}_p^{1+}\}$.

Frequently, the $S_c = 1/2$ ground state of $[\text{Fe}_4\text{S}_4]^{1+}$ clusters yields a diferrous and mixed-valence pair, distinguishable by their ΔE_Q and δ values.¹³ However, some $[\text{Fe}_4\text{S}_4]^{1+}$ clusters,¹⁷ like that of apo- α of Figure S1, have the same ΔE_Q and δ for all sites. The exchange-coupling pathway between the $S_c = 1/2$ $[\text{Fe}_4\text{S}_4]^{1+}$ cluster and the $S_{\text{Ni}} = 1/2$ Ni_p^{1+} involves the cysteinate that links one cluster site, previously labeled Fe_D , to the Ni^{1+} ($\mathcal{A} = \text{JS}_D \cdot S_{\text{Ni}}$, where S_D is the local spin of cluster site D).³ The J-term acts in first-order between the two $S = 1/2$ ground manifolds and mixes in second-order excited $[\text{Fe}_4\text{S}_4]^{1+}$ cluster states into the ground manifold. The second-order term is relevant for understanding the $A_{\text{red-CO}}$ state³ but we ignore it here. Coupling then yields a singlet (A-cluster spin $S = 0$) and a triplet state, split by $K_D J$, where K_D is a spin projection factor ($|K_D| \leq 1.5$ ¹⁸) that depends on the internal coupling of the $[\text{Fe}_4\text{S}_4]^{1+}$ cluster and the nature of Fe_D . For $K_D J > 0$ the ground state is diamagnetic. The solid line in Figure 1D is a spectral simulation assuming that **1** and **2** are diamagnetic.

In a series of preliminary experiments we observed a new EPR signal when samples of Ni-activated α were treated with sub-stoichiometric amounts of Ti(III) citrate. This axial signal (Figure 2A) has $g_{\parallel} = 2.10$ and $g_{\perp} = 2.02$, values typical of Ni^{1+} species. The intensity of the axial signal increases as Ti(III) citrate is initially added (Figure 2B) and maximizes at ~ 0.2 equivalents/ α of Ti(III) citrate (See also Supplementary Information). The qualitative behavior of the titration plot suggests that the signal originates from a semi-reduced $S = 1/2$ $\{[\text{Fe}_4\text{S}_4]^{2+} \text{Ni}_p^{1+}\}$ state which is then further reduced to the $\{[\text{Fe}_4\text{S}_4]^{1+} \text{Ni}_p^{1+}\}$ state.

After the Mössbauer spectra were collected, the two samples of Figure 1 were analyzed for their ability to accept a methyl group from CoFeSP. In each case, 0.3 methyl groups per α were transferred, indicating that only $\sim 30\%$ of the α subunits in these samples arise from the functional $A_{\text{red-act}}$ state.

In summary, our results demonstrate that Ti(III) citrate can reduce the isolated Ni-activated α subunit of the title enzyme to form the $\{[\text{Fe}_4\text{S}_4]^{1+} \text{Ni}_p^{1+}\}$ state, and what appears to be the semi-reduced $\{[\text{Fe}_4\text{S}_4]^{2+} \text{Ni}_p^{1+}\}$ state. The former state has been suggested by DFT calculations, and the latter has been proposed as a catalytic intermediate¹ but here we provide the first experimental evidence for the existence of these states. Although further studies are required to reveal which of the Mössbauer spectral components exhibited by the Ti(III) citrate-treated sample represent the $A_{\text{red-act}}$ state, binding of a methyl cation to $\{[\text{Fe}_4\text{S}_4]^{1+} \text{Ni}_p^{1+}\}$ could result, after internal electron transfer from the $[\text{Fe}_4\text{S}_4]^{1+}$ moiety to Ni_p , in the methyl-bound state $\{[\text{Fe}_4\text{S}_4]^{2+} \text{Ni}_p^{2+}\text{-CH}_3\}$.⁴

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgment

This work was supported by National Institutes of Health grant GM46441 (PAL) and the National Science Foundation grant MCB-042 4494 (EM).

Supporting Information: Sample preparations. Mössbauer and EPR spectra of apo - α . Details of Mössbauer analyses and comments on exchange coupling. This material is available free of charge via the Internet at <http://pubs.acs.org>.

References

1. Ragsdale SW. J. Inorg. Biochem 2007;101:1657–1666. [PubMed: 17716738]
2. Darnault C, Volbeda A, Kim EJ, Legrand P, Vernede X, Lindahl PA, Fontecilla-Camps JC. Nat. Struct. Biol 2003;10:271–279. [PubMed: 12627225]
3. Xia J, Hu Z, Popescu CV, Lindahl PA, Münck E. J. Am. Chem. Soc 1997;119:8301–8312.

4. Bramlett MR, Stubna A, Tan X, Surovtsev IV, Münck E, Lindahl PA. *Biochemistry* 2006;45:8674–8685. [PubMed: 16834342]
5. Lindahl PA. *J. Biol. Inorg. Chem* 2004;9:516–524. [PubMed: 15221478]
6. Tan X, Sewell C, Yang Q, Lindahl PA. *J. Am. Chem. Soc* 2003;125:318–319. [PubMed: 12517128]
7. Kruger HJ, Peng G, Holm RH. *Inorg. Chem* 1991;30:734–742.
8. Eckert NA, Dougherty WG, Yap GPA, Riordan CG. *J. Am. Chem. Soc* 2007;129:9286–9287. [PubMed: 17622143]
9. Schenker RP, Brunold TC. *J. Am. Chem. Soc* 2003;125:13962–13963. [PubMed: 14611224]
10. Amara P, Volbeda A, Fontecilla-Camps JC, Field MJ. *J. Am. Chem. Soc* 2005;127:2776–2784. [PubMed: 15725036]
11. Loke H-K, Tan X, Lindahl PA. *J. Am. Chem. Soc* 2002;124:8667–8672. [PubMed: 12121109]
12. Tan X, Kagiampakis I, Surovtsev IV, Demeler B, Lindahl PA. *Biochemistry* 2007;46:11606–11613. [PubMed: 17887777]
13. Lindahl PA, Day EP, Kent TA, Orme-Johnson WH, Münck E. *J. Biol. Chem* 1985;260:11160–11173. [PubMed: 2993304]
14. Bertini I, Ciurli S, Luchinat C. *Struct. Bond* 1995;83:1–53.
15. Ragsdale SW, Wood HG, Antholine WE. *PNAS* 1985;82:6811–6814. [PubMed: 2995986]
16. The effect of CO is rather complex: for our earlier preparations we observed that adding CO caused oxidation of $[\text{Fe}_4\text{S}_4]^{1+}$ clusters to the 2+ state, see ref. 3 and 4.
17. Ragsdale SW, Lindahl PA, Münck E. *J. Biol. Chem* 1987;262:14289–14297. [PubMed: 2821001]
18. Noodleman L. *Inorg. Chem* 1991;30:246–256.

Chart 1.
Structure of the A-cluster. SR represents cysteinyl residues.

Figure 1.

4.2 K Mössbauer spectra of α dimer taken in applied fields as indicated. (A) Spectrum of 1 mM Ti(III) citrate-reduced α dimer. Red solid line is a superposition of doublets **1** and **2**, using the parameters quoted in the text; blue and green solid lines show doublets separately. (B) Spectrum of (A) after removing the line width contribution of the ^{57}Co source by a Fourier deconvolution. (C) Spectrum of 1 mM Ti(III) citrate-reduced α dimer after exposure to CO. The solid line is a simulation for doublets **3** and **4** superimposed on a simulation for the $A_{\text{red-CO}}$ species (dashed line above data, using published parameters³). (D) 7.0 T spectrum (parallel applied field) of the sample of (A) after removal of a 25 % contribution of apo- α . The solid

line is a spectral simulation assuming that **1** and **2** are diamagnetic; raw data and all parameters are given in Supplementary Information.

Figure 2.

(A) EPR spectrum of Ni-activated α dimer reduced with 0.2 Ti(III) citrate/ α monomer.

Conditions: 9.482 GHz; 1 mT modulation; T = 10 K. (B) Plot of EPR signal intensity (amplitude of derivative feature) versus amount of Ti(III) citrate added.