Published in final edited form as:

J Med Chem. 2008 October 9; 51(19): 6220–6224. doi:10.1021/jm8006195.

Anti-tumor Compounds Based on a Natural Product Consensus Pharmacophore

Chandraiah Lagiseti¹, Alan Pourpak², Qin Jiang², Xiaoli Cui², Tinopiwa Goronga¹, Stephan W. Morris^{2,3}, and Thomas R. Webb^{1,*}

¹Department of Chemical Biology and Therapeutics, St. Jude Children's Research Hospital, MS 1000, 332 North Lauderdale, Memphis, TN 38105.

²Department of Pathology, St. Jude Children's Research Hospital, MS 1000, 332 North Lauderdale, Memphis, TN 38105.

³Department of Oncology, St. Jude Children's Research Hospital, MS 1000, 332 North Lauderdale, Memphis, TN 38105.

Abstract

We report the design and highly enantioselective synthesis of a potent analog of the spliceosome inhibitor FR901464, based on a non-natural product scaffold. The design of this compound was facilitated by a pharmacophore hypothesis that assumed key interaction types that are common to FR901464 and an otherwise unrelated natural product (pladienolide). The synthesis allows for the preparation of numerous novel analogs. We present results on the *in vitro* activity for this compound against several tumor cell lines.

A novel mechanism of anti-tumor activity was recently elucidated for two structurally distinct fermentation products (see Figure 1)¹ and pladienolide² (actually a set of closely related compounds that have been designated pladienolides A-G, isolated from *Streptomyces platensis*)³ and FR901464 (isolated from *Pseudomonas* sp. No 2663).^{4–6} These natural products have previously been reported to have a similar novel effect on the cell cycle in mammalian cell lines; cell cycle arrest at the G1 and G2/M phases.^{3, 4} This effect on the cell cycle distinguished these compounds from other anti-tumor agents that also inhibit proliferation in tumors and act at previously identified molecular targets.^{3–5} This interaction with the spliceosome leads to the inhibition of mRNA splicing and the translation of unspliced mRNA, which ultimately results in the observed effects on the cell cycle.^{2, 7} The first of these natural products to be discovered, FR901464, showed activity in *in vivo* cancer animal models, but showed a relatively narrow therapeutic window.⁴ On the other hand pladienolide-based compounds (first reported in 2004) have shown a very broad therapeutic window.² Of the seven pladienolides, pladienolide B was reported to have the most promising anti-cancer activity *in vivo*, and it has recently been reported that a derivative of pladienolide B has entered human clinical trials for cancer.²

Motivated by the recognition that FR901464 and pladienolide have a very similar (or possibly identical) mechanism of action, we compared overlays of molecular models of the two natural products. Upon inspection of these overlays we recognized that despite the gross dissimilarities

*To whom correspondence should be addressed. Phone 901-495-3928. Fax: 901-495-5715. E-mail: thomas.webb@stjude.org.

Supporting Information Available: The supporting information includes the general experimental methods, procedures for the synthesis of **9–13** and **15–18**, as well as full experimental methods for the cell cycle analysis and the *in vitro* cytotoxicity assays. This material is available free of charge via the Internet at <http://pubs.acs.org>.

between FR901464 and pladienolide, both molecules share certain common key features (which are known to be critical to their activity) that can overlap in low energy conformations in three dimensions (see Figure 2). It is known from previous studies that the orientation of the acetyl group and the presence of the epoxide are important in conferring activity to FR901464 analogs,^{8,9} but that the glycosidic hydroxyl group can be replaced by an alkoxy,¹⁰ or a methyl group,¹¹ to give more stable and more active compounds. Active analogs of pladienolide also contain similar functionality,^{2,3} though less structure-activity information has been reported on these more recently discovered compounds. This led us to propose that the three critical requirements for activity of both FR901464 and pladienolide are an appropriately positioned epoxy and carbonyloxy group connected via an appropriately constrained 'linker' group of the right shape and length. We also assumed that these groups would need to be conformationally constrained in a way similar to that seen in the natural products. Information regarding the requirement of the linker group is not available, but both natural products have significant similarities in the structure of their diene linker group. In FR901464, the linking group includes conformational constraints consisting of two pyran 6-membered rings, whereas in pladienolide the common diene constraint is an unsaturated 12-membered macrocyclic lactone. Using this information we developed a hypothetical 3D pharmacophore model that allowed us to propose that compounds of the general structure **1–3** (see Figure 3) could have similar biological activity, and could therefore serve as starting points for the development of compounds with potential for human cancer therapy. The proposed compound **1** contains the unchanged methylpenta-2,4-diene group and 5-oxopent-3-en-2-yl acetate group along with the modified deoxyspiro-epoxide group, with the pyran ring replaced by a cis-1,4-substituted cyclohexyl group, whereas compounds **2** and **3** include additional polar modifications to the methylpenta-2,4-diene group that are very synthetically assessable. These changes (particularly for compound **1**) are expected to impart similar conformational presentation of the linker, the epoxy and the carbonyloxy groups as that found in FR901464.

It should be noted that these modifications (conversion of the pyran ring oxygen to carbon, removal of two methyl groups and removal of a hydroxyl oxygen) eliminate a total of 6 chiral centers when compared to FR901464, which made the synthesis of all of our target compounds readily achievable. At the outset of our project it was not clear that compounds containing all of these substantial modifications would be active, since there have been no reports providing detailed structural information regarding the interaction of FR901464 and pladienolide with the spliceosome SF3b subunit or of the synthesis of compounds that are similar to **1–3**. We believed, however, that there was a reasonable likelihood that one or more of our compounds could demonstrate substantial activity, especially given that there is an example of a FR901464 analog with a cell-based growth inhibition IC₅₀ as low as 10 picomolar.¹¹

There have been numerous synthetic approaches to FR901464, however since these synthetic approaches target the much more complex natural product these routes were not highly instructive for the construction of our synthetic targets **1–3**. Our approach to these analogs is shown in Scheme 1 and Scheme 2. We found the readily available inexpensive compound **4** (which has commercial application as an insect repellent) to be an attractive starting material for the left hand portion of our synthetic targets **1–3**. In order to introduce the desired stereochemistry we explored numerous transition metal catalysts for the enantioselective reduction of **4** to ketoester **5**; for example we investigated the ruthenium(II) BINAP complex as a chiral hydrogenation catalyst, using the published conditions, but found either no reduction, or at higher temperatures the undesired reduction of the ketone carbonyl in preference to the carbon-carbon double bond.¹² However, we were pleased to find that application of the appropriate MacMillan conditions¹³ gave the desired product in good yield (though contaminated with the starting material **4**, which co-eluted on silica chromatography) and in high enantiomeric excess (91% based on chiral SFC chromatography of a derivative, see Supporting Information). The absolute stereochemistry of our ketoester product **5** is based on

numerous published examples of the stereochemistry obtained using this organocatalyst on similar substrates.^{13, 14} We then turned our attention to the diastereoselective introduction of the desired spiroepoxide; we expected that the addition of dimethylsulfoxonium methylide¹⁵ would occur from an axial approach on the ketone carbonyl of **5**, to give primarily the desired diastereomer (epoxyester **6**). We were very pleased that this was indeed the case; in fact we did not detect any of the undesired epoxide in the crude reaction mixture, and the product was obtained in excellent yield (see Supporting Information for details on the NOE experiments used for assigning the relative stereochemistry). This approach allowed us to set two of the three chiral centers in our targets, and provided intermediates that were appropriately functionalized for conversion to our ultimate targets **1–3**.

The synthesis of the compound **1** could then be accomplished starting with the aldehyde **9** derived from the commercially available *cis*-cyclohexane derivative **8**, followed by homologation to give *trans* alcohol **10** cleanly in high overall yield.¹⁶ This compound was converted to the corresponding 2-thiobenzthiazole derivative **11** using Mitsunobu conditions¹⁷ and subsequently oxidized to the corresponding 2-sulfonylbenzothiazole derivative **12**.¹⁰ Removal of the BOC group in **12** and coupling of the optically pure (*S,Z*)-4-(dimethyl-*t*-butylsilyloxy)pent-2-enoic acid (prepared according to the known procedure)¹⁸ gave the desired functionalized amide **13** in excellent overall yield. Coupling of this reagent with epoxy aldehyde **7a**,¹⁰ gave diene **14** followed by deprotection and acylation of **15** gave final compound **1** cleanly in acceptable yield (the overall yield of **1** from **4** is 2.7%).

In order to give the carbamate **2** and carbonate **3**, as shown in Scheme 2, we used the commercially available alcohol **16** to prepare the azide **17**, which could be deprotected and converted to the amide azide **18** in excellent yield. This derivative could be reduced with triphenylphosphine on solid support to cleanly give amine **19**,¹⁹ which was then converted to the carbamate **2** in excellent yield²⁰ (the overall yield of **2** from **16** is 12.6%). In a similar way the alcohol **16** was converted to the *p*-nitrophenylcarbonate intermediate **22**, which was deprotected and amidation with (*S,Z*)-4-(dimethyl-*t*-butylsilyloxy)pent-2-enoic acid (prepared according to the published method)¹⁰ to give amide **23** in good yield. This amide could then be coupled with alcohol **7b** (from Scheme 1) to give the carbonate **3** (the overall yield of **3** from **16** is 14.2%).

A panel of seven human cancer cell lines was then used to evaluate the cytotoxic potential of compounds **1–3**, however only compound **1** showed potent activity. Compound **2** did not show any significant cytotoxic activity in our initial tumor line screen (in A549, COS-7, MCF-7, WiDr and NIH3T3 cells) at concentrations up to 5 μ M. In this screen, compound **3** inhibited cell growth by only 10–20 % at the highest concentration tested of 5 μ M, but only in the A549 and MCF-7 lines. Compound **1** was then screened in an expanded tumor panel consisting of the following cell lines: A549 lung cancer, WiDr colorectal cancer, MCF-7 breast cancer, PC-3 prostate cancer, OVCAR-3 ovarian cancer, and JeKo-1 and JVM-2 mantle cell lymphoma cell lines. An XTT colorimetric cytotoxicity assay was employed to estimate the IC₅₀ values (see Table 1). The cytotoxicity profile of compound **1** against the A549, MCF-7, OVCAR-3, PC-3 and WiDr cancer cell lines revealed IC₅₀ values of 4.65, 2.29, 1.98, 2.12, and 2.03 μ M, respectively (Table 1). Although compound **1** is substantially less potent than the natural products against several of the cell lines tested, it shows potent and selective killing of two tumor cell lines (JeKo-1 and JVM-2, 100 nM and 130 nM IC₅₀s, respectively).

To further investigate the biological activity of compound **1**, the effect of the compound on the cell cycle was evaluated. Following an 18-hour incubation of compound **1** with the A549 lung cancer, JeKo-1 mantle cell lymphoma, and WiDr colorectal cancer cell lines, all three lines accumulated in the G₂/M phases of the cell cycle (Figure 4). The degree of G₂/M cell cycle arrest correlated with the sensitivity of the cell lines to the cytotoxic activity of compound **1**

(JeKo-1 > WiDr > A549). In addition to the increased fraction of cells in the G₂/M phase of the cell cycle, a concomitant decrease in the S-phase fraction was observed for all three cell lines following treatment with compound **1**. As reported in the literature,^{3,4,7} the spliceosome inhibitor compounds pladienolide B, FR901464 and spliceostatin A all also induce a similar accumulation of cells in the G₂/M phases of the cell cycle, with a simultaneous decrease in the fraction of cells in S phase.

In conclusion, we report the enantioselective and diastereospecific synthesis of compound **1** that shows an activity profile similar to FR901464 and pladienolide, but with reduced potency. The design of this compound is based on insights from the common positioning of similar interacting groups from these unrelated natural products. Application of the judicious modification of atom types and the removal of groups that did not appear to impart interaction features or conformational constraints we were able to develop compounds that showed significant anti-tumor activity, but that have only three stereocenters (as compared to nine in FR901464) and are therefore much more synthetically attractive. Since compound **1** displays a similar bioactivity profile it is quite possible that it also acts in a manner analogous to FR901464 and pladienolide, by targeting the SF3b subunit of the spliceosome. The synthetic scheme that we have developed allows for the introduction of numerous modifications that will allow us to fully elucidate the structure-activity relationships in this series, and will also allow us to prepare compounds that have an improved *in vivo* profile, in order to develop lead compounds as potential human therapeutics.

Experimental Section

(S)-Butyl 6, 6-dimethyl-4-oxotetrahydro-2H-pyran-2-carboxylate (**5**)

A solution of commercially available butopyronoxyl **4** (230 mg, 1.01 mmol) and (2R,5R)-5-butyl-methyl-2-(5-methylfuran-2-yl-) imidazolidin-4-one (55 mg, 0.20 mmol) in anhyd Et₂O (4.6 mL) was stirred and cooled at ice bath temp (0–4 °C) as di-*tert*-butyl 2,6-dimethyl-dicarboxylate (346 mg, 1.11 mmol) and trichloroacetic acid (TCA) (32 mg, 0.20 mmol) were sequentially added. The resulted yellow solution was allowed to stir for 3 days at 4 °C. The reaction mixture was passed through a short pad of silica gel and eluted with Et₂O (60 mL). Concentration of the Et₂O and purification of the residue by flash chromatography (20% EtOAc in hexane) gave 204 mg (88%) of keto ester **5** as an oil. [α]_D²⁵ +15.7°, (*c* = 2.1 in CHCl₃); ¹H NMR (400 MHz, CDCl₃) δ 4.46 (dd, *J* = 4.0, 11.0 Hz, 1H), 4.19 (td, *J* = 2.7, 6.7 Hz, 2H), 2.64–2.47 (m, 3H), 2.32 (dd, *J* = 1.6, 14.0 Hz, 1H), 1.68–1.59 (m, 2H), 1.46 (s, 3H), 1.43–1.33 (m, 2H), 1.23 (s, 3H), 0.94 (t, *J* = 7.4 Hz, 3H); ¹³C NMR (CDCl₃, 100 MHz) δ 205.2, 170.1, 75.2, 70.0, 65.4, 52.9, 43.5, 30.6, 30.5, 23.7, 19.0, 13.6; IR (Neat Film) 2965, 2875, 1734, 1725, 1464, 1379, 1241, 1181 cm^{–1}; HRMS (FAB) *m/z* Calcd for C₁₂H₂₁O₄ (M+1)⁺ 229.1440, found 229.1440.

(3R, 5S)-butyl 7, 7-dimethyl-1, 6-dioxaspiro [2.5] octane-5-carboxylate (**6**)

A suspension of sodium hydride (96 mg, 2.40 mmol, 60% in mineral oil) in anhyd DMSO (4.2 mL) was stirred at rt as trimethylsulfoxonium iodide (491 mg, 2.40 mmol) was added. After 30 min, the keto ester **5** (500 mg, 2.19 mmol) in DMSO (1.0 mL) was added drop-wise. After 1h at rt, the reaction was quenched with ice cold H₂O (5 mL). The product was then extracted with EtOAc (3×30 mL) and the combined extracts were washed with H₂O (2×15 mL) and brine (20 mL), and dried over MgSO₄. Evaporation of the solvent and purification by flash chromatography (20% EtOAc in hexane) gave 430 mg (81%) of epoxide derivative **6** as a liquid. [α]_D²⁵ +41.5° (*c* = 2.2 in CHCl₃); ¹H NMR (400 MHz, CDCl₃) δ 4.53 (dd, *J* = 2.7, 12.1 Hz, 1H), 4.14 (t, *J* = 6.7 Hz, 2H), 2.61 (dd, *J* = 4.5, 13.9 Hz, 2H), 2.14–2.02 (m, 2H), 1.65–1.58 (m, 2H), 1.47 (ddd, *J* = 4.2, 6.7, 10.9 Hz, 1H), 1.39 (s, 3H), 1.38–1.33 (m, 2H), 1.34 (s, 3H), 1.16 (dd, *J* = 2.0, 14.0 Hz, 1H), 0.93 (t, *J* = 7.3 Hz, 3H); ¹³C NMR (CDCl₃, 100 MHz)

δ 171.5, 74.0, 68.3, 64.9, 55.2, 51.1, 42.2, 35.2, 31.0, 30.5, 23.3, 19.0, 13.7; IR (Neat Film) 2966, 2874, 1756, 1465, 1378, 1218; 1180 cm^{-1} ; HRMS (FAB) m/z Calcd for $\text{C}_{13}\text{H}_{23}\text{O}_4$ ($M+1$)⁺ 243.1596, found 243.1595.

(3R,5S)-7,7-dimethyl-1,6-dioxaspiro[2.5]octane-5-carbaldehyde (7a)

To a stirred solution of ester **6** (1 g, 4.1 mmol) in anhyd hexane (15 mL) and THF (9 mL) was added DIBALH (4.7 mL, 4.75 mmol, 1.0 M in hexane) at -78°C . The resulting solution was allowed to stir for 30 min at -78°C by which time TLC indicated that the alcohol has been consumed. MeOH (1.5 mL) was added and diluted with saturated aqueous sodium/potassium tartarate (30 mL) and EtOAc (80 mL). The solution was stirred for 15 min at rt to get a clear solution. The organic layer was separated and the aqueous layer was extracted with EtOAc (2 $\times$ 100 mL). The combined organic layers were washed with brine (45 mL), and dried over Na_2SO_4 . Evaporation of the solvent and purification of the residue by flash chromatography (50% EtOAc in hexane) gave 390 mg (56%) of aldehyde as a liquid. ^1H NMR (400 MHz, CDCl_3) δ 9.65 (s, 1H), 4.35 (dd, $J = 3.0, 12.0$ Hz, 1H), 2.60 (q, $J = 4.6$ Hz, 2H), 2.04–1.86 (m, 2H), 1.46–1.40 (m, 1H), 1.39 (s, 3H), 1.33 (s, 3H), 1.20 (dd, $J = 2.0, 14.0$ Hz, 1H); ^{13}C NMR (100 MHz, CDCl_3) δ 201.7, 74.0, 73.6, 54.7, 51.1, 42.5, 32.6, 30.6, 30.9, 23.3; IR (neat film) ν 2970, 2913, 1732, 1184, 1084 cm^{-1} ; HRMS (FAB) m/z Calcd for $\text{C}_9\text{H}_{14}\text{O}_3$ (M^+) 170.0943, found 170.0943.

7,7-dimethyl-1,6-dioxaspiro[2.5]octan-5-yl) methanol (7b)

A solution of epoxy ester **6** (320 mg, 1.32 mmol) in anhyd THF (5.0 mL) was stirred and cooled at -78°C as DIBALH (4.6 mL, 1.0 M in hexane) was slowly added drop-wise. The solution was stirred at the same temperature for 3 h. Saturated aqueous NH_4Cl (4 mL) was introduced to destroy excess reagent at -78°C , this mixture was allowed to warm to rt and stir for 30 min to get a clear mixture. The organic layer was then separated and the aqueous layer was extracted with EtOAc (3 $\times$ 30 mL). The combined layers were washed with saturated aqueous NaHCO_3 (15 mL) and brine (15 mL), and dried over Na_2SO_4 . Concentration and purification by flash chromatography (50% EtOAc in hexane) gave the alcohol **7b** in quantitative yield as a liquid. ^1H NMR (400 MHz, CDCl_3) δ 4.07–3.98 (m, 1H), 3.70–3.61 (m, 1H), 3.53–3.43 (m, 1H), 2.58 (q, $J = 4.6$ Hz, 2H), 2.15 (t, $J = 5.8$ Hz, 1H), 1.99–1.88 (m, 2H), 1.38 (s, 3H), 1.25 (s, 3H), 1.17 (dd, $J = 2.0, 13.9$ Hz, 1H), 1.08–1.01 (m, 1H); ^{13}C NMR (100 MHz, CDCl_3) δ 73.1, 68.7, 65.9, 55.5, 51.0, 42.7, 33.7, 31.3, 23.8; IR (Neat Film) 3435, 2973, 2914, 1376, 1250, 1221, 1186, 1093, 1052 cm^{-1} ; HRMS (FAB) m/z Calcd for $\text{C}_9\text{H}_{16}\text{O}_3$ (M)⁺ 172.1100, found 172.1099.

(S,Z)-4-(tert-butyl dimethylsilyloxy)-N-((1R,4R)-4-((2E,4E)-5-((3R,5S)-7,7-dimethyl-1,6-dioxaspiro[2.5]octan-5-yl)-3-methylpenta-2,4-dienyl)cyclohexyl)pent-2-enamide (14)

A solution of sulfone derivative **13** (430 mg, 0.75 mmol) in anhyd THF (3 mL) was stirred and cooled at -78°C as LHMDS (0.92 mL, 0.92 mmol, 1.0 M in THF) was added drop-wise. The solution turned orange during the addition. After stirring for 10 min at -78°C , the aldehyde **7a** (75 mg, 0.75 mmol) in anhyd THF (2 mL) was introduced drop-wise. The resulting suspension was again stirred for 1 h at -78°C . The reaction was then slowly allowed to warm to rt and stirred for 2 h. Saturated aqueous NH_4Cl solution (15 mL) was added and the product was extracted with EtOAc (3 $\times$ 35 mL). The combined extracts were washed with brine (40 mL), and dried over MgSO_4 . Evaporation of the solvent and purification of the residue by flash chromatography (25% EtOAc in hexane) gave 70 mg of diene **14** as a liquid. ^1H NMR (400 MHz, CDCl_3) δ 6.27 (d, $J = 15.7$ Hz, 1H), 5.99 (dd, $J = 7.8, 11.6$ Hz, 1H), 5.61–5.45 (m, 5H), 4.48–4.44 (m, 1H), 4.03 (s, 1H), 2.57 (s, 2H), 2.08 (t, $J = 7.2$ Hz, 2H), 2.01–1.89 (m, 2H), 1.72 (s, 3H), 1.57–1.53 (m, 6H), 1.46–1.36 (m, 1H), 1.40 (s, 3H), 1.29–1.27 (m, 6H), 1.18–1.08 (m, 4H), 0.87 (s, 9H), -0.01 (d, $J = 5.3$ Hz, 6H); IR (Neat Film) 3394, 2926, 2856, 1735, 1666,

1629, 1532, 1448, 1320, 1244, 1049 cm^{-1} ; HRMS (ESI) m/z Calcd for $\text{C}_{31}\text{H}_{54}\text{NO}_4\text{Si}$ ($\text{M}+1$)⁺ 532.3822, found 532.3825.

(S,Z)-N-((1R,4R)-4-((2E,4E)-5-((3R,5S)-7,7-dimethyl-1,6-dioxaspiro[2.5]octan-5-yl)-3-methylpenta-2,4-dienyl)cyclohexyl)-4-hydroxypent-2-enamide (15)

A solution of TBS protected diene **14** (13 mg, 0.02 mmol) in THF (1.0 mL) was stirred and cooled at 0 °C as TBAF (49 μL , 0.04 mmol, 1.0 M in THF) was added. The resulting yellow solution was allowed to stir for 2h at rt by which time the TLC showed that all of starting material has been consumed. Evaporation of THF and purification of the residue by flash chromatography (60–80% EtOAc in hexane) gave 9.5 mg (95%) of **15** as a liquid. $[\alpha]_{\text{D}}^{25} +78^\circ$ ($c = 0.86$ in CHCl_3); ^1H NMR (400 MHz, CDCl_3) δ 6.27 (d, $J = 15.7$ Hz, 1H), 6.15 (dd, $J = 5.4, 11.9$ Hz, 1H), 5.87 (bd, 1H), 5.75 (dd, $J = 1.6, 12.0$ Hz, 1H), 5.56 (dd, $J = 6.7, 15.6$ Hz, 1H), 5.47 (t, $J = 7.7$ Hz, 1H), 4.80–4.73 (m, 1H), 4.50–4.43 (m, 1H), 4.09–4.02 (m, 1H), 2.57 (s, 2H), 2.09 (t, $J = 7.3$ Hz, 2H), 2.04–1.88 (m, 2H), 1.72 (s, 3H), 1.69–1.57 (m, 6H), 1.52–1.44 (m, 1H), 1.40 (s, 3H), 1.34 (d, $J = 6.7$ Hz, 3H), 1.28 (s, 3H), 1.26–1.23 (m, 2H), 1.18–1.11 (m, 2H); ^{13}C NMR (100 MHz, CDCl_3) δ 165.9, 150.0, 136.1, 134.1, 131.6, 127.1, 123.1, 73.0, 69.6, 64.6, 55.7, 51.0, 45.8, 42.4, 38.6, 36.4, 34.1, 31.5, 29.2, 27.8, 27.7, 23.8, 22.8, 12.5; IR (Neat Film) 3307, 2926, 2857, 1656, 1622, 1536, 1447, 1372, 1258, 1183, 1108 cm^{-1} ; HRMS (ESI) m/z Calcd for $\text{C}_{25}\text{H}_{40}\text{NO}_4$ ($\text{M}+1$)⁺ 418.2957, found 418.2948.

(S,Z)-5-((1R,4R)-4-((2E,4E)-5-((3R,5S)-7,7-dimethyl-1,6-dioxaspiro[2.5]octan-5-yl)-3-methylpenta-2,4-dienyl)cyclohexylamino)-5-oxopent-3-en-2-yl acetate (1)

A solution of alcohol **15** (7 mg, 0.01 mmol) in CH_2Cl_2 (0.9 mL) was allowed to stir at 0 °C as Et_3N (16 μL , 0.11 mmol) and DMAP (2.0 mg, 0.01 mmol) were sequentially added. After 5 min, acetic anhydride (5.4 μL , 0.05 mmol) was added using a micropipette. The solution was stirred for 1h at 0 °C. Evaporation of the solvent and purification of the residue by flash chromatography (50% EtOAc in hexane) gave 6.1 mg (79%) of **1** as a viscous oil. $[\alpha]_{\text{D}}^{25} -1.0^\circ$, ($c = 0.21$ in CHCl_3); ^1H NMR (400 MHz, CDCl_3) δ 7.04 (br d, 1H), 6.27 (d, $J = 15.7$ Hz, 1H), 5.81 (m, 2H), 5.66 (dd, $J = 9.2, 11.6$ Hz, 1H), 5.55 (dd, $J = 6.7, 15.7$ Hz, 1H), 5.49 (t, $J = 7.5$ Hz, 1H), 4.50–4.43 (m, 1H), 4.12–4.06 (m, 1H), 2.57 (s, 2H), 2.12–2.06 (m, 2H), 2.07 (s, 3H), 2.01–1.88 (m, 2H), 1.72 (s, 3H), 1.72–1.66 (m, 1H), 1.64–1.57 (m, 6H), 1.40 (s, 3H), 1.36 (d, $J = 6.4$ Hz, 3H), 1.28 (s, 3H), 1.27–1.19 (m, 3H), 1.15 (dd, $J = 2.0, 13.8$ Hz, 1H); ^{13}C NMR (100 MHz, CDCl_3) δ 171.1, 164.8, 137.9, 136.2, 133.8, 131.9, 126.9, 125.4, 73.0, 69.6, 69.2, 55.6, 51.0, 45.5, 42.4, 38.6, 36.4, 34.2, 31.5, 29.5, 29.4, 27.8, 27.6, 23.7, 21.2, 20.3, 12.5; 21 IR (Neat Film) 3310, 2928, 2856, 1660, 1626, 1534, 1445, 1365, 1252, 1219, 1185, 1116, 1072 cm^{-1} ; HRMS (ESI) m/z Calcd for $\text{C}_{27}\text{H}_{42}\text{NO}_5$ ($\text{M}+1$)⁺ 460.3063, found 460.3049.

7,7-dimethyl-1,6-dioxaspiro[2.5]octan-5-yl)methyl((1R,4s)-4-((S,Z)-4-(tert-butylidimethylsilyloxy)pent-2-enamido)cyclohexyl)methylcarbamate (20)

A solution of alcohol **7b** (40 mg, 0.23 mmol) was allowed to stir in anhyd THF (2.5 mL) at rt as Et_3N (0.23 mmol) and DMAP (2.8 mg, 0.02 mmol) were added. After 10 min, *p*-nitrophenyl chloroformate (45 mg, 0.465 mmol) was added and the resulting white suspension was stirred for overnight at rt. The reaction mixture was then diluted with EtOAc (40 mL) and washed with aqueous 1N HCl (10 mL) and saturated aqueous NaHCO_3 (3 $\times$ 10 mL), brine (10 mL), and dried over Na_2SO_4 . Evaporation of the solvent and purification of the residue by flash chromatography (15% EtOAc in hexane) gave (7,7-dimethyl-1,6-dioxaspiro[2.5]octan-5-yl)methyl 4-nitrophenyl carbonate (70 mg, 90%) as a viscous liquid. The amine derivative **19** (16 mg, 0.04 mmol) in anhyd THF (0.4 mL) and Et_3N (1 drop) was added to a stirred mixture of the above activated carbonate (23 mg, 0.07 mmol). After 12 h stirring at rt, the solvent was evaporated *in vacuo*. Purification of the residue by flash chromatography (30% EtOAc in hexane) gave 8 mg of **20** in quantitative yield as a liquid. ^1H NMR (400 MHz, CDCl_3) δ 6.00

(dd, $J = 7.8, 11.6$ Hz, 1H), 5.60-5.51 (m, 3H), 4.79 (t, $J = 5.8$ Hz, 1H), 4.15-4.00 (m, 4H), 3.11 (t, $J = 6.5$ Hz, 2H), 2.57 (q, $J = 4.6$ Hz, 2H), 1.97 (d, $J = 14.0$ Hz, 1H), 1.84 (dd, $J = 11.5, 13.4$ Hz, 1H), 1.68-1.60 (m, 6H), 1.36 (s, 3H), 1.30-1.21 (m, 9H), 1.18-1.11 (m, 2H), 0.87 (s, 9H), 0.04 (d, $J = 5.5$ Hz, 6H); ^{13}C NMR (100 MHz, CDCl_3) δ 164.8, 156.5, 150.5, 119.6, 73.2, 67.4, 67.1, 65.5, 53.3, 51.0, 45.5, 45.3, 42.4, 36.1, 34.3, 31.3, 28.9, 28.8, 25.9, 25.4, 23.7, 23.5, 18.1, -4.6, -4.7; IR (Neat Film) 3328, 2929, 2857, 1701, 1659, 1532, 1452, 1368, 1337, 1251, 1117, 1074 cm^{-1} ; HRMS (ESI) m/z Calcd for $\text{C}_{28}\text{H}_{51}\text{N}_2\text{O}_6\text{Si}$ ($\text{M}+1$) $^+$ 539.3516, found 539.3506.

(7,7-dimethyl-1,6-dioxaspiro[2.5]octan-5-yl)methyl ((1R,4s)-4-((S,Z)-4-hydroxypent-2-enamido)cyclohexyl)methyl carbamate (21)

Removal of the TBS group in **20** (7 mg, 0.01 mmol) used the same procedure as for the conversion of **14** to **15**, to give 5 mg (90%) of **21** as an oil. ^1H NMR (400 MHz, CDCl_3) δ 6.15 (dd, $J = 5.5, 11.9$ Hz, 1H), 5.93 (s, 1H), 5.74 (d, $J = 12.0$ Hz, 1H), 5.43 (s, 1H), 4.86-4.72 (m, 2H), 4.11-4.04 (m, 3H), 3.12 (t, $J = 6.5$ Hz, 2H), 2.57 (q, $J = 4.6$ Hz, 2H), 1.97 (d, $J = 14.0$ Hz, 1H), 1.84 (dd, $J = 11.6, 13.5$ Hz, 1H), 1.70-1.62 (m, 6H), 1.36 (s, 3H), 1.35 (d, $J = 6.7$ Hz, 3H), 1.29-1.24 (m, 6H), 1.18-1.11 (m, 2H); ^{13}C NMR (100 MHz, CDCl_3) δ 165.8, 156.6, 149.9, 123.2, 73.2, 67.5, 67.0, 64.6, 55.4, 50.9, 45.2, 42.5, 36.1, 34.3, 31.3, 29.7, 25.4, 22.8; IR (Neat Film) 3333, 2924, 2856, 1707, 1657, 1535, 1454, 1376, 1251, 1115, 1022 cm^{-1} ; HRMS (ESI) m/z Calcd for $\text{C}_{22}\text{H}_{37}\text{N}_2\text{O}_6$ ($\text{M}+1$) $^+$ 425.2652, found 425.2649.

(2S,Z)-5-(((1s,4R)-4-(((7,7-dimethyl-1,6-dioxaspiro[2.5]octan-5-yl)methoxy)carbonylamino)methyl)cyclohexylamino)-5-oxopent-3-en-2-yl acetate (2)

Acetylation of **21** (6 mg, 0.01 mmol) used the same procedure as for conversion of **15** to the corresponding acetate **1**, to give 5.9 mg (90%) of **2** as an oil. ^1H NMR (400 MHz, CDCl_3) δ 7.19 (br d, 1H), 5.82-5.74 (m, 2H), 5.65 (dd, $J = 9.1, 11.8$ Hz, 1H), 4.89 (br t, 1H), 4.17-4.04 (m, 4H), 3.16-3.08 (m, 2H), 2.57 (q, $J = 4.6$ Hz, 2H), 2.09 (s, 3H), 1.97 (d, $J = 13.9$ Hz, 1H), 1.84 (dd, $J = 11.6, 13.5$ Hz, 1H), 1.77-1.69 (m, 2H), 1.68-1.59 (m, 4H), 1.37-1.34 (m, 6H), 1.28-1.24 (m, 6H), 1.18-1.11 (m, 2H); ^{13}C NMR (100 MHz, CDCl_3) δ 164.9, 152.6, 137.3, 125.8, 73.1, 69.1, 67.1, 55.4, 50.9, 45.8, 45.2, 42.5, 36.2, 31.3, 29.7, 29.1, 25.4, 25.1, 23.6, 21.3; ^{21}IR (Neat Film) 3334, 2928, 2859, 1717, 1666, 1533, 1452, 1371, 1247, 1122, 1047 cm^{-1} ; HRMS (ESI) m/z Calcd for $\text{C}_{24}\text{H}_{39}\text{N}_2\text{O}_7$ ($\text{M}+1$) $^+$ 467.2757, found 467.2759.

((1R,4s)-4-((S,Z)-4-(tert-butyldimethylsilyloxy)pent-2-enamido)cyclohexyl)methyl (7,7-dimethyl-1,6-dioxaspiro[2.5]octan-5-yl)methyl carbonate (24)

A solution of carbonate derivative **23** (36 mg, 0.07 mmol) in CH_3CN (1 mL) was stirred at rt, followed by the addition of alcohol **7b** (20 mg, 0.10 mmol) in CH_3CN (1 mL) and DMAP (10 mg, 0.07 mmol). After 12h at rt, the solvent was evaporated under reduced pressure. Purification of the residue by flash chromatography (30% EtOAc in hexane) gave 20 mg (53%) of **24** as a liquid. ^1H NMR (400 MHz, CDCl_3) δ 6.00 (dd, $J = 7.8, 11.6$ Hz, 1H), 5.61 (d, $J = 7.5$ Hz, 1H), 5.53 (7-5.48, 2H), 4.20-4.06 (m, 3H), 4.03 (d, $J = 6.8$ Hz, 2H), 2.57 (q, $J = 4.6$ Hz, 2H), 2.04-1.80 (m, 3H), 1.70-1.62 (m, 6H), 1.36 (s, 3H), 1.35-1.24 (m, 3H), 1.28-1.25 (m, 6H), 1.19-1.11 (m, 2H), 0.87 (s, 9H), 0.04 (d, $J = 5.5$ Hz, 6H); ^{13}C NMR (100 MHz, CDCl_3) δ 165.0, 155.3, 150.4, 119.7, 73.3, 71.3, 70.4, 66.4, 65.4, 55.3, 50.9, 45.2, 34.9, 34.3, 31.2, 28.9, 28.8, 25.9, 24.5, 23.8, 23.5, 18.2, -4.6, -4.7; IR (Neat Film) 3331, 2927, 2855, 1746, 1662, 1627, 1531, 1459, 1385, 1254, 1191, 1116 cm^{-1} ; HRMS (ESI) m/z Calcd for $\text{C}_{28}\text{H}_{50}\text{NO}_7\text{Si}$ ($\text{M}+1$) $^+$ 540.3357, found 540.3361.

(7,7-dimethyl-1,6-dioxaspiro[2.5]octan-5-yl)methyl((1R,4s)-4-((S,Z)-4-hydroxypent-2-enamido)cyclohexyl)methyl carbonate (25)

Removal of TBS group in **24** (10 mg, 0.01 mmol) was accomplished following the procedure as for conversion of **14** to **15**, to give 7.5 mg (95%) of **25** as an oil. ^1H NMR (400 MHz,

CDCl₃) δ 6.15 (dd, *J* = 5.5, 11.9 Hz, 1H), 5.94 (br d, 1H), 5.75 (dd, *J* = 1.5, 12.0 Hz, 1H), 5.35 (s, 1H), 4.80-4.72 (m, 1H), 4.20-4.07 (m, 4H), 4.04 (d, *J* = 6.6 Hz, 2H), 2.58 (q, *J* = 4.6 Hz, 2H), 2.05-1.82 (m, 4H), 1.74-1.63 (m, 7H), 1.38-1.33 (m, 6H), 1.24 (s, 3H), 1.19-1.13 (m, 2H); ¹³C NMR (100 MHz, CDCl₃) δ 164.8, 154.3, 148.9, 122.2, 73.3, 70.3, 69.4, 65.4, 63.6, 54.3, 49.9, 44.5, 41.5, 33.9, 33.3, 30.2, 28.7, 27.7, 23.3, 22.2, 21.8; IR (Neat Film) 3321, 2931, 2859, 1745, 1657, 1535, 1453, 1263, 1190 cm⁻¹; HRMS (ESI) *m/z* Calcd for C₂₂H₃₆NO₇ (M + 1)⁺ 426.2492, found 426.2475.

(2S,Z)-5-(((1S,4R)-4-((((7,7-dimethyl-1,6-dioxaspiro[2.5]octan-5-yl)methoxy)carbonyloxy)methyl)cyclohexylamino)-5-oxopent-3-en-2-yl acetate (3)

Acetylation of **25** (7 mg, 0.01 mmol) was achieved following the same procedure used for conversion of **15** to the corresponding acetate **1**, to give 6 mg (78%) of **3** as an oil. ¹H NMR (400 MHz, CDCl₃) δ 7.20 (br d, 1H), 5.84-5.74 (m, 2H), 5.65 (dd, *J* = 9.1, 11.8 Hz, 1H), 4.20-4.08 (m, 4H), 4.02 (d, *J* = 6.9 Hz, 2H), 2.57 (q, *J* = 4.6 Hz, 2H), 2.06 (s, 3H), 2.04-1.85 (m, 2H), 1.84-1.80 (m, 1H), 1.78-1.63 (m, 6H), 1.44-1.35 (m, 8H), 1.25 (s, 3H), 1.20-1.12 (m, 2H); ¹³C NMR (100 MHz, CDCl₃) δ 171.4, 164.9, 155.3, 137.4, 125.6, 73.3, 71.7, 70.4, 69.1, 66.4, 55.4, 50.9, 45.3, 42.4, 35.1, 34.3, 31.2, 28.9, 28.8, 24.4, 24.2, 23.5, 21.2, 20.3; ²¹IR (Neat Film) 3326, 2932, 1742, 1668, 1631, 1532, 1451, 1371, 1252, 1119, 1048 cm⁻¹; HRMS (ESI) *m/z* Calcd for C₂₄H₃₈NO₈ (M + 1)⁺ 468.2597, found 468.2584.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgment

This work was supported by NCI Cancer Center Core grant CA21765 and by the American Lebanese Syrian Associated Charities (ALSAC), St. Jude Children's Research Hospital.

References

1. Rymond B. Targeting the spliceosome. *Nat. Chem. Biol* 2007;3:533–535. [PubMed: 17710096]
2. Kotake Y, Sagane K, Owa T, Mimori-Kiyosue Y, Shimizu H, Uesugi M, Ishihama Y, Iwata M, Mizui Y. Splicing factor SF3b as a target of the antitumor natural product pladienolide. *Nat. Chem. Biol* 2007;3:570–575. [PubMed: 17643112]
3. Mizui Y, Sakai T, Iwata M, Uenaka T, Okamoto K, Shimizu H, Yamori T, Yoshimatsu K, Asada M. Pladienolides, new substances from culture of *Streptomyces platensis* Mer-11107. III. In vitro and in vivo antitumor activities. *J. Antibiot* 2004;57:188–196. [PubMed: 15152804]
4. Nakajima H, Hori Y, Terano H, Okuhara M, Manda T, Matsumoto S, Shimomura K. New antitumor substances, FR901463, FR901464 and FR901465. II. Activities against experimental tumors in mice and mechanism of action. *J. Antibiot* 1996;49:1204–1211. [PubMed: 9031665]
5. Nakajima H, Sato B, Fujita T, Takase S, Terano H, Okuhara M. New antitumor substances, FR901463, FR901464 and FR901465. I. Taxonomy, fermentation, isolation, physicochemical properties and biological activities. *J. Antibiot* 1996;49:1196–1203. [PubMed: 9031664]
6. Nakajima H, Takase S, Terano H, Tanaka H. New antitumor substances, FR901463, FR901464 and FR901465. III. Structures of FR901463, FR901464 and FR901465. *J. Antibiot* 1997;50:96–99. [PubMed: 9066774]
7. Kaida D, Motoyoshi H, Tashiro E, Nojima T, Hagiwara M, Ishigami K, Watanabe H, Kitahara T, Yoshida T, Nakajima H, Tani T, Horinouchi S, Yoshida M. Spliceostatin A targets SF3b and inhibits both splicing and nuclear retention of pre-mRNA. *Nat Chem Biol* 2007;3:576–583. [PubMed: 17643111]
8. Thompson CF, Jamison TF, Jacobsen EN. FR901464: total synthesis, proof of structure, and evaluation of synthetic analogues. *J. Am. Chem. Soc* 2001;123:9974–9983. [PubMed: 11592876]

9. Motoyoshi H, Horigome M, Ishigami K, Yoshida T, Horinouchi S, Yoshida M, Watanabe H, Kitahara T. Structure-activity relationship for FR901464: a versatile method for the conversion and preparation of biologically active biotinylated probes. *Biosci. Biotechnol. Biochem* 2004;68:2178–2182. [PubMed: 15502365]
10. Motoyoshi H, Horigome M, Watanabe H, Kitahara T. Total synthesis of FR901464: second generation. *Tetrahedron* 2006;62:1378–1389.
11. Albert Brian J, Sivaramakrishnan A, Naka T, Czaicki Nancy L, Koide K. Total syntheses, fragmentation studies, and antitumor/antiproliferative activities of FR901464 and its low picomolar analogue. *J. Am. Chem. Soc* 2007;129:2648–2659. [PubMed: 17279752]
12. Ohta T, Miyake T, Seido N, Kumobayashi H, Takaya H. Asymmetric Hydrogenation of Olefins with Aprotic Oxygen Functionalities Catalyzed by BINAP-Ru(II) Complexes. *J. Org. Chem* 1995;60:357–363.
13. Tuttle JB, Ouellet SG, MacMillan DWC. Organocatalytic Transfer Hydrogenation of Cyclic Enones. *J. Am. Chem. Soc* 2006;128:12662–12663. [PubMed: 17002356]
14. Ouellet SG, Walji AM, Macmillan DWC. Enantioselective organocatalytic transfer hydrogenation reactions using Hantzsch esters. *Acc. Chem. Res* 2007;40:1327–1339. [PubMed: 18085748]
15. Corey EJ, Chaykovsky M. Dimethyloxosulfonium methylide and dimethylsulfonium methylide. Formation and application to organic synthesis. *J. Am. Chem. Soc* 1965;87:1353–1364.
16. Yoon NM, Pak CS, Brown HC, Krishnamurthy S, Stocky TP. Selective reductions. XIX. Rapid reaction of carboxylic acids with borane-tetrahydrofuran. Remarkably convenient procedure for the selective conversion of carboxylic acids to the corresponding alcohols in the presence of other functional groups. *J. Org. Chem* 1973;38:2786–2792.
17. Mitsunobu O. The use of diethyl azodicarboxylate and triphenylphosphine in synthesis and transformation of natural products. *Synthesis* 1981:1–28.
18. Valverde S, Martin-Lomas M, Herradon B, Garcia-Ochoa S. The reaction of carbohydrate derived alkoxyaldehydes with methoxycarbonylmethylenetriphenylphosphorane: stereoselective synthesis of b-unsaturated esters. *Tetrahedron* 1987;43:1895–1901.
19. Nagarajan S, Ganem B. Chemistry of naturally occurring polyamines. 11. Unsaturated spermidine and spermine derivatives. *J. Org. Chem* 1987;52:5044–5046.
20. Pessah N, Reznik M, Shamis M, Yantiri F, Xin H, Bowdish K, Shomron N, Ast G, Shabat D. Bioactivation of carbamate-based 20(S)-camptothecin prodrugs. *Bioorg. Med. Chem* 2004;12:1859–1866. [PubMed: 15051055]
21. We have seen that compounds **1**, **2**, and **3** show evidence of energetically similar chair and twist-boat conformations in the cyclohexane rings, which appear to equilibrate on the NMR timescale, as has been previously observed in the ¹³C NMR spectra for some cis-1,4-substituted cyclohexane derivatives with bulky or heteroatom substituents, see for example: (a) Gordon MD, Quin LD. Carbon-13 Nuclear Magnetic Resonance Spectroscopy in the Study of Conformational Effects among Cyclohexyl Phosphorus Compounds. *J. Org. Chem* 1976;41:1690–1694. 1694 (b) Gill G, Pawar DM, Noe EA. Conformational Study of cis-1,4-Di-tert-butylcyclohexane by Dynamic NMR Spectroscopy and Computational Methods. Observation of Chair and Twist-Boat Conformations. *J. Org. Chem* 2005;70:10727–10731. 10731

Figure 1.
The structures of pladienolide B (left) and FR901464 (right)

Figure 2.

A 3D representation of an example of an overlay of the presumed key interaction groups in a low energy conformation of pladienolide and FR901464, showing that the epoxy group and the carbonyloxy groups are the same distance in both molecules, and that the methyl-2,4-pentadienyl groups can readily adopt a similar positioning. The overlays were prepared using the Molecular Operating System (MOE 2007.09, Chemical Computing Group, Inc.) using the Flexible Alignment function with both molecules following a conformational minimization using MOE default settings.

Figure 3.
The initial synthetic targets in our study that matched our hypothetical pharmacophore.

Figure 4. Cell cycle analysis following treatment with compound 1. The A) A549, B) JeKo-1, and C) WiDr cancer cell lines were treated with compound 1 for 18 hours, then permeabilized and stained with propidium iodide for cell cycle analysis by flow cytometry.

Scheme 1.

The Synthesis of Analog 1a

^aReagents and conditions: (a) (2*R*,5*R*)-5-benzyl-3-methyl- (5-methylfuran-2-yl-imidazolidin-4-one, *t*-butyl Hantzsch ester, Cl₃CCO₂H, Et₂O, 4° C; (b) NaH, (CH₃)₃SOI, DMSO; (c) DIBALH, THF, -78° C; (d) BH₃·THF, THF; (e) Oxalyl chloride, DMSO, Hunig's base, -78° C, CH₂Cl₂; (f) i) Ph₃PC(CH₃)CO₂Et, benzene; ii) DIBALH, -78° C, toluene; (g) 2-mecapto benzothiazole, DIAD, THF; (h) Ammonium molybdate, H₂O₂, EtOH; (i) i) TFA, CH₂Cl₂ ii) (S,Z)-4-(TBSO)pent-2-enoic acid, HBTU, Hunig's base, CH₃CN; (j) LHMDs, **7a**, THF, -78° C to rt; (k) Bu₄NF, THF, 0° C; l) Ac₂O, Et₃N, CH₂Cl₂.

Scheme 2. The Synthesis of Analogs **2** and **3b**

^bReagents and conditions: (a) i) MsCl, Et₃N, 0° C, CH₂Cl₂; ii) NaN₃, DMF, 70° C; (b) i) TFA, CH₂Cl₂; ii) (S,Z)-4-(TBSO)pent-2-enoic acid, HBTU, Hunig's base, CH₃CN; (c) Ph₃P, H₂O, THF; (d) **7b**, 4-nitrophenyl chloroformate, Et₃N, THF; (e) Bu₄NF, THF, 0° C; (f) Ac₂O, Et₃N, CH₂Cl₂; (g) 4-nitrophenyl chloroformate, Et₃N, THF; (h) **7b**, DMAP, CH₃CN.

Table 1*In vitro* cytotoxicity of compound **1** following a 72-hour exposure.

Cell line	Pladienolide B ^{2,3}	FR901464 ⁵	Compound 1
	Published IC ₅₀ (μ M)	Published IC ₅₀ (μ M)	IC ₅₀ $\pm$ S.E.M (μ M) [*]
A549	0.0014	0.0013	4.65 $\pm$ 0.37
JeKo-1	NA [†]	NA	0.10 $\pm$ 0.002
JVM-2	NA	NA	0.13 $\pm$ 0.03
MCF-7	0.0004	0.0018	2.29 $\pm$ 0.38
OVCAR-3	0.0004	NA	1.98 $\pm$ 0.63
PC-3	0.0023	NA	2.12 $\pm$ 0.57
WiDr	0.00086	NA	2.03 $\pm$ 0.20

^{*} Experimental IC₅₀ values represent the mean $\pm$ S.E.M. from three independent experiments.

[†] NA=Data not available.