Published in final edited form as:

Int J Cancer. 2009 April 1; 124(7): 1721–1726. doi:10.1002/ijc.24014.

Children with Endemic Burkitt Lymphoma are Deficient in EBNA1-Specific IFN- γ T Cell Responses

Ann M. Moormann¹, Kevin N. Heller², Kiprotich Chelimo³, Paula Embury¹, Robert Ploutz-Snyder⁴, Juliana A. Otieno⁵, Margaret Oduor⁵, Christian Münz², and Rosemary Rochford⁴

¹Center for Global Health and Diseases, Case Western Reserve University, Cleveland, OH

²Laboratory of Viral Immunobiology and Christopher H. Browne Center for Immunology and Immune Diseases, The Rockefeller University, New York NY

³Kenya Medical Research Institute, Center for Global Health and Research, Kisumu, Kenya

⁴Department of Microbiology and Immunology, SUNY Upstate Medical University, Syracuse, NY

⁵Nyanza Provincial General Hospital, Kenya Ministry of Health, Kisumu, Kenya

Abstract

Endemic Burkitt lymphoma (eBL) is the most common childhood cancer in equatorial Africa and is linked to Epstein-Barr virus (EBV) and *Plasmodium falciparum* co-infections early in life. Epstein-Barr nuclear antigen 1 (EBNA1) is the sole viral latent antigen expressed in BL tumors. Loss of EBNA1-specific immune surveillance could allow eBL emergence. Therefore, EBNA1-specific T cell responses were analyzed by IFN- γ ELISPOT in Kenyan children with eBL and compared to healthy children with divergent malaria exposure. Significantly fewer children with eBL, 16% (7/44) had EBNA1-specific IFN- γ responses in contrast to healthy children living in a malaria holoendemic area or in an area with sporadic malaria transmission, 67% (40/60) and 72% (43/60) responders, respectively ($p < 0.003$). Children with eBL maintained IgG₁ dominated antibody responses to EBNA1 similar to healthy children suggesting a selective loss of IFN- γ secreting EBNA1-specific T cells in the presence of intact humoral immunity. CD8⁺ T cell responses to EBV lytic and latent antigens not expressed in the tumors were similarly robust in eBL patients compared to healthy children. In addition, CD4⁺ T cell responses to a malaria protein, merozoite surface protein 1, were present in lymphoma patients. This study demonstrates a selective loss of EBNA1-specific T cell responses in children with eBL, and suggests a potential immunotherapeutic target for this EBV-associated lymphoma.

Keywords

Burkitt lymphoma; malaria; EBNA1; IFN-gamma

¹Correspondence should be addressed to Ann M Moormann, Case Western Reserve University, Center for Global Health and Diseases, 2103 Cornell Road WRB 4-130, Cleveland Ohio 44106-7286; moorms@case.edu; phone: (216) 368-5144; fax: (216) 368-4825..

Author contributions A.M.M. designed experiments and wrote the manuscript; K.N. H performed EBNA1 ELISAs; K.C performed ELISPOTS, P.E. performed ELISPOTS, R.P.-S. performed biostatistical analysis; J.O. recruited patients for study and provided clinical input; M. O. performed histopathology; C. M. designed experiments; and R. R., designed experiments and wrote the manuscript.

Statement of novelty and impact: Our preliminary and novel discovery of a select deficiency in EBNA1-specific IFN- γ mediated T cell immunity in children with endemic Burkitt lymphoma will shed light on etiologic mechanisms of this cancer and potentially drive clinical research efforts focused on immunotherapies for EBV-associated lymphomas.

INTRODUCTION

Epstein-Barr virus (EBV) is etiologically associated with a number of malignancies that occur in immune competent hosts including nasopharyngeal carcinoma, mixed-cellularity Hodgkin's disease, and endemic Burkitt lymphoma (eBL)¹. While there are different patterns of viral gene expression in EBV-associated malignancies, all EBV tumors express the viral latent protein Epstein-Barr nuclear antigen 1 (EBNA1)¹. An important question is how these tumors escape the host immune response when viral antigens are present.

Endemic BL is the most common childhood malignancy in Equatorial Africa and occurs in children between the ages of 2 and 14 years with a peak incidence of 6 years of age². Holoendemic malaria is thought to be an etiologic co-factor in this malignancy. While the exact mechanisms are unknown, repeated malaria infections in young children are thought to drive B-cell expansion and suppress T cell immunity thus allowing for the emergence of a malignant clone. We recently reported that 5-9 year old children living in a holoendemic malaria region had depressed CD8⁺ T cell responses against both EBV lytic and latent antigens³, supporting a model for malaria driven immune modulation as one potential etiologic mechanism. However, nothing is known of the EBV- specific T cell responses in children with eBL.

EBNA1 is the only latent protein expressed in eBL tumors. Escape from immune surveillance by eBL was originally thought to occur because BL cells have defects in MHC Class I antigen processing and the gly-ala repeat domain of the EBNA1 inhibits proteasomal processing and EBNA1 mRNA translation⁴⁻⁶. In contrast to this lack of CD8⁺ T cell recognition of EBNA1 and eBL, recent studies have shown that EBNA1-specific CD4⁺ T cells can be isolated from nearly all EBV-seropositive healthy adults^{7, 8}, and these CD4⁺ T cells are cytolytic for constitutively HLA class II expressing BL cell lines which express EBNA1 protein⁹. However, if the host is able to mount a cytolytic CD4⁺ T cell response to EBNA1, why do eBL tumors escape host immune control? In this study, we examined EBNA1-specific CD4⁺ T cell responses in comparison to EBV-specific CD8⁺ and malaria-specific CD4⁺ T cell responses in Kenyan children with eBL, and compared these to healthy Kenyan children with divergent patterns of malaria exposure. This allowed us to address the question of whether children with eBL were able to mount an EBNA1-specific T cell response.

MATERIALS AND METHODS

Study Population and Design

From August 2005 to March 2006, we prospectively enrolled 44 Kenyan children admitted to Nyanza Provincial General Hospital (NPGH), Kisumu, Kenya. Confirmation of BL diagnosis was made by histology from a fine needle aspirate stained with May-Grunewald Giemsa and reviewed by Dr. Margaret Oduor, the Clinical Pathologist at hospital. Blood samples were taken from children with eBL upon admission to the hospital and prior to administration of the first dose of cytotoxic chemotherapy.

Since eBL is the most prevalent pediatric cancer in this region enrolling enough children diagnosed with cancer other than eBL was not feasible. Therefore, we enrolled 120 healthy children from two regions of Kenya with divergent risk for BL and malaria exposure. The first region is Kisumu District where malaria is holoendemic, i.e. intense, perennial malaria transmission and the incidence of eBL is high^{2, 10}. The second region is the Nandi District in the highlands of western Kenya where malaria transmission is sporadic and unstable and the risk of eBL is very low^{2, 10}. In addition, adults from the Kisumu region were included as a referent group and as a basis of comparison for EBNA1-specific immunity in healthy

US adults⁸. The mean ages of children enrolled in this study were 5.7, 6.2 and 6.9 years for Nandi, Kisumu and eBL children, respectively. The three groups of children were compared for hematological indices and found to have similar values within normal ranges for their age. The mean white blood cell counts were 8.4, 11.7 and 10×10^3 per microliter and mean hemoglobin was 10.4, 11.3 and 13.7 gm/dl for Nandi, Kisumu and eBL children, respectively.

Ethical approval for the study was obtained from Institutional Review Boards at Case Western Reserve University (CWRU) and SUNY Upstate Medical University and the Ethical Review Committee at the Kenya Medical Research Institute (KEMRI).

Isolation of Peripheral Blood Mononuclear Cells

Peripheral blood was collected in sodium heparinized tubes and transported to the CWRU/KEMRI laboratory located at the Centre for Global Health Research, in Kisumu for processing the same day. Peripheral blood mononuclear cells (PBMCs) were separated from whole blood by Ficoll-Hypaque density gradient centrifugation and suspended in RPMI-1640 (GIBCO, Invitrogen, Paisley, Scotland, UK) supplemented with 10% heat inactivated human AB serum, 50 µg/ml gentamicin, 10mM Hepes and 2 mM glutamine for cytokine stimulation assays. Plasma and whole blood was frozen for later serologic and DNA analyses.

Assessment of EBV- and malaria-specific T cell responses by IFN-γ ELISPOT

IFN-γ ELISPOT assays and selection of HLA Class I peptides from EBV lytic (BRLF1, BZLF1, BMLF1) and EBV latent (EBNA 3A, 3B, 3C) antigens has been previously described³. Fifty-one overlapping EBNA1 peptides covering the C' terminal domain (amino acids 400-641) as previously described⁸ were used as a total pool (EATP) and 5 sub-pools consisting of 10-11 peptides in each sub-pool (EAI-V). IFN-γ responses to phytohemagglutinin (Sigma-Aldrich, St. Louis, MO) at 1 µg/ml and a recombinant blood stage malaria protein (merozoite surface protein-1, MSP1, 42kDa, 3D7 strain)¹¹ were controls. The specificity of EBV- and malaria-antigen responses was tested on PBMCs from 20 malaria unexposed adult American volunteers. EBV seropositive American PBMC donors had robust cytokine responses to EBV peptides but lacked responses to malaria peptides (data not shown). PBS was added to 0.5×10^6 PBMCs per well as the negative control and represented the background level of IFN-γ produced by unstimulated cells. The number of spot forming units (SFU) per well was counted using ImmunoSpot Scanning & Imaging Software, Version 4.0 (Pharmingen).

Determination of EBNA1-specific IgG subclass

ELISA for IgG subclass titers were done as described by the manufacturer (Sigma/Diamedix; Miami, FL). Plasma samples were diluted at 1:20 and 1:10 and run in triplicate. EBNA1 and IgG subclass titers were assessed by evaluating plasma samples from patients and healthy volunteers at serial dilutions from 1:25 to 1:3200. After plotting absorption values, the point of inflection was calculated to determine the IgG titer for each subclass.

Determination of EBV viral load

EBV viral load was determined by whole peripheral blood DNA extraction as previously described with minor modifications¹². For the real-time quantitative PCR, EBV and β-actin were amplified using previously described primer/probes¹² using BioRad MasterMix in the BioRad IQ3 cyclor. To quantitate EBV viral load, serial dilutions of cellular DNA obtained from an EBV negative BL line (BL41) were used to generate a standard curve for total cellular DNA. EBV viral load was calculated using serial dilutions of an EBV plasmid

containing the amplified EBV region. For statistical analysis, samples below the threshold of detection were changed to 0.5 before natural log (Ln) transformation.

Statistical Analysis

Analyses were conducted using SAS Version 8.2 (Carey, NC). An IFN- γ ELISPOT was considered positive if the proportion of SFU in the stimulated well was significantly greater compared to the unstimulated background well using a chi-square Fisher's exact test ($p < 0.05$); a more stringent criteria than merely counting a positive response as more than 5 SFC above background or as two-fold above the background. Proportions of responders were compared across groups (eBL, Kisumu, Nandi) using Logistic Regression Analysis, with Odds Ratios and significance values calculated comparing the percent of eBL responders relative to each the other groups. P-values less than 0.05 were taken to indicate significance while P-values between 0.06 and 0.08 were taken to be indicative of a trend. Responses to EBV peptides were found only from PBMC of EBV seropositive donors (data not shown for EBV seronegative donors). The magnitude of IFN- γ responses was generated using Analysis of Variance to compare outcomes by group, with Post-Hoc tests set to compare all pairs. Tukey's post-hoc were used for pairwise comparisons where we met homogeneity of variance assumption, otherwise Games-Howell was used (which adjusts for heterogeneity of variance).

RESULTS

Children diagnosed with eBL are deficient in EBNA1-specific, IFN- γ producing T cell responses

To evaluate EBV-specific T cell immunity in children with eBL, three groups of Kenyan children were enrolled in this study: children diagnosed with eBL ($n=44$), healthy children living in the holoendemic malaria region of Kisumu District (herein referred to as Kisumu) ($n=60$), and healthy children living in the epidemic-prone malaria region of Nandi District (herein referred to as Nandi) ($n=60$). We first asked whether there were differences in frequency of EBNA1-specific, IFN- γ producing T cell responses in the children with eBL compared to healthy children regardless of malaria exposure. To do this, freshly isolated PBMC from each of the groups were stimulated with an overlapping peptide library of the C-terminal domain of EBNA1 (aa400-641), which contains all but one described CD4⁺ T cell epitopes of this antigen. This peptide library was divided in five EBNA1 peptide sub-pools consisting of 10 or 11 peptides each (EAI, EAII, EAIII, EAIIV and EAV), or used as a total pool containing all 51 peptides (EATP). The peptide pools were previously shown to elicit IFN- γ from EBNA1-specific CD4⁺ T cells in the majority of healthy adult EBV carriers⁸. Responses to the EBNA1 peptides were measured by IFN- γ ELISPOT assays. The proportion of positive responders in each group was compared and is presented in Table 1. Children with eBL had significantly fewer IFN- γ responders to the total pool of EBNA1 peptide (7/44) compared to healthy Nandi (43/60) and Kisumu (40/60) children. When we examined responses to the EBNA1 peptide sub-pools (i.e. EAI-V), we also observed fewer responders in the eBL patients compared to the healthy children from both Nandi and Kisumu for all pools although this achieved statistical significance for only 1 of the 5 pools for Kisumu and 3 of 5 pools for Nandi children (Table 1). Overall, the odds of having an IFN- γ response to EBNA1 are significantly greater in healthy children compared to children with eBL (Table 1).

When PBMCs isolated from the eBL patients were stimulated with PHA, a general immune stimulant, 95% responded, similar to percent response observed in Kisumu (97%) and Nandi (92%) children, suggesting that children with eBL were not globally immune deficient. As a second test of immune competency, the proportion of eBL children responding to the

malaria antigen MSP-1—primarily recognized by CD4⁺ T cells¹¹— was similar to the proportion of responders observed in both Kisumu children (36.4% and 25.0%, respectively) who reside in a malaria holoendemic area (Table 1). The proportion of responders from Nandi was lower with only 13.8% responders as would be expected from children living in an area with unstable malaria transmission ($p=0.01$).

We next compared the capacity of children with eBL and healthy children to respond to known EBV-specific CD8⁺ T cell epitopes using peptide pools (EP1/EP2) that were previously described and contain peptides for both EBV lytic (BRLF1, BZLF1, BMLF1) and latent (EBNA3A, 3B and 3C) proteins³. No differences in the frequencies of IFN- γ responses to EBV-specific CD8⁺ T cell epitopes in children with eBL compared to Kisumu or Nandi children were observed (Table 1) suggesting that EBV CD8⁺ T cell responses are not deficient in the children with eBL. Our previous results showed fewer IFN- γ responses to EBV-specific CD8⁺ T cell epitopes restricted to 5-9 year old Kisumu children compared to Nandi children³. The healthy children examined here spanned the same age range (i.e. 2-14 years) as the eBL children. Therefore the age-specific (5-9 year olds) deficiency in IFN- γ responses to EBV-specific CD8⁺ T cell epitopes was balanced by responses from the older and younger Kisumu children included in this analysis. When we examined IFN- γ EBV-specific CD8⁺ T cell responses only from 5-9 year old eBL children, they were comparable to the healthy non-malaria exposed healthy 5-9 year olds from Nandi (data not shown).

Although the frequency of EBNA1-specific IFN- γ responders was lower in eBL children, we wanted to determine if the magnitude of this response in the 7 eBL children who did respond was similar to children from Kisumu or Nandi. Comparisons between eBL, Kisumu and Nandi children in the magnitude of their IFN- γ producing T cell responses to the individual EBNA1 peptide pools, the EBNA total pool (EATP) and to the malaria recombinant protein, MSP1 are shown in Figure 1. We found magnitude of IFN- γ responses to EATP to be significantly lower in eBL children compared to Nandi ($p<0.001$) and Kisumu ($p<0.001$) children, while Kisumu and Nandi children did not differ from each other ($p=0.562$). When examining each EBNA1 peptide subpool, we found a significant difference overall between the three groups for EAI ($p<0.003$) but the post-hoc comparisons revealed that Nandi children had significantly higher IFN- γ levels compared to eBL ($p<0.012$) and Kisumu ($p<0.02$) children, while the eBL and Kisumu children were not significantly different from one another ($p=.701$). For the EAIII pool there was a significant difference overall ($p<0.002$), with post-hoc comparisons revealing that responses in eBL children were significantly lower than in Nandi ($p<0.003$) and Kisumu ($p<0.011$) children, while the level of responses between the Kisumu and Nandi children did not differ ($p=0.591$). This was also true for EAV, with a significant difference overall ($p<0.007$) and post-hoc comparisons showing the level of IFN- γ responses from eBL children significantly lower than from Nandi ($p<0.007$) and Kisumu ($p<0.029$) children, but healthy children with divergent malaria exposure did not differ from one another ($p=0.836$). There were no significant differences in IFN- γ responses to EAI and EAIV among the three groups.

When we compared the magnitude of IFN- γ expressed to MSP1, children with eBL had a significantly higher mean cytokine level compared to Kisumu and Nandi children ($p<0.05$) as shown in Figure 1, demonstrating that children with eBL were not deficient in other CD4⁺ T cell mediated responses.

Children with eBL have similar humoral immune response to EBNA1 as healthy children

One possible explanation for the inability of eBL patients to respond to EBNA1 peptides is that the T cell mediated IFN- γ response to EBNA1 was absent prior to BL tumorigenesis. If this was the case, then we would predict an absence of EBNA1 specific IgG₁ antibodies, because Th1 CD4⁺ T cells assist B cells to make IgG₁ antibodies. To evaluate this

possibility, IgG isotype specific ELISAs were done to determine if children with eBL had an EBNA1 antibody response and if so, what the dominant IgG isotype of this response was. Children with eBL produced IgG₁ subclass antibodies to EBNA1 similar to healthy children and adults (Figure 2). This also argues against Th2 polarization by means of alternate cytokines, such as IL-10, being produced in response to EBNA1. Interestingly, the levels of anti-EBNA1 IgG₁ were higher in the eBL patients than in the other groups. IgG₄ was detected in a small percentage of children with eBL but not IgG₂ or IgG₃. IgG₂ was only detected in one of the Nandi children (Figure 2).

EBV viral load tends to be higher in children without EBNA1-specific IFN- γ responses

A recent study suggested that control of EBV correlates with CD4⁺ T cell rather than CD8⁺ T cell responses¹³. To test whether there was a link between EBNA1- specific, IFN- γ producing T cell responses and EBV viral load, we extracted DNA from whole blood and measured EBV viral load by RTQ-PCR¹². The mean EBV viral load was then compared within each group of children stratified by EBNA1-specific IFN- γ response (Figure 3). EBV viral load data were positively skewed, with a few extremely high values, and several very low or undetectable levels. As such, we analyzed natural log transformed EBV viral load using distribution-free non-parametric statistics. We employed the Mann-Whitney non-parametric statistic to compare IFN- γ producing EBNA1 (i.e. EATP) responders to non-responders within our three groups (Nandi, Kisumu, eBL), with Fischer's exact tests of significance. A near-significant trend emerged for eBL and Kisumu children, showing that EBV viral load tended to be higher in children without EBNA1-specific IFN- γ responses in these two groups.

DISCUSSION

Children in Equatorial Africa are at high risk for eBL, and this is linked to both holoendemic malaria exposure and early-age of EBV infection. A long-standing hypothesis is that the repeated malaria infections experienced by children living in malaria holoendemic regions suppresses EBV specific T-cell immunity. Our earlier studies have supported this malariainduced immune modulation model where we showed that 5-9 year old children, living in malaria holoendemic regions have diminished EBV-specific CD8⁺ T cell responses relative to children living in malaria epidemic regions. What wasn't known was whether this diminished response also extended to children with eBL. In addition, although the EBV-specific CD8⁺ T cell responses are targeted towards several of the EBV latent and lytic proteins, only the EBNA1 protein is expressed by eBL tumors. We show here that while children with eBL have a comparable EBV-specific CD8⁺ T cell response to healthy children, the eBL patients show a selective loss of the EBNA1-specific, IFN- γ producing T cells. EBNA1-specific CD4⁺ T cells suppress tumor growth in a BL mouse model, suggesting that the absence of IFN- γ responses to EBNA1 in eBL patients might allow the emergence of the malignant clone. When the loss of the EBNA1-specific IFN- γ response occurred cannot be determined from a cross-sectional study such as this. However, the presence of EBNA1 IgG in most of the eBL patients is suggestive that these children had an EBNA1 IFN- γ response at some point prior to the emergence of the tumor. The key question will be to understand why that loss occurred. One possibility is that EBV viral loads CD4⁺ T cell responses may be exquisitely sensitive to the immunoregulatory influence of *P. falciparum*, particularly with respect to dendritic cell function¹⁶ and that children chronically or repeatedly infected with malaria have impaired antigen presentation whereby T cell responses to EBV antigens are impaired due to a bystander effect. Alternatively, the high levels of EBNA-1 protein expressed by BL tumors could down-regulate the IFN- γ T cell response due to T cell exhaustion.

In this study we compared T responses in children from Kisumu region (both eBL patients and healthy children) and children from the Nandi region. While it was not possible to HLA type the study participants, previous work by our group has shown that there is no significant skewing of the HLA types between these two populations¹⁷. In addition, the EBNA-1 total pool of peptides span the entire C-terminal region of the protein encompassing all possible epitopes with the potential to be presented by any HLA type. Thus, the lack of EBNA1 IFN- γ responders in eBL patients is unlikely to be due to variance in HLA types between populations. We also examined whether there were clinical differences (e.g. site of tumor location, response to treatment) between the 7 of 44 children that did respond and those that did not however, no differences were noted between the two groups (data not shown).

We found that children with elevated EBV viral load tended to lack EBNA-1 specific IFN- γ responses. Since EBNA1 is the only antigen expressed in latently infected memory B cells, this would suggest that those children with lower EBNA1 CD4⁺ T cell responses might have a more difficult time controlling EBV persistence. Clearly more data is necessary but the trend we have observed is novel and intriguing. Ongoing longitudinal studies in our laboratories will allow us to more clearly define the temporal association between elevated EBNA1 antibody levels, EBV-specific T cell responses and EBV viral loads.

The EBNA-1 protein has gly-ala repeats that prevent processing and proper antigen presentation via MHC Class I ⁴⁻⁶. However, there could be cross-reactive epitopes that are exogenously presented to CD8⁺ T cells as has been described in mouse models¹⁸. So we cannot exclude a loss of EBNA1 to CD8 epitopes in our eBL patients. Of note, however is that the eBL patients were able to respond to EBV peptide pools that contain epitopes specific for well described EBV CD8⁺ T cell antigens (e.g. BRLF1, BZLF1, BMLF1, EBNA- 3A, -3B, -3C) thus suggesting that the lack of EBNA1-specific IFN- γ responses in the eBL patients are likely to be CD4⁺ T mediated. In addition, the eBL patients mounted robust IFN- γ responses to the malaria antigen which is mediated by CD4⁺ T cells¹⁹.

Consistent with our result in eBL, the report by Heller and colleagues shows that there is loss of EBNA1-specific CD4⁺ T cell responses in patients with EBV-positive (IJC in press), but not EBV-negative Hodgkin lymphoma. Together, these studies argue that selective loss of EBNA1-specific IFN- γ response may be critical to the emergence of EBV-associated lymphomas. It is tempting to speculate that the absence of EBNA1- specific CD4⁺ T cells predisposes EBV carriers to lymphoma development, even though the mechanisms leading to this select dysfunction most likely differ between eBL and Hodgkin lymphoma. Longitudinal studies of eBL patients will determine if restoration of EBNA1-specific T cell responses is associated with long-term survival. A universal loss in EBNA1-specific immune control could be exploited for immunotherapeutic approaches that induce cytolytic responses to EBNA1 in patients with EBV-associated malignancies.

Acknowledgments

We thank the families and children for their participation in this study. We thank Dorine Omenha for project coordination and Nancy Fiore and Steven Asito Amolo for technical assistance. We thank Carole Long for the use of the recombinant malaria proteins for this study. We thank Mrs. Somia and the Equatorial Africa Children's Cancer Fund (<http://www.eachcancerfund.org/>) for securing the cytotoxic chemotherapy to treat all eBL patients admitted to NNPGH. This manuscript was approved for publication by the Director of the Kenya Medical Research Institute. Research funded by NIH K08-AI51565 (Moormann), NIAID-AI43906 (Kazura), Children's Cancer Research Fund, a California non-profit organization (Heller), NCI CA108609 and NCI CA101741 (Münz), NCI CA102667 (Rochford) and Fogarty International Center D43 TW006576-05 (King).

REFERENCES

1. Crawford, DH. Biology and disease associations of Epstein-Barr virus; *Philos Trans R Soc Lond B Biol Sci.* 2001. p. 461-73.2001;356(1408):461-73. Available from http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=11313005
2. Mwanda, OW.; Rochford, R.; Moormann, AM.; MacNeil, A.; Whalen, C.; Wilson, ML. Burkitt's lymphoma in Kenya: geographical, age, gender and ethnic distribution; *East Afr Med J.* 2004. p. S68-77.2004(8 Suppl):S68-77. Available from http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=15622605
3. Moormann, AM.; Chelimo, K.; Sumba, PO.; Tisch, DJ.; Rochford, R.; Kazura, JW. Exposure to holoendemic malaria results in suppression of Epstein-Barr virus-specific T cell immunosurveillance in Kenyan children; *J Infect Dis.* 2007. p. 799-808.2007;195(6):799-808. Available from http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=17299709
4. Blake, N.; Lee, S.; Redchenko, I.; Thomas, W.; Steven, N.; Leese, A.; Steigerwald-Mullen, P.; Kurilla, MG.; Frappier, L.; Rickinson, A. Human CD8+ T cell responses to EBV EBNA1: HLA class I presentation of the (Gly-Ala)-containing protein requires exogenous processing; *Immunity.* 1997. p. 791-802.1997;7(6):791-802. Available from http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=9430224
5. Levitskaya, J.; Coram, M.; Levitsky, V.; Imreh, S.; Steigerwald-Mullen, PM.; Klein, G.; Kurilla, MG.; Masucci, MG. Inhibition of antigen processing by the internal repeat region of the Epstein-Barr virus nuclear antigen-1; *Nature.* 1995. p. 685-8.1995;375(6533):685-8. Available from http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=7540727
6. Yin, Y.; Manoury, B.; Fahraeus, R. Self-inhibition of synthesis and antigen presentation by Epstein-Barr virus-encoded EBNA1; *Science.* 2003. p. 1371-4.2003;301(5638):1371-4. Available from http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=12958359
7. Nikiforow, S.; Bottomly, K.; Miller, G.; Munz, C. Cytolytic CD4(+)-T-cell clones reactive to EBNA1 inhibit Epstein-Barr virus-induced B-cell proliferation; *J Virol.* 2003. p. 12088-104.2003;77(22):12088-104. Available from http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=14581546
8. Heller, KN.; Upshaw, J.; Seyoum, B.; Zebroski, H.; Munz, C. Distinct memory CD4+ T-cell subsets mediate immune recognition of Epstein Barr virus nuclear antigen 1 in healthy virus carriers; *Blood.* 2007. p. 1138-46.2007;109(3):1138-46. Available from http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=16985171
9. Paludan, C.; Bickham, K.; Nikiforow, S.; Tsang, ML.; Goodman, K.; Hanekom, WA.; Fonteneau, JF.; Stevanovic, S.; Munz, C. Epstein-Barr nuclear antigen 1-specific CD4(+) Th1 cells kill Burkitt's lymphoma cells; *J Immunol.* 2002. p. 1593-603.2002;169(3):1593-603. Available from http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=12133989
10. Rainey, JJ.; Omenah, D.; Sumba, PO.; Moormann, AM.; Rochford, R.; Wilson, ML. Spatial clustering of endemic Burkitt's lymphoma in high-risk regions of Kenya; *Int J Cancer.* 2007. p. 121-7.2007;120(1):121-7. Available from http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=17019706
11. Singh, S.; Kennedy, MC.; Long, CA.; Saul, AJ.; Miller, LH.; Stowers, AW. Biochemical and immunological characterization of bacterially expressed and refolded Plasmodium falciparum 42-kilodalton C-terminal merozoite surface protein 1; *Infect Immun.* 2003. p. 6766-74.2003;71(12):6766-74. Available from

http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=14638762

12. Moormann, AM.; Chelimo, K.; Sumba, OP.; Lutzke, ML.; Ploutz-Snyder, R.; Newton, D.; Kazura, J.; Rochford, R. Exposure to holoendemic malaria results in elevated Epstein-Barr virus loads in children; *J Infect Dis.* 2005. p. 1233-8.2005;191(8):1233-8. Available from http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=15776368
13. Piriou, E.; van Dort, K.; Nanlohy, NM.; van Oers, MH.; Miedema, F.; van Baarle, D. Loss of EBNA1-specific memory CD4+ and CD8+ T cells in HIV-infected patients progressing to AIDS-related non-Hodgkin lymphoma; *Blood.* 2005. p. 3166-74.2005;106(9):3166-74. Available from http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=16014568
14. de-The G. Epstein-Barr virus and Burkitt's lymphoma worldwide: the causal relationship revisited. *IARC Sci Publ.* 1985:165-76. [PubMed: 2998987]
15. Voo, KS.; Fu, T.; Wang, HY.; Tellam, J.; Heslop, HE.; Brenner, MK.; Rooney, CM.; Wang, RF. Evidence for the presentation of major histocompatibility complex class I-restricted Epstein-Barr virus nuclear antigen 1 peptides to CD8+ T lymphocytes; *J Exp Med.* 2004. p. 459-70.142004;199(4):459-70. Available from http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=14769850
15. Fachiroh, J.; Paramita, DK.; Hariwiyanto, B.; Harijadi, A.; Dahlia, HL.; Indrasari, SR.; Kusumo, H.; Zeng, YS.; Schouten, T.; Mubarika, S.; Middeldorp, JM. Single-assay combination of Epstein-Barr Virus (EBV) EBNA1- and viral capsid antigen-p18-derived synthetic peptides for measuring anti-EBV immunoglobulin G (IgG) and IgA antibody levels in sera from nasopharyngeal carcinoma patients: options for field screening; *J Clin Microbiol.* 2006. p. 1459-67. Available from http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=16597877
16. Urban, BC.; Ferguson, DJ.; Pain, A.; Willcox, N.; Plebanski, M.; Austyn, JM.; Roberts, DJ. Plasmodium falciparum-infected erythrocytes modulate the maturation of dendritic cells; *Nature.* 1999. p. 73-7.1999;400(6739):73-7. Available from http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=10403251
17. Cao K, Moormann AM, Lyke KE, Masaberg C, Sumba OP, Doumbo OK, Koech D, Lancaster A, Nelson M, Meyer D, Single R, Hartzman RJ, et al. Differentiation between African populations is evidenced by the diversity of alleles and haplotypes of HLA class I loci. *Tissue Antigens.* 2004; 63:293-325. [PubMed: 15009803]
18. Wilson NS, Behrens GM, Lundie RJ, Smith CM, Waithman J, Young L, Forehan SP, Mount A, Steptoe RJ, Shortman KD, de Koning-Ward TF, Belz GT, et al. Systemic activation of dendritic cells by Toll-like receptor ligands or malaria infection impairs cross-presentation and antiviral immunity. *Nat Immunol.* 2006; 7:165-72. [PubMed: 16415871]
19. Huaman MC, Martin LB, Malkin E, Narum DL, Miller LH, Mahanty S, Long CA. Ex vivo cytokine and memory T cell responses to the 42-kDa fragment of Plasmodium falciparum merozoite surface protein-1 in vaccinated volunteers. *J Immunol.* 2008; 180:1451-61. [PubMed: 18209040]

Figure 1. Comparison of the magnitude of IFN- γ ELISPOT responses in children with eBL and healthy children

The magnitude of IFN- γ ELISPOT responses was compared between children with eBL (n=44) and healthy children from Kisumu (n=60) and Nandi (n=60) for EBNA1 total peptide pool (EATP) and EBNA1 sub-pools, (i.e., EAI, EAIL, EAIII, EAIV, EAV) and for malaria antigen MSP1 (3D7). IFN- γ spot forming units (SFU) minus PBS control SFU per 1×10^6 PBMCs are displayed (range of SFU in PBS wells was 0-12 SFU/ 10^6 PBMCs). Only responses above background are shown. Each dot represents an individual sample. Lines indicate the mean and SEM for each pool tested. There were significantly lower EATP levels in eBL children compared to Nandi ($p < 0.001$) and Kisumu ($p < 0.001$) children, and Kisumu and Nandi children did not differ from each other ($p = 0.562$). Significantly lower responses for children with eBL were also observed for EBNA1 sub-pools EAI, EAIII and EAV whereas there were no differences in responses for EAIL and EAIV. For MSP1, children with eBL had significantly higher IFN- γ responses compared to Kisumu and Nandi children ($p < 0.05$).

Figure 2. Mean EBNA1 IgG1 subclass levels in different populations
(A) Mean EBNA1 IgG1 levels in plasma were determined by ELISA for healthy Kisumu and Nandi children, children with eBL and healthy adults. Scatter plot show individual points for IgG1 titer and SEM is indicated for each group. (B) Frequency of EBNA1 IgG subclass positive samples. The frequency of response is shown for each EBNA1 IgG isotype tested.

Figure 3. Mean EBV viral load stratified by EBNA1-specific IFN- γ ELISPOT response among eBL patients and healthy children with divergent malaria exposure histories

EBV viral load was natural log transformed and then the mean EBV viral load was determined. The mean EBV viral load was compared within each group of children (eBL, Kisumu and Nandi) stratified by EBNA1-specific total pool (EATP) IFN- γ responders and non-responders (indicated as EBNA1 IFN- γ response). A near-significant trend emerged for eBL and Kisumu (but not the Nandi children). EBV viral load tended to be higher in children without IFN- γ responses to EBNA1 in these two groups.

Table 1

Percent IFN- γ responders following stimulation of PBMC with EBNA1 peptide pools (i.e. EATP, EAI, EAIL, EAIIL, EAIIV, EAV), malaria recombinant protein MSP1 (3D7), or EBVpooled peptides (EP1/EP2).

Antigens	eBL [/]			Kisumu			Nandi		
	% Positive (total n=44)	% Positive (total n=60)	Odds Ratio	% Positive (total n=60)	Odds Ratio	p-value	% Positive (total n=60)	Odds Ratio	p-value
EATP	16.1	66.7	10.57			.00	71.7	13.37	.00
EAI	4.5	11.7	2.77			.22	23.7	6.53	.02
EAIL	13.6	30.0	2.71			.06	25.4	2.16	.15
EAIIL	13.6	41.7	4.52			.00	44.1	4.99	.00
EAIIV	6.8	15.0	2.41			.21	20.3	3.49	.07
EAV	4.5	15.0	3.71			.11	22.0	5.94	.02
MSP1 (3D7)	36.4	25.0	.58			.21	13.8	.28	.01
EP1/EP2	43.2	30.0	.56			.17	50.8	1.36	.44

[/] Reference group for comparison to Kisumu and Nandi groups.