Published in final edited form as:

Clin Pharmacol Ther. 2009 January ; 85(1): 36–44. doi:10.1038/clpt.2008.194.

Promoter Polymorphisms in ACE (Angiotensin I–Converting Enzyme) Associated With Clinical Outcomes in Hypertension

AD Johnson^{1,2}, Y Gong^{3,4}, D Wang¹, TY Langaee^{3,4}, J Shin^{3,4}, RM Cooper-DeHoff^{3,4,5}, NJ Schork⁶, P Binkley⁷, CJ Pepine⁵, JA Johnson^{3,4,5}, and W Sadee^{1,7}

¹Program in Pharmacogenomics, Department of Pharmacology, College of Medicine, The Ohio State University, Columbus, Ohio, USA

²The Framingham Heart Study, Framingham, Massachusetts, USA

³Department of Pharmacy Practice, College of Pharmacy, University of Florida, Gainesville, Florida, USA

⁴Center for Pharmacogenomics, University of Florida, Gainesville, Florida, USA

⁵Division of Cardiovascular Medicine, College of Medicine, University of Florida, Gainesville, Florida, USA

⁶Scripps Genomic Medicine, Scripps Health and The Scripps Research Institute, La Jolla, California, USA

⁷Division of Cardiovascular Medicine and the Dorothy Davis Heart and Lung Research Institute, College of Medicine, The Ohio State University, Columbus, Ohio, USA.

Abstract

Genetic variants of *ACE* are suspected risk factors in cardiovascular disease, but the alleles responsible for the variations remain unidentified. To search for regulatory polymorphisms, allelic angiotensin I–converting enzyme (ACE) mRNA expression was measured in 65 heart tissues, followed by genotype scanning of the *ACE* locus. Marked allelic expression imbalance (AEI) detected in five African-American subjects was associated with single-nucleotide polymorphisms (SNPs) (rs7213516, rs7214530, and rs4290) residing in conserved regions 2–3 kb upstream of *ACE*. Moreover, each of the SNPs affected transcription in reporter gene assays. SNPs rs4290 and rs7213516 were tested for associations with adverse cardiovascular outcomes in hypertensive patients with coronary disease (International Verapamil SR Trandolapril Study Genetic Substudy (INVEST-GENES), $n = 1,032$). Both SNPs were associated with adverse cardiovascular outcomes, largely attributable to nonfatal myocardial infarction in African Americans, showing an odds ratio of 6.16 (2.43–15.60) ($P < 0.0001$) for rs7213516. The high allele frequency in African Americans (16%) compared to Hispanics (4%) and Caucasians (<1%) suggests that these alleles contribute to variation between populations in cardiovascular risk and treatment outcomes.

Angiotensin I–converting enzyme (ACE) plays a key role in cardiovascular biology. Its functions are formation of angiotensin II and inactivation of bradykinin, resulting in vasoconstriction and increased blood pressure. ACE inhibitors are recommended as first-line agents in the treatment of hypertension and heart failure. Expressed in many tissues, ACE further affects a broad spectrum of physiological processes. As a result, the *ACE* gene has been

© 2008 American Society for Clinical Pharmacology and Therapeutics

Correspondence: W Sadee (E-mail: wolfgang.sadee@osumc.edu).

SUPPLEMENTARY MATERIAL is linked to the online version of the paper at <http://www.nature.com/cpt>

CONFLICT OF INTEREST

A.D.J., D.W., and W.S. are all coinventors on a submitted patent application pertaining to the *ACE* polymorphisms reported here. The other authors declared no conflict of interest.

implicated in susceptibility to hypertension, myocardial infarction (MI), renal pathophysiology, diabetes, and Alzheimer's disease. The suggestion of a heritable component to serum ACE activity¹ led to extensive phenotype–genotype studies with ACE-related pathophysiology and response to ACE inhibitors.² Numerous studies have focused on an insertion/deletion (I/D) polymorphism in intron 15. However, meta-analyses of phenotypic associations largely failed to confirm a role for I/D,³ and *in vitro* experiments did not reveal any effect on transcription⁴ or splicing.⁵ Therefore, genetic factors contributing to differential ACE expression remain unidentified. As ACE is one of the main therapeutic targets of antihypertensive therapies, a better understanding of genetic differences is required to assess disease risk and response to therapy.

Genetic family studies map the heritable contribution to ACE activity and blood pressure to the region of the ACE gene, particularly in subjects of African ancestry.^{6,7} Moreover, allele frequencies at the ACE locus vary greatly between African Americans and European Americans.⁸ African Americans are at higher risk of hypertension⁹ and its target organ sequelae¹⁰ and less responsive to ACE inhibitors.^{11,12} In addition, African Americans are more likely to experience adverse drug effects.¹³ In this study, we searched for functional polymorphism in the ACE gene locus that might account for differences in disease susceptibility or response to therapy with ACE inhibitors.

The ACE gene consists of 25 exons spanning ~25 kb and encoding a soluble or a membrane-bound protein variant with two peptidase domains (Figure 1). ACE harbors a number of polymorphisms; however, frequent nonsynonymous single-nucleotide polymorphisms (SNPs) that affect the protein sequence are lacking, suggesting that regulatory polymorphisms that are yet to be discovered may contribute to genetic susceptibility in cardiovascular diseases involving ACE. To search for regulatory polymorphisms, we measured allelic mRNA expression of ACE in human cardiac tissues. In contrast to total mRNA levels, allelic mRNA ratios cancel out *trans*-acting factors, so that any detectable allelic expression imbalance (AEI) is a strong indicator of *cis*-acting regulatory factors.^{14–19} Allelic mRNA ratios, as the most proximate and accurate phenotype for SNP scanning, can then be exploited in a search for regulatory polymorphisms, followed by molecular genetic studies to address underlying mechanisms.^{14–16,19}

In this study, we report regulatory alleles affecting ACE expression that are common among African Americans, discovered in a screen of human myocardial tissues. To assess the clinical relevance of these alleles, we conducted a clinical genetic association study in the International Verapamil SR Trandolapril Study Genetic Substudy (INVEST-GENES).²⁰

RESULTS

Allelic mRNA expression of ACE and association with promoter SNPs in heart tissues

We selected two marker SNPs (rs4309 and rs4343)—located in exon 8 and exon 16, respectively, of ACE (Figure 1)—to measure allelic ratios of genomic DNA (gDNA) and mRNA in heart tissues using the SNaPshot method (Applied Biosciences, Foster City, CA). Standard curves performed with mixtures of DNA alleles were linear over the observed range ($r^2 = 0.996–0.999$). As gDNA ratios varied within a small range (less than ± 2 SD), no variable copy number polymorphisms were detectable (although the SNaPshot method used would have missed hemizygous subjects). Therefore, the mean allelic gDNA ratios were normalized to 1. Allelic ACE mRNA expression in heart tissues varied up to fourfold compared to gDNA ratios, indicating the presence of strong *cis*-acting regulatory factors (Figure 2; using a log₂ scale). Allelic expression ratios obtained with the two marker SNPs in compound heterozygotes ($n = 20$) indicate that the results are reproducible. Allelic mRNA ratios deviated significantly from unity in 5 of 33 subjects. Strikingly, each of the five tissues showing strong AEI was obtained

from African-American subjects, even though only 8 African Americans were heterozygous for a marker SNP. In contrast, none of the Caucasians displayed significant AEI (Figure 2), showing a significant difference between ethnic groups.

To ascertain the responsible regulatory polymorphisms, we sequenced the *ACE* locus in gDNA from the eight African-American subjects. No polymorphisms within the transcribed mRNA region (untranslated regions or protein-coding regions) matched the pattern of allelic expression. On the other hand, three polymorphisms (rs7213516, rs7214530, and rs4290) (Table 1) in a region 2–3 kb upstream of the *ACE* transcription start site were strongly associated with allelic mRNA expression imbalance in African Americans ($P < 10^{-7}$). Moreover, these three SNPs were absent in Caucasian cardiac tissues, which also failed to show detectable AEI (Figure 2b). In the cardiac tissues surveyed, rs7213516, rs7214530, and rs4290 were in extensive but incomplete linkage disequilibrium (LD), so we cannot exclude any of the three SNPs from contributing to the AEI ratios. In the HapMap data for the Yuruba population in Ibadan, Nigeria, rs4290 was in complete LD with rs7214530 ($D' = 1.0$, $r^2 = 1.0$) but not with rs7213516 ($D' = 1.0$, $r^2 = 0.55$) (Figure 1). The incomplete LD in HapMap between rs4290 and rs7213516 motivated later selection of these two markers for the clinical association study.

We next genotyped additional *ACE* markers (rs4291, rs4292, rs4357, rs4363, rs13447447, and rs4366) for all samples with allelic mRNA data (Table 1). The only additional SNP showing significant association with AEI, rs4357 ($P < 10^{-7}$) located in intron 21 (Figure 1), was in partial LD with the upstream SNPs (with rs4290: $D' = 1.0$, $r^2 = 0.77$; with rs7213516: $D' = 0.71$, $r^2 = 0.38$). Because several subjects with AEI were homozygous for rs4357, this argues against a functional role. The commonly studied I/D variant (rs13447447) had a P value of 0.09 for association with AEI, again owing to LD with the promoter SNPs, but it was also ruled out because many I/D heterozygotes failed to show AEI.

The value of allelic mRNA ratios <1 in the five African-American subjects showing AEI indicated that the less frequent allele had reduced mRNA expression (considering the inferred phasing between the marker SNP alleles and those of the promoter SNPs). To test this further, we measured overall *ACE* mRNA levels using reverse transcriptase–PCR. Although no association with mRNA levels was observed with the I/D variant (Figure 3a), carrying the minor allele of the promoter SNPs was associated with decreased *ACE* mRNA expression (rs4290 T; $P < 0.02$ (Figure 3b), rs7213516 A; $P < 0.04$). This result indicates that the minor alleles of the promoter SNPs reduce expression.

Reporter gene analysis of three *ACE* promoter SNPs

To determine whether the promoter SNPs rs7213516, rs7214530, and rs4290 affect transcription, we compared the activities of a 4.3-kb fragment from the *ACE* promoter region, which contained either the reference sequence (*G-T-C*) or different combinations of the three SNPs, using a reporter gene assay in HEK293 and bovine aortic endothelial cells (BAEC). As shown in Figure 4a, the expression constructs containing any of the minor alleles of the three promoter SNPs significantly reduced reporter gene expression in BAEC, using two different transfection reagents. Although there were differences in the degree of reduction between the various constructs, no single SNP could account for all results. To test for cell context–dependent effects, we also measured promoter activity in HEK293 cells. In contrast to the results with BAEC, none of the SNPs had an effect on promoter activity in HEK293 cells regardless of plasmid amounts used for transfection (Figure 4b). As the experiments in BAEC and HEK293 cells were performed side by side with the same plasmid preparations, the negative results in HEK293 cells further indicate that the plasmid preparations had similar transfection efficiencies, which can be a source of error if not controlled for. Taken together with the mRNA analysis in heart tissues (Figures 2 and 3), we conclude that each of the three

promoter SNPs appears to reduce *ACE* gene expression, although any effects appear tissue dependent.

Genetic association of *ACE* with adverse cardiovascular outcomes in INVEST-GENES

We genotyped rs7213516, rs4290, and three additional polymorphisms (Table 2) in 258 subjects who experienced a primary outcome event (first occurrence of all-cause death, nonfatal MI, or nonfatal stroke) and 774 hypertensive controls who lacked primary outcome events in the genetic substudy (INVEST-GENES) of the randomized controlled clinical trial INVEST. All genotype frequencies were in Hardy–Weinberg equilibrium in all three race/ethnicity groups and displayed substantial differences between ethnic groups. LD is shown in Figure 5, illustrating the relationships between the geno-typed SNPs. For the five polymorphisms tested in the INVEST-GENES, allele frequencies in Hispanics were intermediate between those of Caucasians and African Americans. Minor allele frequencies of both rs7213516 and rs4290 differed significantly between African Americans (16%), Hispanics (4%), and Caucasians (<1%). Genotyping quality-control checks showed >99.5% concordance between different assays for the same polymorphisms (see also **Supplementary Materials and Methods** online).

Consistent with our finding that the rs7213516 A and rs4290 T alleles are associated with *ACE* expression differences, these alleles were also robustly associated in the INVEST-GENES cohort with increased risk of a primary outcome event (Figure 6). The main effect was strongest in African Americans for both SNPs, with similar trends in Hispanics and Caucasians, despite limited power in these latter groups because of low allele frequency. In African Americans, rs7213516 A and rs4290 T carriers had four times higher odds of experiencing a primary outcome event (odds ratio (OR): 4.13, 95% confidence interval (CI): 1.52–11.21 ($P = 0.0054$), and OR: 3.91, 95% CI: 1.54–9.90 ($P = 0.0041$), respectively). Subgroup analysis by treatment category indicated that the A allele of rs7213516 was associated with higher risk (i.e., OR > 1) in all drug combinations, although this reached statistical significance only in patients receiving both atenolol and trandolapril (OR: 4.31, 95% CI: 1.56–11.89 ($P < 0.005$)) (Supplementary Table S1 online).

In secondary outcomes analysis, rs7213516 conferred the highest risk for nonfatal MI (OR: 6.16, 95% CI: 2.43–15.60 ($P = 0.0001$)), whereas there was no significantly higher risk for all-cause mortality ($P = 0.92$) or nonfatal stroke ($P = 0.19$) (Table 3). Similarly, the association with rs4290 is also largely driven by nonfatal MI (OR 2.37); however, it reached only marginal significance.

The *ACE* I/D polymorphism (rs13447447) was inconsistently associated with outcomes (Figure 6). Associations were not directionally similar in the different racial/ethnic groups, nor was there a linear trend between I/D heterozygotes and I/I homozygotes. Finally, the I/D was not associated with any of the individual components of the composite outcome (Table 3). We therefore conclude that there is no meaningful association with clinical outcomes analyzed here, consistent with a lack of effects on *ACE* mRNA level in heart tissue (Figure 3). There was also no evidence for association of the primary outcome with polymorphisms rs4291 and rs4366.

DISCUSSION

This study used allelic mRNA expression analysis of *ACE* in human heart tissues, followed by SNP scanning, to identify regulatory polymorphisms in the *ACE* locus, long suspected of conferring genetic risk for cardiovascular disease and possibly affecting treatment outcomes. This approach revealed strong effects on *ACE* mRNA expression attributable to three promoter SNPs—rs7213516, rs7214530, and rs4290—located in conserved regions 2–3 kb upstream of the transcription start site. The excellent congruence between AEI ratios and clearly identifiable

polymorphisms, and a significant association between genotypes and total ACE mRNA expression in human heart tissues, support the notion that these promoter SNPs reduce ACE mRNA expression. This conclusion is further buttressed by results from reporter gene assays. The three *ACE* promoter SNPs are common in individuals of African-American ancestry (Table 1) but rare in Caucasians and intermediate in Hispanics. As the promoter SNP effects on ACE mRNA expression were observed only in tissues from heart failure patients, we cannot exclude the possibility that AEI is a result of the disease state. However, consistent with our gene expression results, a clinical association study in a non-heart failure group of hypertensive patients revealed a robust genetic effect on outcomes, suggesting that the heart failure state was not an important determinant of the findings.

Three promoter SNPs linked to ACE expression

Allelic mRNA expression ratios were strongly linked with the three promoter SNPs ($P \leq 0.0001$), but because of the extensive LD among them, a conclusion cannot be drawn regarding which polymorphism is functional. All three SNPs reside in conserved regions (Figure 7). Moreover, rs7214530 is part of a predicted recognition site for MEF2, a cardiac transcription factor previously implicated in cardiovascular disease and MI.^{21–23} It is therefore possible that all three SNPs have co-evolved as part of a haplotype block prevalent in subjects of African origin, each contributing to gene regulation, possibly to different extents in different tissues. Reporter gene assays with an *ACE* promoter fragment containing various combinations of the three suspected SNPs demonstrated decreased promoter activity for each combination of variant alleles, compared to the reference sequence in endothelial cells (BAEC) but not in HEK293 cells, indicating that these effects can be tissue specific. It is therefore likely that these *ACE* promoter polymorphisms have different effects in different target tissues and therefore could be associated with different pathophysiologies. The genetic effects on expression observed here in cardiac ventricular tissues and endothelial cells are interesting because prior evidence from animal models supports tissue-specific effects of ACE. ACE is expressed in endothelial cells as a membrane-bound enzyme on the luminal surface of the vasculature and cleaved to derive an active, soluble form in the plasma. The specific regulation of ACE in endothelial cells plays a key role in blood pressure regulation.²⁴ Furthermore, animal models of ACE overexpression in cardiac tissues and experiments in cell lines suggest that cardiac-specific gene expression of ACE may contribute to cardiac hypertrophy.²⁵ Together these results suggest that further experiments to elucidate the mechanisms and contribution of functional alleles in specific tissues may provide valuable insights with clinical relevance.

Other polymorphisms in ACE

We found no evidence for a functional effect of the *ACE* I/D polymorphism in intron 15 on mRNA expression in human heart tissue, consistent with previous negative *in vitro* studies.^{4, 5} In addition, our clinical association data did not support an effect of I/D on outcomes across the various ethnic populations, despite an allele frequency and power that was substantially higher than for the promoter SNPs. Yet countless genetic association studies are based on the I/D polymorphism, even though evidence for a physiological function is lacking and clinical associations are borderline at best.³

Association of ACE promoter SNPs with clinical outcomes

We tested several *ACE* polymorphisms for association with clinical outcomes in hypertensive patients with coronary artery disease (INVEST-GENES). The promoter SNPs identified in our mechanism-based screen (rs7213516 and rs4290; rs7214530 was not genotyped because of strong LD with rs4290) were highly associated with cardiovascular disease outcomes ($P < 0.001$) (Figure 6) and in particular with MI (Table 4). The ORs ranging from 4 to 6 suggest an unexpectedly strong genetic effect, which must be further assessed in replication studies.

We also assessed relative risk of primary outcome as a function of drug treatment, showing association with the *ACE* promoter SNPs in individuals receiving ACE inhibitor and/or β -blocker therapy (**Supplementary Materials and Methods** online). However, the INVEST-GENES design was not optimal for assessing the effects of genetic factors on drug treatment outcomes, and therefore these results can be taken only as a guide for future experiments. Nevertheless, this association may have a biological basis, given that trandolopril and atenolol target overlapping systems of blood pressure control, of which ACE is a critical component. Results from the Valsartan Heart Failure (Val-HeFT) trial raise the possibility that excessive neurohormonal inhibition may contribute to adverse outcomes in heart failure treatment.²⁶ Because the promoter alleles identified here are associated with decreased ACE expression, we hypothesize that they may potentiate pharmacological ACE inhibition plus β -blockade, resulting in higher event rates via excessive neurohormonal inhibition. A prospective study of the *ACE* promoter SNP effect on drug treatment outcomes is required to resolve these issues.

Given that the promoter alleles are common in African Americans, they may partially account for phenotypic variation in ACE levels, blood pressure (e.g., ref. ⁶), and response to ACE inhibitors^{11–13} in individuals of African ancestry. Although these alleles were found at lower frequencies in Hispanics and Caucasians, they could be clinically relevant in the population at large, but we had limited statistical power to address this question. Further investigation will determine whether these alleles may have clinical utility as biomarkers in the selection of therapeutic options for individual patients. The use of race in guiding treatment is controversial but does play a role in clinical practice.²⁷ Ultimately, therapy may be best optimized for individual patients with tests for functional biomarkers instead of relying on assumptions related to apparent, or self-identified, race or ethnicity.²⁸

Physiological roles for ACE include blood pressure regulation, kidney function, processing of kinins and other peptides, and degradation of amyloid β -protein,^{29,30} suggesting that the new *ACE* promoter alleles may be relevant in other human pathologies. Among the heart tissues obtained from 12 African-American heart transplant patients, only 8 were eligible for AEI analysis. Among the 12 samples, the minor allele frequencies of rs7213516 and rs4290 (25–27%) were higher than expected (16.0% in INVEST-GENES), with one patient homozygous for the minor alleles, arguing for a larger study of heart failure patients.

In conclusion, the discovery of regulatory alleles in key genes through allelic mRNA expression analysis, followed by clinical association studies, has broad potential to lead to viable biomarkers guiding an individual's therapy.³¹

METHODS

Analysis of ACE mRNA expression in heart tissues

Approval for use of human subjects was obtained from the Ohio State University Institutional Review Board. Left-ventricle tissue from 65 heart transplant patients was obtained through the Cooperative Human Tissue Network, Midwestern Division, at Ohio State University and stored at -80°C until extraction. gDNA and RNA were isolated, and cDNA was prepared from 1.0 μg RNA in three independent preparations, using oligo dT and gene-specific primers close to the two marker SNPs to minimize the effects of mRNA decay in postextract tissues (see **Supplementary Materials and Methods** online for details).

Total mRNA expression levels

Overall ACE mRNA expression was measured using reverse transcriptase–PCR for each sample (details in **Supplementary Materials and Methods** online). Gene expression results by genotype were analyzed by *t*-test for mean differences using SPSS 14 (Chicago, IL).

Measurement of allelic ACE mRNA expression

We measured allelic mRNA expression, as described previously,^{14–19} amplifying short regions of gDNA and cDNA around *ACE* exonic marker SNPs from heart tissues of heterozygous individuals (rs4309, located in exon 8, $n = 28$; rs4343, located in exon 16, $n = 24$). Primer extension with fluorescent dideoxy-nucleotides by SNaPshot (Applied Biosciences, Foster City, CA) allowed quantitation of relative amounts of each allele by capillary electrophoresis on an ABI 3730 DNA Analyzer (Applied Biosciences, Foster City, CA). Corrected allelic mRNA expression ratios for individual cDNAs were calculated by normalizing to the mean ratio of gDNA peaks (SD for gDNA: rs4309 $\pm 12.4\%$, rs4343 $\pm 8.6\%$). Each sample was assayed from three independent cDNA syntheses, each performed at least in duplicate (details in **Supplementary Materials and Methods** online).

Scanning the ACE locus for functional polymorphisms

To link SNPs with allelic mRNA expression ratios, we genotyped SNPs selected to represent the major haplotype blocks (Table 1) in all 65 heart tissues. SNPs were genotyped as described in **Supplementary Materials and Methods** online. In addition, we sequenced full-length cDNAs and the 5'-upstream region over 3 kb in eight African Americans, detecting five SNPs in the upstream region (Table 1). The presence of AEI was set at allelic mRNA ratios >1.5 or $<1/1.5$ as a cutoff. Association between genotype status (heterozygous or homozygous) and AEI was determined using HelixTree (Golden Helix, Bozeman, MT). LD between SNPs (expressed as D') and haplotypes was calculated using HelixTree (Golden Helix).

ACE reporter gene assay

A promoter fragment ranging from $-4,335$ to $+1$ (the major transcription start site) in PGL3.basic vector (Promega, Madison, WI) was provided by Melanie Eyries.³² Various combinations of rs7213516/rs7214530/rs4290 haplotypes were obtained using site-directed mutagenesis or restriction digest of amplified gDNA with *MscI* and *BstEII* and subsequent cloning (see **Supplementary Materials and Methods** online for details). All inserts were fully sequenced to verify the intended sequence. The constructs were transfected into HEK293 and BAEC, cultured in Dulbecco's modified Eagle's medium/F12 media containing 10% fetal bovine serum, penicillin (10 U/ml), and streptomycin (10 μ g/ml), at 37°C with 5% CO₂. Twenty-four hours before transfection, 1×10^5 to 2×10^5 cells were plated into 24-well plates and transiently transfected with FuGENE HD Transfection Reagent (Roche Applied Science, Indianapolis, IN) or Lipofectamine (Invitrogen, Carlsbad, CA) in serum-free medium for 5 h. As a control, *Renilla* luciferase constructs were cotransfected with PGL3.basic-fused constructs at a 1:20 ratio. Cells were harvested after 48 h and transferred to 96-well plates, and luciferase activity was detected using Dual-Glo luciferase assays (Promega, Madison, WI) on a fluorescence plate reader (PerkinElmer, Waltham, MA). Two independent transfections and triplicate luciferase assays were performed for each construct and cell line. Results were analyzed using Prism (GraphPad, La Jolla, CA).

Clinical genetic association study: INVEST and INVEST-GENES

The INVEST trial evaluated adverse cardiovascular outcomes in patients randomized to atenolol or verapamil SR hypertension treatment strategy in 22,576 patients with documented coronary artery disease and hyper-tension²⁰ (details in **Supplementary Materials and Methods** online). The primary outcome was the first occurrence of death (all cause), nonfatal MI, or nonfatal stroke. These events were taken separately as secondary outcomes. In the genetic substudy (INVEST-GENES), gDNA was collected from 5,979 patients using buccal cells from mouthwash samples.³³ All patients provided written informed consent, as approved by the University of Florida Institutional Review Board. The present case-control study focused on the 258 INVEST-GENES patients who experienced a primary outcome event

during study follow-up (cases), frequency matched 3:1 for age, sex, and race/ethnicity with 774 individuals who were event-free during study follow-up (controls) (baseline characteristics in Table 4). The patients had a mean age of 71 years, 50% were female, 25% were of Hispanic ethnicity, and, 13% were African Americans. Previous analyses showed that case-control analysis in this group matched findings from the entire INVEST cohort, the inclusion of which increases only the number of controls.³⁴

We genotyped promoter SNPs rs7213516 and rs4290 and the tagging markers (rs4291, rs13447447, and rs4366) (genotyping details in **Supplementary Materials and Methods** online) to sample major haplotype blocks. Quality-control procedures included blind duplicate genotyping of 5% of samples using the same or an alternative method, assessment of Hardy-Weinberg equilibrium, and assay validation using Coriell samples previously genotyped as part of HapMap. To address potential population stratification, we genotyped 87 autosomal ancestry informative markers interspaced with large interlocus distances across the genome to give independent association with the disease and genetic background (see **Supplementary Materials and Methods** online).^{35,36}

Statistical analysis of clinical genetic associations

Baseline characteristics between cases and controls in INVEST-GENES were compared using *t*-test for continuous variables and χ^2 -test for categorical variables. Hardy-Weinberg equilibrium of genotype frequencies within each racial/ethnic group was tested using χ^2 -test with one degree of freedom. Because of the low minor allele frequencies for rs7213516 and rs4290 in the entire INVEST cohort, we decided *a priori* to combine heterozygous patients with those homozygous for the variant alleles for all analyses. Logistic regression was performed to assess the association of genotypes/haplotypes with the primary and secondary outcomes after adjusting for ancestry and prespecified confounding factors, namely age (by decades), gender, race/ethnicity, history of MI and heart failure, and drug treatments.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

ACKNOWLEDGMENTS

This study was supported by National Institutes of Health grants HL 74730, HL 69758, and RR017568, grants from Abbott Laboratories and the University of Florida Opportunity Fund, and a Biomedical Research and Technology Transfer Partnership Program grant from the State of Ohio. A.D.J. was initially supported by an American Heart Association Pre-Doctoral Fellowship (0515157B) and completed the work at Ohio State University while being supported by an Ohio State University Distinguished University Fellowship. J.S. was an American Heart Association Post-Doctoral Fellow at the time of the work. We thank M. Eyries for kindly providing the ACE reference upstream construct.

References

1. Cambien F, et al. Familial resemblance of plasma angiotensin-converting enzyme level: the Nancy study. *Am. J. Hum. Genet* 1988;43:774–780. [PubMed: 2847529]
2. Arnett DK, et al. Pharmacogenetic association of the angiotensin-converting enzyme insertion/deletion polymorphism on blood pressure and cardiovascular risk in relation to antihypertensive treatment: the Genetics of Hypertension-Associated Treatment (GenHAT) study. *Circulation* 2005;111:3374–3383. [PubMed: 15967849]
3. Sayed-Tabatabaei FA, Oostra BA, Isaacs A, van Duijn CM, Witteman JC. ACE polymorphism. *Circ. Res* 2006;98:1123–1133. [PubMed: 16690893]
4. Rosatto N, Pontremoli R, de Ferrari G, Ravazzolo R. Intron 16 insertion of the angiotensin converting enzyme gene and transcriptional regulation. *Nephrol. Dial. Transplant* 1999;14:868–871. [PubMed: 10328461]

5. Lei H, Day IN, Vorechocský I. Exonization of *Alu* Ya5 in the human *ACE* gene requires mutations in both 3' and 5' splice sites and is facilitated by a conserved splicing enhancer. *Nucleic Acids Res* 2005;33:3897–3906. [PubMed: 16027113]
6. Zhu X, et al. Linkage and association analysis of angiotensin I-converting enzyme (ACE)-gene polymorphisms with ACE concentration and blood pressure. *Am. J. Hum. Genet* 2001;68:1139–1148. [PubMed: 11283791]
7. Bouzekri N, et al. Angiotensin I-converting enzyme polymorphisms, ACE level and blood pressure among Nigerians, Jamaicans and African-Americans. *Eur. J. Hum. Gen* 2004;12:460–468.
8. Rieder MJ, Taylor SL, Clark AG, Nickerson DA. Sequence variation in the human angiotensin converting enzyme. *Nat. Genet* 1999;22:59–62. [PubMed: 10319862]
9. Hajjar I, Kotchen TA. Trends in prevalence, awareness, treatment, and control of hypertension in the United States, 1998–2000. *JAMA* 2003;290:199–206. [PubMed: 12851274]
10. Rosamond W, et al. Heart disease and stroke statistics—2007 update: a report from the American Heart Association statistics committee and stroke statistics subcommittee. *Circulation* 2007;115:e69–e171. [PubMed: 17194875]
11. Exner DV, Dries DL, Domanski MJ, Cohn JN. Lesser response to angiotensin-converting-enzyme inhibitor therapy in black as compared with white patients with left ventricular dysfunction. *N. Engl. J. Med* 2001;344:1351–1357. [PubMed: 11333991]
12. Sehgal AR. Overlap between whites and blacks in response to antihypertensive drugs. *Hypertension* 2004;43:566–572. [PubMed: 14757779]
13. McDowell SE, Coleman JJ, Ferner RE. Systematic review and meta-analysis of ethnic differences in risks of adverse reactions to drugs used in cardiovascular medicine. *BMJ* 2006;332:1177–1181. [PubMed: 16679330]
14. Johnson AD, et al. Polymorphisms affecting gene transcription and mRNA processing in pharmacogenetic candidate genes: detection through allelic expression imbalance in human target tissues. *Pharmacogenet. Genomics* 2008;18:781–791. [PubMed: 18698231]
15. Wang D, Johnson AD, Papp AC, Kroetz DL, Sadée W. Multidrug resistance polypeptide 1 (*MDR1*, *ABCB1*) variant 3435C>T affects mRNA stability. *Pharmacogenet. Genomics* 2005;15:693–704. [PubMed: 16141795]
16. Zhang Y, Wang D, Johnson AD, Papp AC, Sadée W. Allelic expression imbalance of human mu opioid receptor (*OPRM1*) caused by variant A118G. *J. Biol. Chem* 2005;280:32618–32624. [PubMed: 16046395]
17. Lim JE, Pinsonneault J, Sadée W, Saffen D. Tryptophan hydroxylase 2 (TPH2) haplotypes predict levels of TPH2 mRNA expression in human pons. *Mol. Psychiatry* 2007;12:491–501. [PubMed: 17453063]
18. Pinsonneault JK, Papp AC, Sadée W. Allelic mRNA expression of X-linked monoamine oxidase A (MAOA) in human brain: dissection of epigenetic and genetic factors. *Hum. Mol. Genet* 2006;15:2636–2649. [PubMed: 16893905]
19. Zhang Y, et al. Novel polymorphisms in the human dopamine D2 receptor gene affect gene expression, splicing, and neuronal activity. *Proc. Natl. Acad. Sci. USA* 2007;104:20552–20557. [PubMed: 18077373]
20. Pepine CJ, et al. A calcium antagonist versus a non-calcium antagonist hypertension treatment strategy for patients with coronary heart disease. The International Verapamil-Trandolopril Study (INVEST): a randomized control trial. *JAMA* 2003;290:2805–2816. [PubMed: 14657064]
21. Naya FJ, et al. Mitochondrial deficiency and cardiac sudden death in mice lacking the MEF2A transcription factor. *Nat. Med* 2002;8:1303–1309. [PubMed: 12379849]
22. Wang L, Fan C, Topol SE, Topol EJ, Wang Q. Mutation of MEF2A in an inherited disorder with features of coronary artery disease. *Science* 2003;302:1578–1581. [PubMed: 14645853]
23. van Oort RJ, et al. MEF2 activates a genetic program promoting chamber dilation and contractile dysfunction in calcineurin-induced heart failure. *Circulation* 2006;114:298–308. [PubMed: 16847152]
24. Kessler SP, deS Senanayake P, Scheidemantel TS, Gomos JB, Rowe TM, Sen GC. Maintenance of normal blood pressure and renal functions are independent effects of angiotensin-converting enzyme. *J. Biol. Chem* 2003;278:21105–21112. [PubMed: 12777443]

25. Tian XL, et al. Over-expression of angiotensin converting enzyme-1 augments cardiac hypertrophy in transgenic rats. *Hum. Mol. Genet* 2004;13:1441–1450. [PubMed: 15128700]
26. Cohn JN, Tognoni G, Valsartan Heart Failure Trial Investigators. A randomized trial of the angiotensin-receptor blocker valsartan in chronic heart failure. *N. Engl. J. Med* 2001;345:1667–1675. [PubMed: 11759645]
27. Taylor AL, et al. Combination of isosorbide dinitrate and hydralazine in blacks with heart failure. *N. Engl. J. Med* 2004;351:2049–2057. [PubMed: 15533851][Erratum, *N. Engl. J. Med.* **352**, 1276 (2005).]
28. Bamshed M. Genetic influences on health: does race matter? *JAMA* 2005;294:937–946. [PubMed: 16118384][Erratum, *JAMA* **294**, 1620 (2005).]
29. Hemming ML, Selkoe DJ. Amyloid beta-protein is degraded by cellular angiotensin-converting enzyme (ACE) and elevated by an ACE inhibitor. *J. Biol. Chem* 2005;280:37644–37650. [PubMed: 16154999]
30. Meng Y, et al. Association of polymorphisms in the angiotensin-converting enzyme gene with Alzheimer disease in an Israeli Arab community. *Am. J. Hum. Genet* 2006;78:871–877. [PubMed: 16642441]
31. Johnson AD, Wang D, Sadée W. Polymorphisms affecting gene regulation and mRNA processing: broad implications for pharmacogenetics. *Pharmacol. Ther* 2005;106:19–38. [PubMed: 15781120]
32. Eyries M, Agrapart M, Alonso A, Soubrier F. Phorbol ester induction of angiotensin-converting enzyme transcription is mediated by Egr-1 and AP-1 in human endothelial cells via ERK1/2 pathway. *Circ. Res* 2002;91:899–906. [PubMed: 12433834]
33. Andrisin TE, Humma LM, Johnson JA. Collection of genomic DNA by the noninvasive mouthwash method for use in pharmacogenetic studies. *Pharmacotherapy* 2002;22:954–960. [PubMed: 12173798]
34. Beitelshes AL, et al. *KCNMB1* genotype influences response to verapamil and adverse outcomes in the International Verapamil SR/Trandolapril Study (INVEST). *Pharmacogenet. Genomics* 2007;17:719–729. [PubMed: 17700361]
35. Shriver MD, et al. Large-scale SNP analysis reveals clustered and continuous patterns of human genetic variation. *Hum. Genomics* 2005;2:81–89. [PubMed: 16004724]
36. Myakishev MV, Khripin Y, Hu S, Hamer DH. High-throughput SNP genotyping by allele-specific PCR with universal energy-transfer-labeled primers. *Genome Res* 2001;11:163–169. [PubMed: 11156625]

Figure 1.

ACE gene structure (University of California–Santa Cruz genome browser) and location of polymorphisms tested in this study. Overviews of HapMap linkage disequilibrium (LD) in the gene region for individuals from Utah of northern-European ancestry (CEU) and individuals from Yoruba, Nigeria (YRI), are shown at bottom (Haploview). ACE, angiotensin I–converting enzyme.

Figure 2.

ACE allelic mRNA expression in left ventricular heart tissues from (a) African Americans and (b) Caucasians. Allelic mRNA expression ratios (major/minor allele for marker single-nucleotide polymorphisms (SNPs) rs4309 (C/T), rs4343 (A/G)) are averages of results using both markers. Allelic expression imbalance was prevalent in (a) African-American but not (b) Caucasian heart tissues. Genotypes for the promoter SNPs are indicated above the African-American samples. Data are mean $\pm$ SD, *** P < 0.001 vs. pooled DNA ratios. ACE, angiotensin I-converting enzyme.

Figure 3. Total mRNA expression levels of angiotensin I-converting enzyme (ACE) in 65 heart tissues. Box plots display median expression $\pm$ one quartile. Results are grouped by (a) genotype of the insertion/deletion (I/D) ($P = 0.93$) and (b) carriers of the promoter rs4290 T allele. ACE mRNA levels are relative to β -actin. * $P < 0.05$ vs. CC genotype.

Figure 4.

Luciferase reporter gene assay of the *ACE* promoter in bovine aortic endothelial cells (BAEC) (a) and HEK293 cells (b). An *ACE* promoter fragment from -4,335 to +1 was cloned into pGL3.basic vector containing the promoter single-nucleotide polymorphisms (SNPs) rs7213516 (G/A), rs7214530 (T/G), and rs4290 (C/T). The reference haplotype was G-T-C, whereas the variant constructs contained one to three minor alleles. In BAEC, 0.8 µg plasmids were cotransfected with 40 ng *Renilla* luciferase plasmid using either Lipofectamine or FuGENE reagent. Luciferase activities from fused-pGL3 vector were normalized using *Renilla* luciferase activity as an internal control. * $P < 0.05$; ** $P < 0.001$ compared to reference haplotype G-T-C. In HEK293 cells, various amounts of plasmid were transfected using Lipofectamine, with no differences observed between any conditions. ACE, angiotensin I-converting enzyme.

Figure 5. Linkage disequilibrium (LD) structure of polymorphisms in the International Verapamil SR Trandolapril Study Genetic Substudy clinical genetic association study. Values for D' and r^2 are shaded on a gradient based on the strength of correlation. Boxes without number values and shaded dark gray indicate perfect correlation. Boxes without number values and shaded light gray indicate that low allele frequency prevented calculation of D' .

Figure 6.

Odds ratios of three polymorphisms for primary outcome in the overall population and within each group. Odds ratios were adjusted for age, sex, race/ethnicity, body mass index, smoking, International Verapamil SR Trandolapril Study treatment strategy, previous myocardial infarction, previous stroke, heart failure, diabetes, renal insufficiency, baseline systolic blood pressure, diuretic use, and angiotensin I-converting enzyme inhibitor use. AA, African American; Cauc, Caucasian; Hisp, Hispanic.

Figure 7.

Promoter sequence alignments and transcription factor-binding sites. The three promoter single-nucleotide polymorphisms rs7213516, rs7214530, and rs4290 are located –2883, –2828, and –2306 bp upstream of the transcription start site (+1). Predicted MEF2A-binding sites (JASPAR position-weight matrices) are shown. Sequence alignments (made with CLUSTALW) are based on Basic Local Alignment Search Tool of the human promoter region. *1-bp insert in rhesus, dog, elephant, and armadillo; †9-bp insert in dog and armadillo.

Table 1

Polymorphisms analyzed in this study and minor allele frequencies observed in heart tissues, sorted by race/ethnicity

	SNP location	Minor allele frequencies, human heart tissues	
		Caucasians (%)	African Americans (%)
rs7213516 (G>A)	−2883 from TSS	0.0	27.3 (<i>n</i> = 12)
rs7214530 (T>G)	−2828 from TSS	n.d.	29.0 (<i>n</i> = 12)
rs4290 (C>T)	−2305 from TSS	0.0	25.0 (<i>n</i> = 12)
rs4291 (A>T)	−240 from TSS	34.2	31.3 (<i>n</i> = 8)
rs4292 (T>C)	−93 from TSS	31.6	6.3 (<i>n</i> = 8)
rs4309 (C>T) AEI marker SNP	Exon 8 (Pro>Pro)	44.4	37.5 (<i>n</i> = 12)
rs13447447 (I/D, insertion/deletion)	Intron15	49.1 ^b	45.5 (<i>n</i> = 11)
rs4343 (A>G) AEI marker SNP	Exon 16 (Thr>Thr)	48.1 ^a	20.8 (<i>n</i> = 12)
rs4357 (C>T)	Intron 21	0.0	22.7 (<i>n</i> = 11)
rs4363 (A>G)	Intron 24	47.2	36.4 (<i>n</i> = 11)
rs4366 (22/33)	3' Downstream	49.1	27.3 (<i>n</i> = 11) ^a

AEI, allelic expression imbalance; n.d., not detected; SNP, single-nucleotide polymorphism.

^ars13447447 (I) and rs4343 (A) were the minor alleles in the Caucasian heart tissue samples.

^b22 is the minor allele in Hispanics and African Americans; 33 is the minor allele in Caucasians.

Table 2
Polymorphisms analyzed in this study and minor allele frequencies observed in the INVEST-GENES cohort, sorted by race/ethnicity

	Minor allele	overall (%)	Minor allele frequencies INVEST-GENES			P value ^a
			Caucasians (%)	Hispanics (%)	African Americans (%)	
rs7213516 (G>A)	A	3.37	0.17	4.33	16.00	<0.0001
rs4290 (C>T)	T	3.60	0.57	4.21	16.03	<0.0001
rs4291 (A>T)	T	37.12	38.78	34.68	32.92	0.11
rs13447447 (U/D)	I	41.16	41.67	39.62	40.57	0.38
rs4366 (22/33)	22 ^b	47.91	55.89	49.79	40.16	0.0004

The P values indicate the level of significance for interethnic differences in minor allele frequencies. INVEST-GENES, International Verapamil SR Trandolapril Study Genetic Substudy.

^a P value for χ^2 -tests or Fisher's exact test for the genotype frequencies by race/ethnicity, as appropriate.

^b 22 is the minor allele in Hispanics and African Americans; 33 is the minor allele in Caucasians.

Table 3
Unadjusted and adjusted odds ratios (ORs) and 95% confidence intervals (CIs) for secondary outcomes by genotype

	Unadjusted OR (95% CI)	Adjusted OR ^a (95% CI)
All-cause death (<i>n</i> = 90)		
rs7213516 (A carriers vs. GG)	1.04 (0.44–2.49)	1.68 (0.62–4.54)
rs4290 (T carriers vs. CC)	1.41 (0.68–2.94)	1.92 (0.83–4.46)
rs13447447 (ID vs. DD)	1.49 (0.87–2.55)	1.45 (0.83–2.52)
rs13447447 (II vs. DD)	1.34 (0.65–2.77)	1.34 (0.63–2.82)
Nonfatal MI (<i>n</i> = 75)		
rs7213516 (A carriers vs. GG)	2.81 (1.40–5.64) [*]	6.16 (2.43–15.60) ^{**}
rs4290 (T carriers vs. CC)	1.78 (0.85–3.73)	2.37 (0.99–5.69)
rs13447447 (ID vs. DD)	0.59 (0.34–1.02)	0.61 (0.34–1.07)
rs13447447 (II vs. DD)	1.41 (0.76–2.65)	1.44 (0.75–2.79)
Nonfatal stroke (<i>n</i> = 81)		
rs7213516 (A carriers vs. GG)	1.68 (0.77–3.65)	1.72 (0.70–4.24)
rs4290 (T carriers vs. CC)	1.43 (0.66–3.09)	1.27 (0.54–3.02)
rs13447447 (ID vs. DD)	0.69 (0.40–1.20)	0.66 (0.37–1.16)
rs13447447 (II vs. DD)	1.34 (0.70–2.59)	1.41 (0.72–2.76)

MI, myocardial infarction.

^a Adjusted for: age, sex, race/ethnicity, body mass index, smoking, previous myocardial infarction, heart failure, diabetes, and ancestry.

^{*} *P* = 0.004

^{**} *P* = 0.0001.

Table 4

Baseline characteristics for INVEST-GENES case and control patients

Characteristic (N, % unless otherwise noted)	Cases (N = 258)	Controls (N = 774)
Age, mean (SD), years	71.5 (9.9)	70.2 (9.3)
Women	131 (50.8)	393 (50.8)
BP, mean (SD), mm Hg		
Systolic	150.6 (19.0)	147.4 (19.0)
Diastolic	83.6 (11.1)	83.3 (11.1)
Race/ethnicity		
White	158 (61.2)	472 (61.0)
Black	36 (14.0)	101 (13.1)
Hispanic	63 (24.4)	198 (25.6)
Other/multiracial	1 (0.4)	3 (0.4)
BMI, mean (SD), kg/m ²	27.4 (4.8)	29.0 (5.5)
Medical history		
Prior myocardial infarction	96 (37.2)	230 (29.7)
Angina pectoris	153 (59.3)	483 (62.4)
Prior stroke/TIA	36 (14.0)	71 (9.2)
Left ventricular hypertrophy	46 (17.8)	136 (17.6)
Heart failure (classes I–III)	28 (10.9)	29 (3.8)
Peripheral vascular disease	43 (16.7)	88 (11.4)
Smoking		
Past	133 (51.6)	355 (45.9)
Within 30 days	34 (13.2)	83 (10.7)
Diabetes ^a	102 (39.5)	224 (28.9)
Hypercholesterolemia ^a	161 (62.4)	485 (62.7)
Renal impairment ^b	14 (5.4)	18 (2.3)
Cancer	20 (7.8)	46 (5.9)
Medication at enrollment		
Aspirin/other antiplatelet agent	162 (62.8)	451 (58.3)
Antidiabetic medication	86 (33.3)	188 (24.3)
Any lipid-lowering agent	106 (41.1)	331 (42.8)
Nitrates	92 (35.7)	232 (30.0)

Percentages may not add up to 100 due to rounding.

BMI, body mass index; BP, blood pressure; INVEST-GENES, International Verapamil SR Trandolapril Study Genetic Substudy; TIA, transient ischemic attack.

^aHistory of or currently taking antidiabetic or lipid-lowering medications.^bHistory of or current serum creatinine level that is elevated but <4 mg/dl (<354 μmol/l).