Published in final edited form as:

J Comb Chem. 2008 ; 10(5): 721–725. doi:10.1021/cc800078h.

Explorations of *Stemona* Alkaloid-Inspired Analogues: Skeletal Modification and Functional Group Diversification

Kevin J. Frankowski, Benjamin Neuenswander, and Jeffrey Aubé*

Department of Medicinal Chemistry and Center for Chemical Methodologies and Library Development, University of Kansas, Malott Hall, 1251 Wescoe Hall Drive, Lawrence, Kansas 66045-7582

Abstract

A tandem Diels-Alder/Schmidt reaction provided an efficient route for the exploration of unnatural *Stemona* alkaloid analogues. Thus, a series of tricyclic scaffolds were efficiently prepared and then elaborated into seven sets of functionalized analogues. These derivatives incorporated appended heterocycles such as indoles and quinolines or other diversity-incorporating moieties such as carbamates and amines. Both the scaffold generation sequence and the diversification steps could be manipulated to provide regio- and diastereochemically-pure products.

Keywords

Diels–Alder; Schmidt reaction; Fischer indole synthesis; Friedländer quinoline synthesis; reductive amination; carbamate formation; azides; *Stemona* alkaloid analogues

There is considerable interest in the synthesis of structurally diverse analogues of biologically active natural products.¹ *Stemona* alkaloid congeners should provide an excellent focus for such an approach due to their extensive history in Chinese traditional medicine. Diverse activities such as antitussive, antiparasitic, and the ability to treat ocular disease have been ascribed to various members of this extensive class.² In one recent study, the antitussive activities of several individual alkaloids were verified in an animal model.³ However, the systematic preparation of *Stemona* alkaloid analogues is challenging due to the structural complexity of these alkaloids.

The tandem Diels–Alder/azido Schmidt reaction has proved useful for the rapid construction of multicyclic scaffolds containing appropriate functional groups for further elaboration.⁴ In particular, this reaction sequence was used in the target-directed synthesis of the *Stemona* alkaloids stenine and neostenine.⁵ In this work, we report an efficient approach to variations of the *Stemona* alkaloid core and demonstrate the diversification of these compounds into a series of previously unreported *Stemona* alkaloid chemotypes.

E-mail: jaube@ku.edu.

Supporting Information **Available** Experimental details and full characterization data (including ¹H and ¹³C NMR spectra) for all new individual compounds synthesized and representative members of carbamate library compounds, HPLC purity data for all library compounds and X-ray crystallographic data (CIF file) for **9**, **10/1b** and **13/3**. This material is available free of charge via the internet at <http://www.acs.pubs.org>.

Scaffold synthesis

We began by preparing scaffolds based on the potent antitussive agents^{3a} neostenine **1** and neotuberostemonine **2** in which the lactone ring(s) was eliminated (Figure 1). A tandem Diels–Alder/azido Schmidt reaction using siloxy diene **3** converted simple cycloalkenone precursors into the 7,6,5 nitrogenous core of the natural products as well as two 6,6,5 analogues (Scheme 1). The relative stereochemistry at the ring junction of each scaffold arises from an *endo* selective Diels–Alder step coupled with retention of stereochemistry in the ring expansion step. Previous studies had established that $\text{BF}_3 \cdot \text{OEt}_2$ was necessary to obtain clean *endo* selectivity in trisubstituted dienes (i.e., **3** where R = alkyl).^{5b} In most other cases, the more reactive SnCl_4 could be used. For example, treating **3** (R = H) and cyclohexanone with SnCl_4 , a mixture of *endo* and *exo* tricycles were obtained in 66% total yield (the major *endo* isomer could be isolated by recrystallization). Alternatively, $\text{BF}_3 \cdot \text{OEt}_2$ conditions provided the *endo* isomer exclusively in 27% yield. In all examples, only the *endo* isomers were purified for use in subsequent steps. Where appropriate, α -substituted ketones were treated with base to epimerize the alkyl group into the thermodynamically favored β orientation, corresponding to that observed in the natural series. Scheme 1 displays the full range of scaffolds produced; 0.5–2.0 g quantities of each were routinely obtained. Importantly, these scaffolds contained a ketone useful for diversification; providing opportunities for modification not readily available in the natural products themselves.

Diversification studies: Introduction of basic nitrogen

We now turned our attention to examining different ways of diversifying the scaffolds, beginning with the reactions designed to introduce a basic nitrogen atom into the structures. Thus, tricyclic amides **5{1b}**, **5{5a}**, and **5{6a}** were converted to the corresponding tertiary amines via a previously reported⁵ two-step sequence (Scheme 2 and Table 1).

We had envisioned that direct reductive amination of the ketone carbonyl using $\text{NaBH}(\text{OAc})_3$ ⁶ would provide another tack for introducing basic nitrogen. However, scaffolds with substitution adjacent to the ketone carbonyl proved difficult substrates for direct reductive amination conditions, proceeding with low to no conversion. Acceptable yields⁷ of amines could be obtained across the full range of ketone substrates by preforming the imine using a mixture of TiCl_4 and $\text{Ti}(\text{O}-i\text{-Pr})_4$ ⁸ followed by reduction with NaCNBH_3 to afford amine products (Scheme 3 and Table 2). Unfortunately, inseparable mixtures of diastereomers were obtained using either the two-step reductive amination approach or by direct reductive amination with $\text{NaBH}(\text{OAc})_3$. All ring system examined gave nonselective aminations, presumably due to insufficient diastereofacial direction in the iminium ion intermediates.

One route to diastereomerically-pure amine derivatives is outlined in Scheme 4. Preparative TLC purification of the reductive amination product **7{2}** afforded diastereomerically-pure secondary amines **7{2a}** and **7{2b}**. The relative stereochemistry of the phenylamine group was assigned from the X-ray crystal structure of the amide **9**, afforded by coupling of 3,5-dichlorobenzoic acid with amine **7{2a}** (equation 1). Amines **7{2a}** and **7{2b}** were then reacted with a selection of aldehydes in a second reductive amination to obtain diastereomerically-pure tertiary amines **8{1–4}** (Table 3).

Synthesis of alcohol derivatives

In contrast to the problematic amine reductions, the ketone moiety of the tricycles **5** could be selectively reduced to either diastereomeric alcohol by varying reaction conditions (Scheme 5). Thus, reduction of ketone **5{1b}** with $\text{NaBH}_4/\text{CeCl}_3$ occurred with axial addition of hydride⁹ stereoselectively afforded **10{1b}** as verified by X-ray crystallography (equation 2).

Alternatively, Selectide reduction afforded scaffolds **10{1a-6a}**, which were then converted into a library of carbamates **11{1-24}** (Table 4).

Additional stereoisomeric carbamates were prepared by pairing the alkyl epimerization step with the subsequent reduction step (Scheme 6). Both unepimerized **5{3a}** and epimerized **5{3b}** were reduced by the two complementary reduction methods and subsequently coupled with butyl isocyanate to provide four isomeric carbamate derivatives.

The tricyclic scaffolds **5{1b}** and **3b}** could be regio- and stereoselectively alkylated with primary bromides or iodides (Scheme 7 and Table 5). This method provides a straightforward route to regioisomeric ketones such as **12{1}** and **12{2}** as well as introduces functionalized side chains (e.g., R₂ = *p*-bromophenyl) suitable for follow up chemistry (e.g., by palladium-catalyzed cross coupling). Terminating the side chain with an ester functionality would allow for the synthesis of lactone-containing derivatives that more closely mimic the natural *Stemona* alkaloids by a ketone reduction/lactonization sequence analogous to that used for the syntheses of stenine^{5b} and neostenine^{5c}. The stereochemical assignments of the cis alkylation product are based on precedent for similar alkylations in the syntheses of stenine^{5b} and neostenine^{5c}.

Incorporation of additional heterocycles

Finally, we were interested in ways of appending additional heterocyclic rings to our carbocyclic skeleton. Our initial investigations were focused on applying the Fischer indole synthesis to ketones **5** (Scheme 8). Three sets of conditions for the Fischer indole synthesis were surveyed: zinc chloride in acetic acid (method A), *p*-toluenesulfonic acid in anhydrous ethanol (method B), and simple dissolution in anhydrous ethanol (method C). No one method was superior to the others for every ketone/hydrazine combination. Table 6 summarizes the indole products produced using these methods. The regiochemistry of the reaction was established by obtained an X-ray crystal structure of a representative indole product (**13{3}**). The amide carbonyl on the indole products could be reduced with DIBAL-H to give the indole tertiary amines **14** (Scheme 8 and Table 7) without affecting the indole ring. In all cases, the yields are the results from unoptimized, single attempt trials, which were sufficient in providing material for submission to biological screening collaborations. In one case using method C, no indole product was isolated but instead an isomeric side product **15** was formed in 37% yield (Scheme 9).

Scheme 10 and Table 8 highlight one additional demonstration of the utility of *Stemona* alkaloid scaffolds for the construction of novel polycyclic compounds. The scaffolds with two methylene carbons adjacent to the ketone such as **5{2a}** or **5{6a}** were found to be excellent substrates for the Friedländer quinoline synthesis under the conditions of Miller and coworkers.¹⁰

In summary, we have applied an *endo* selective tandem Diels-Alder/Schmidt reaction to the construction of a series of unnatural neostenine-like scaffolds. The utility of these scaffolds was demonstrated by their elaboration to six sets of functionalized derivatives. The biological evaluation of these compounds as well as the synthesis of additional compound libraries are currently underway and will be reported in due course.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

The authors thank Victor Day for X-ray crystallography and Frank Schoenen for discussions on parallel synthesis technology. This research was funded by the National Institute for General Medical Sciences (KU-CMLD PO50-GM069663).

References

1. For lead references see: (a)Breinbauer R, Vetter IR, Waldmann H. *Angew Chem, Int Ed* 2002;41:2878–2890.2890(b)Tan DS. *Comb Chem High Throughput Screening* 2004;7:631–643.643(c)Oguri H, Schreiber SL. *Org Lett* 2005;7:47–50.50 [PubMed: 15624974](d)Hanessian S, Kothakonda KK. *J Comb Chem* 2005;7:837–842.842 [PubMed: 16283793](e)Shang S, Tan DS. *Curr Opin Chem Biol* 2005;9:248–258.258 [PubMed: 15939326](f)Zacuto MJ, Leighton JL. *Org Lett* 2005;7:5525–5527.5527 [PubMed: 16288547](g)Cohen JL, Limon A, Miledi R, Chamberlin AR. *Bioorg Med Chem Lett* 2006;16:2189–2194.2194 [PubMed: 16455241](h)Zinzalla G, Milroy LG, Ley SV. *Org Biomol Chem* 2006;4:1977–2002.2002 [PubMed: 16688343](i)Barluenga S, Dakas PY, Ferandin Y, Meijer L, Winssinger N. *Angew Chem, Int Ed* 2006;45:3951–3954.3954(j)Dai CF, Cheng F, Xu HC, Ruan YP, Huang PQ. *J Comb Chem* 2007;9:386–394.394 [PubMed: 17358084](k)Wipf P, Graham TH, Xiao J. *Pure Appl Chem* 2007;79:753–761.761(l)Tan, Derek S. *Chem Biol* 2007;483–518.518(m)Shang S, Iwadare H, Macks DE, Ambrosini LM, Tan DS. *Org Lett* 2007;9:1895–1898.1898 [PubMed: 17439132](n)Umarye JD, Lessmann T, Garcia AB, Victor M, Sommer S, Waldmann H. *Chem Eur J* 2007;13:3305–3319.3319(o)Mentel M, Breinbauer R. *Topics Curr Chem* 2007;278:209–241.241(p)Winkler JD, Londregan AT, Hamann MT. *Org Lett* 2007;9:4467–4469.4469 [PubMed: 17918854](q)Kesavan S, Panek JS, Porco JA. *Org Lett* 2007;9:5203–5206.5206 [PubMed: 17997565](r)Galliford CV, Scheidt KA. *Angew Chem, Int Ed* 2007;46:8748–8758.8758(s)Nielsen TE, Schreiber SL. *Angew Chem, Int Ed* 2008;47:48–56.56(t)Yendapally R, Hurdle JG, Carson EI, Lee RB, Lee RE. *J Med Chem* 2008;51:1487–1491.1491 [PubMed: 18281930]
2. For reviews see: (a)Götz M, Edwards OE. *The Alkaloids* Manske RHF. Academic PressNew York1967;9:545–551.551(b)Götz M, Strunz GM. *Alkaloids* Wiesner K. ButterworthLondon1973;9:143–160.160(c)Lin WH, Ye Y, Xu RS. *J Nat Prod* 1992;55:571–576.576 (d)Xu RS. *Stud Nat Prod Chem* 2000;21:729–772.772(e)Pilli RA, Ferreira de Oliveira MdC. *Nat Prod Rep* 2000;17:117–127.127 [PubMed: 10714902](f)Greger H. *Planta Med* 2006;72:99–113.113 [PubMed: 16491444](g)Xu YT, Hon PM, Jiang RW, Cheng L, Li SH, Chan YP, Xu HX, Shaw PC, But PPH. *J Ethnopharmacology* 2006;108:46–53.53
3. (a) Chung HS, Hon PM, Lin G, But PPH, Dong H. *Planta Med* 2003;69:914–920. [PubMed: 14648394] (b) Leung PHH, Zhang L, Zuo Z, Lin G. *Planta Med* 2006;72:211–216. [PubMed: 16534724]
4. Zeng Y, Reddy S, Hirt E, Aubé J. *Org Lett* 2004;6:4993–4995. [PubMed: 15606118]
5. (a) Golden JE, Aubé J. *Angew Chem Int Ed Engl* 2002;41:4316–4318. [PubMed: 12434374] (b) Zeng Y, Aubé J. *J Am Chem Soc* 2005;127:15712–15713. [PubMed: 16277507] (c) Frankowski KJ, Golden JE, Zeng Y, Lei Y, Aubé J. *J Am Chem Soc* 2008;130ASAP
6. Abdel-Magid AF, Carson KG, Harris BD, Maryanoff CA, Shah R. *J Org Chem* 1996;61:3849–3862. [PubMed: 11667239]
7. To satisfy submission requirements to both the KU-CMLD repository and the Molecular Libraries Small Molecule Repository (MLSMR) the criteria for success of analogue production were defined as the generation of greater than 10 mg of product in greater than 90% purity.
8. For leading references on titanium Lewis acid facilitated reductive aminations see: (a)Mattson RJ, Pham KM, Leuck DJ, Cowen KA. *J Org Chem* 1990;55:2552–2554.2554(b)Bhattacharyya S. *Tetrahedron Lett* 1994;35:2401–2404.2404(c)Neidigh KA, Avery MA, Williamson JS, Bhattacharyya S. *J Chem Soc Perkin Trans* 11998;:2527–2532.2532
9. Gemal A, Luche JL. *J Am Chem Soc* 1981;103:5454–5459.
10. McNaughton BR, Miller BL. *Org Lett* 2003;5:4257–4259. [PubMed: 14601974]

neostenine (1)

neotuberostemonine (2)

Figure 1.
Selected antitussive *Stemona* alkaloids.

Scheme 1.

Scheme 2.

Scheme 3.

method A: $NaBH(OAc)_3$

method B: 1. $Ti(OiPr)_4$, $TiCl_4$ 2. $NaCNBH_3$

Scheme 4.

Scheme 5.

Scheme 6.

Scheme 7.

Method A: ZnCl₂ in AcOH

Method B: TsOH in EtOH

Method C: EtOH, no acid

Scheme 8.

Scheme 9.

Scheme 10.

Table 1Reduction of the tricyclic amides **5** to the tertiary amines **6**.

entry	amide	product	yield (%)
1	<i>5{1b}</i>	<i>6{1}</i>	74
2	<i>5{5a}</i>	<i>6{2}</i>	22
3	<i>5{6a}</i>	<i>6{3}</i>	32

Table 2Reductive amination of the tricycles **5**.

entry	scaffold	amine	product	yield (%)
1	5{1b}	aniline	7{1}	55
2	5{2a}	aniline	7{2}	79
3	5{3b}	aniline	7{3}	55
4	5{4b}	aniline	7{4}	60
5	5{5a}	aniline	7{5}	31
6	5{6a}	aniline	7{6}	35
7	5{1b}	4-phenylpiperizine	7{7}	22
8	5{2a}	4-phenylpiperizine	7{8}	54
9	5{3b}	4-phenylpiperizine	7{9}	38
10	5{4b}	4-phenylpiperizine	7{10}	30
11	5{5a}	4-phenylpiperizine	7{11}	41
12	5{6a}	4-phenylpiperizine	7{12}	33

Table 3
Reductive amination of the diastereomerically-pure amines **7** with aldehydes.

entry	substrate	R	method	product	yield
1	7[2a]	<i>n</i> -hexyl	A	8[1]	85%
2	7[2b]	2-benzo[b]thiophene	A	8[2]	51%
3	7[2b]	2-naphthyl	B	8[3]	91%
4	7[2b]	<i>n</i> -hexyl	A	8[4]	75%

Table 4

Construction of a 24-member carbamate library.^a

entry	alcohol	R ¹	R ²	carbamate	yield (%)	purity (%)
1	10{1a}	Et	<i>n</i> -butyl	11{1a}	25	95
2	10{1a}	Et	phenyl	11{2a}	67	90
3	10{1a}	Et	4-bromophenyl	11{3a}	25	100
4	10{1a}	Et	5-benzo[d][1,3]dioxole	11{4a}	43	93
5	10{2a}	H	<i>n</i> -butyl	11{5a}	90	100
6	10{2a}	H	phenyl	11{6a}	0	—
7	10{2a}	H	4-bromophenyl	11{7a}	5	100
8	10{2a}	H	5-benzo[d][1,3]dioxole	11{8a}	30	100
9	10{3a}	Bn	<i>n</i> -butyl	11{9a}	35	100
10	10{3a}	Bn	phenyl	11{10a}	28	96
11	10{3a}	Bn	4-bromophenyl	11{11a}	2	100
12	10{3a}	Bn	5-benzo[d][1,3]dioxole	11{12a}	33	100
13	10{4a}	Et	<i>n</i> -butyl	11{13a}	0	—
14	10{4a}	Et	phenyl	11{14a}	19	100
15	10{4a}	Et	4-bromophenyl	11{15a}	37	100
16	10{4a}	Et	5-benzo[d][1,3]dioxole	11{16a}	26	85
17	10{5a}	H	<i>n</i> -butyl	11{17a}	49	100
18	10{5a}	H	phenyl	11{18a}	45	100
19	10{5a}	H	4-bromophenyl	11{19a}	15	100
20	10{5a}	H	5-benzo[d][1,3]dioxole	11{20a}	34	100
21	10{6a}	H	<i>n</i> -butyl	11{21a}	52	100
22	10{6a}	H	phenyl	11{22a}	82	99
23	10{6a}	H	4-bromophenyl	11{23a}	20	100
24	10{6a}	H	5-benzo[d][1,3]dioxole	11{24a}	98	81

^aSee Scheme 5 for the structures of **10** and **11**.

Table 5

Regioselective alkylation of tricyclic ketones **5**.

entry	substrate	R ¹	R ²	product	yield (%)
1	5(1b)	Et	Ph	12(1)	74
2	5(3b)	Bn	Me	12(2)	30
3	5(1b)	Et	CO ₂ Et	12(3)	95
4	5(1b)	Et	2-Br-C ₆ H ₅	12(4)	74

Table 6

The Fischer indole synthesis on tricyclic ketones **5**.

entry	R ¹	R ²	method	product	yield (%)
1	Et	4- <i>i</i> -Pr	C	13{1}	26
2	Et	4-OMe	C	13{2}	77
3	H	H	A	13{3}	76
4	H	4- <i>i</i> -Pr	B	13{4}	24
5	H	4-OMe	B	13{5}	18
6	H	2-Cl	B	13{6}	37
7	H	2,5-Cl ₂ Cl	B	13{7}	37

Table 7
Combined Fischer indole synthesis and subsequent amide reduction of ketones **5**.

entry	R ¹	R ²	method	product	yield (%)
1	H	4- <i>i</i> -Pr	B	14[1]	13
2	H	4-OMe	B	14[2]	8
3	H	2-Cl	B	14[3]	23

Table 8The Friedländer quinoline synthesis on tricyclic ketone **5**.

entry	ketone	R	product	yield (%)
1	5{2a}	H	16{1}	68
2	5{2a}	5-OH	16{2}	83
3	5{2a}	5-Cl	16{3}	86
4	5{2a}	4-CF ₃	16{4}	73
5	5{2a}	[4,5]-dioxole	16{5}	48
6	5{2a}	5-CO ₂ Me	16{6}	76
7	5{6a}	H	16{7}	53