Published in final edited form as:

Rejuvenation Res. 2008 February ; 11(1): 251–257. doi:10.1089/rej.2007.0611.

AN INVERSE ASSOCIATION BETWEEN SELF-REPORTED ARTHRITIS AND MORTALITY IN THE ELDERLY: FINDINGS FROM THE NATIONAL LONG TERM CARE SURVEY

Alexander M Kulminski, Irina V Kulminskaya, Svetlana V Ukraintseva, Kenneth Land, and Anatoli I Yashin, PhD

Center for Population Health and Aging, Duke University Population Research Institute, Trent Hall, Room 002, Box 90408, Durham, NC, 27708 and Department of Sociology, Duke University, Trent Hall, Room 002, Box 90408, Durham, NC, 27708

Abstract

Major musculoskeletal conditions including arthritis represent an increasing burden on individuals and societies. We analyze the association between self-reported arthritis and mortality in the U.S. elderly disabled and non-disabled individuals using unique disability-focused data from the large-scale population-based National Long Term Care Survey. It is found that males and females who reported arthritis/rheumatism have, generally, smaller risks of death than those who did not report those conditions. This inverse relationship is more pronounced in disabled individuals. This finding holds for both short- (Relative Risk [RR]=0.81; 95% Confidence Interval [CI]:0.75–0.88 for males and RR=0.76; CI:0.71–0.82 for females) and long- (RR=0.82; CI:0.78–0.87 for males and RR=0.83; CI:0.79–0.87 for females) term follow-ups. For females, this effect is age insensitive, while for males it is limited to ages below 85. Demographic and 19 major geriatric conditions have trivial effect on these risks supporting the view that a better survival of diseased individuals can be attributed to the effects of medical treatment. Given the widespread prevalence of arthritis/rheumatism and disability in elderly populations and the increasing population of the elderly, these findings call for comprehensive analyses of factors driving better survival and medical costs associated with extended lives.

Keywords

Aging; disease; longevity; mortality

INTRODUCTION

Major musculoskeletal conditions including arthritis represent an increasing burden on individuals and societies in terms of worsening health and increasing health care consumption with associated medical costs [1]. Millions Americans are affected by various types of arthritis with osteoarthritis being the most frequent form among the elderly [2–7]. Projections show that the prevalence of arthritis in American population will likely substantially increase during the next 30 to 40 years, particularly among the elderly (a 2.3-fold increase in the U.S. to the year 2050 compared to 1.3 for adults aged 20 to 65) [6,8]. In the U.S., osteoarthritis is one of the most common causes of disability [9], which is as great as that attributable to cardiovascular disease and is greater than the risk attributable to any other health disorder in the elderly [2,

10]. Global trends toward increasing life expectancy can make osteoarthritis the fourth leading cause of disability in the world by the year 2020 [1].

Rheumatoid arthritis is the second common type of arthritis (after osteoarthritis), which is, however, typical for younger individuals [11]. While rheumatoid arthritis has been extensively studied for its association with excessive mortality [12–15], knowledge about the association between mortality and osteoarthritis remains limited [16]. Generally, studies do not support association between increased risk of mortality and osteoarthritis [17,18]. Some studies examined mortality among individuals with self-reported arthritis. For example, no significant increase in mortality during 15 years of follow-up has been found among individuals, who reported arthritis or osteoarthritis as compared with those without these disorders [19]. The Epidemiological Follow-up of the National Health and Nutrition Examination Survey I also has found no significant association between arthritis and mortality excess [20]. The latter study even provides evidence for significant negative relationship between certain types of osteoarthritis and mortality. Some other studies, however, have reported positive associations between particular types of osteoarthritis and mortality [21]. The results are, therefore, controversial and require better understanding of the relationships between arthritis and survival chances, particularly in the elderly individuals.

In this paper, we analyze relationships between arthritis and mortality using unique disability-focused data from the National Long Term Care Survey (NLTCS) linked with Medicare vital statistics files to comprehensively assess the risk of death among individuals who reported and who did not report arthritis or rheumatism. Over-sampling of disabled individuals in the detailed NLTCS instruments provided an opportunity to assess these connections separately for disabled and non-disabled individuals. We also analyzed how major health conditions and demographic factors modulate these associations. Since the NLTCS participants answered a question concerning whether they have rheumatism or arthritis, no distinction was made between these disorders.

METHODS

The data are from the 1982, 1984, 1989, 1994, and 1999 NLTCS and the linked Medicare vital statistics files (from 1982 to August, 2003). The NLTCS is considered to be one of the best designed surveys to assess chronic disability (activity limitations that last 90+ days) in the U.S. elderly (65+) individuals [22,23]. To complete the NLTCS, a two-stage interviewing process was used. A screening interview assessing chronic disability was given to all participants. A detailed interview was given to i) those who reported at least one chronic impairment in (instrumental) activities of daily living, ii) institutionalized individuals, and iii) those who received a detailed interview in a previous survey. For each new wave of the survey, a cohort sample of about 5,000 persons was added to the surviving sample to replace deaths occurring since the previous survey and to ensure that the new sample was representative of the U.S. elderly population. Such a procedure ensures a valid longitudinal and cross-sectional design for the survey. The 1994 and 1999 surveys also explicitly included samples of individuals who were designated for detailed interviews before being give a test on disability (see [24] for details).

We performed sex-specific analyses assessing relative risks of death using Cox regression models adjusted for demographic (age, race, education, marital status) and self-reported health-related (diabetes, cancer, overweight/obesity, heart diseases, hypertension, stroke, paralysis, other permanent numbness/stiffness besides paralysis/rheumatism/arthritis, multiple sclerosis, cerebral palsy, epilepsy, Parkinson's disease, atherosclerosis, pneumonia, bronchitis, flu, emphysema, asthma, and fractures) conditions. In preliminary analyses, we found that the

effect of disability is best quantified as presence of any impairment vs. no disability. Consequently, only dichotomous disability indicator is used.

Time of follow up measured was used as the time scale in the Cox regressions. We verified that the proportionality assumption of the hazard in the models was a reasonable approximation even for long term follow-ups. Nevertheless, we calculated relative risks of death which occurred during different periods of follow up. The results reported below concern representative analyses of short- (4 years) and long- (up to 22 years, depending on the wave) term follow-up periods. To test robustness of the estimates, we analyzed each of the five NLTCs waves and the entire pooled sample.

RESULTS

Table 1 shows descriptive characteristics for participants of each of the five NLTCs waves and for the pooled sample. The proportion of deaths that occurred within the 22-year observation period decreases as the length of this period decreases for the later waves. The sharp decrease of the percentage of disabled individuals in 1994 and 1999 NLTCs is due to the change of the survey design when the “healthy” supplements were added (individuals who were designated to receive community detail interview before screening on disability). Given over-sampling of disabled individuals, the arthritis prevalence is consistent with the CDC estimates for self-reported arthritis in the U.S. for year 2001, which is about 60% among the elderly 65+ (both sexes), and higher among females [25].

Univariate regression analysis (Table 2, M0) shows that both males and females in the pooled sample who reported on arthritis/rheumatism have lower risks of death than those who did not report those conditions. These estimates are highly significant for females while for males they can be non-significant for long follow-ups. Survey-specific analysis shows that the relative risks of death are significantly lower for males and females with arthritis/rheumatism from the early NLTCs irrespective of the follow-up period. For the later NLTCs waves, these risks are less convincing. Given that the number of disabled individuals decreases in the later surveys, it is reasonable to expect that disability might affect the estimated risks. If so, then adjustment for disability might provide more convincing numbers for the later waves. Indeed, Table 2 (M1) shows that relative risks of death for females become significant irrespective of the NLTCs wave and follow-up period. For males who reported arthritis conditions, the risks of death are either significantly lower or the same as for those who did not report those conditions. Adjusting models for other possible confounders (demographic and health-related factors; see Section “Analysis”) does not substantially change the estimates. For shorter follow-up periods, the relative risks of death for individuals with arthritis/rheumatism tend to be lower than those for longer periods.

Multivariate analysis applying the full model (M2) to the pooled sample for each sex stratified according to the disability status shows significantly lower risks of death for disabled males and females with arthritis/rheumatism compared to disabled individuals without arthritis (Table 3, Disability). For non-disabled males, the risks do not significantly differ between diseased and non-diseased groups; they, however, tend to be lower for diseased individuals. For non-disabled females, the risks of death are significantly lower for longer follow-up periods for those who reported arthritis conditions. Stratification for more homogeneous age groups shows that age is not an essential factor for females, while better survival for diseased males is limited to 85 years (Table 3, Age).

Stratification by disability and age (Table 4) shows that younger (65–84) males with arthritis and disability have significantly lower risks of death than disabled individuals without arthritis. For older males who are disabled and for non-disabled males irrespective of age, these risks

do not significantly differ between diseased and non-diseased individuals irrespective of follow-up period. However, for the oldest-old non-disabled males (85+), the relative risks tend to be well below unity. Disabling conditions increase the chances of death of diseased oldest-old males making them the same irrespective of arthritis/rheumatism and follow-up period.

For non-disabled females (Table 4) for short-term follow-up periods, the risks of death do not significantly differ from unity. For longer follow-up periods, younger diseased females (65–74) have lower chances of death than non-diseased females. Contrarily to males, the oldest-old (85+) diseased and non-disabled females tend to have larger chances of death than non-diseased and non-disabled counterparts. For disabled females, the relative risks are highly significant being lower for females with arthritis or rheumatism.

DISCUSSION AND CONCLUSIONS

This study uses unique data from the large-scale, population-based NLTCS, which focuses on disabled individuals. The study shows very intriguing effect of arthritis/rheumatism and disability on survival. We found that both males and females having arthritis or rheumatism (self-reports; “Do you have arthritis or rheumatism?”) tend to have significantly *lower* chances to die than those who do not suffer of those disorders. Disability is found to be a strong modulating factor which *decreases* the chances of death for diseased males and females. The estimates for disabled individuals are robust and highly significant, indicating, for instance, that diseased females and males have 24% (Relative Risk [RR]=0.76; 95% Confidence Interval [CI]: 0.71–0.82) and 19% (RR=0.81; CI: 0.75–0.88), respectively, lower chances for short-term deaths than non-diseased counterparts. For long-term deaths, the risks become practically identical for both sexes (RR=0.83; CI: 0.79–0.87 for females and RR=0.82; CI: 0.78–0.87 for males). For females, this effect is practically age insensitive. For males, however, this effect is limited basically to ages below 85. The risks of death for diseased individuals are lower both for deaths occurred during short- and long-term periods, i.e., this effect is long-standing. Other factors besides disability (i.e., demographic characteristics and 19 health-related conditions) do not alter the revealed relationships contributing little to their explanation.

One obvious explanation could be that arthritis causes less fatal types of disability compared to other factors (e.g., cardiovascular conditions, cancer). Therefore, disabled individuals with arthritis could be of better health than disabled individuals without arthritis. This explanation is, however, not entirely convincing since: i) adjustments of the analyses for other potential causes of disability including more fatal diseases (e.g., cardiovascular conditions, cancer) had trivial effects on the estimates (compare Table 2, models 1 and 2), ii) for stratified samples (Table 3 and Table 4), the risks were evaluated for model with adjustment for major health-related conditions for the elderly, and iii) this effect holds for individuals (females) having no disabilities (i.e., who are presumably in better health).

Another possible explanation for lower risks of death for individuals with arthritis could be as side-effects of treatment of these conditions. This possibility is supported by the fact that this effect is highly significant among disabled individuals. Given that arthritis/rheumatism are among the leading causes of disability in the elderly [2,9], such individuals can receive more attention from health care providers and, thus, experience better treatment due to their worse conditions than those who do not have disabilities. Typically, current treatment recommendations include the use of various drugs (e.g., acetaminophen, NSAIDs), exercise, education interventions, and joint arthroplasty [7,11]. Moreover, conventional regimes use intensive treatments that combine different types of treatments (e.g., drugs with exercises). Thus, intensive or radical (e.g., surgery) treatments can lead to improving health of such individuals, which increases their chances of surviving. On the other hand, the effects of certain drugs can promote better survival. For instance, treating arthritis with NSAIDs can reduce the

risk of death from all causes, although such treatment can increase the risk of cardiovascular diseases [26]. Some anti-inflammatory compounds also appear to be factors positively affecting aging processes [27].

These findings have implications both for understanding perspectives of regular treatment of the elderly patients as a potential anti-aging intervention and for public health, caregivers, and policymakers. Immediate important concerns that should be addressed are which specific factors contribute to life span extension for these individuals and whether the life extension results in a corresponding high load on Medicare spending or not. These concerns, however, deserve separate analyses.

ACKNOWLEDGMENTS

The research reported in this paper was supported by the National Institute on Aging grants 1R01 AG028259, 1R01-AG-027019 and 5P01-AG-008761.

REFERENCES

1. Davenport G. Rheumatology and musculoskeletal medicine. *Br J Gen Pract* 2004;54:457–464. [PubMed: 15186570]
2. Felson DT, Lawrence RC, Dieppe PA, Hirsch R, Helmick CG, Jordan JM, Kington RS, Lane NE, Nevitt MC, Zhang Y, Sowers M, McAlindon T, Spector TD, Poole AR, Yanovski SZ, Ateshian G, Sharma L, Buckwalter JA, Brandt KD, Fries JF. Osteoarthritis: new insights. Part 1: the disease and its risk factors. *Ann Intern Med* 2000;133:635–646. [PubMed: 11033593]
3. Abyad A, Boyer JT. Arthritis and aging. *Curr Opin Rheumatol* 1992;4:153–159. [PubMed: 1581142]
4. Reginster JY. The prevalence and burden of arthritis. *Rheumatology (Oxford)* 2002;41:3–6. [PubMed: 12173279]
5. CDC. State prevalence of self-reported doctor-diagnosed arthritis and arthritis-attributable activity limitation—United States, 2003. *MMWR Morb Mortal Wkly Rep* 2006;55:477–481. [PubMed: 16675943]
6. Fontaine KR, Haaz S, Heo M. Projected prevalence of US adults with self-reported doctor-diagnosed arthritis, 2005 to 2050. *Clin Rheumatol*. 2007
7. Felson DT, Lawrence RC, Hochberg MC, McAlindon T, Dieppe PA, Minor MA, Blair SN, Berman BM, Fries JF, Weinberger M, Lorig KR, Jacobs JJ, Goldberg V. Osteoarthritis: new insights. Part 2: treatment approaches. *Ann Intern Med* 2000;133:726–737. [PubMed: 11074906]
8. Hootman JM, Helmick CG. Projections of US prevalence of arthritis and associated activity limitations. *Arthritis Rheum* 2006;54:226–229. [PubMed: 16385518]
9. Birchfield PC. Osteoarthritis overview. *Geriatr Nurs* 2001;22:124–130. [PubMed: 11410762]quiz 130–121
10. Jamison, DT.; World Bank. Disease control priorities in developing countries. Vol. ed 2nd. New York, NY, Washington, DC: Oxford University Press; World Bank; 2006.
11. Ross C. A comparison of osteoarthritis and rheumatoid arthritis: diagnosis and treatment. *Nurse Pract* 1997;22:20, 23–24, 27–28 passim. [PubMed: 9314163]quiz 39–41
12. Goodson N, Symmons D. Rheumatoid arthritis in women: still associated with an increased mortality. *Ann Rheum Dis* 2002;61:955–956. [PubMed: 12379514]
13. Guedes C, Dumont-Fischer D, Leichter-Nakache S, Boissier MC. Mortality in rheumatoid arthritis. *Rev Rhum Engl Ed* 1999;66:492–498. [PubMed: 10567978]
14. Bjornadal L, Baecklund E, Yin L, Granath F, Klareskog L, Ekbom A. Decreasing mortality in patients with rheumatoid arthritis: results from a large population based cohort in Sweden, 1964–95. *J Rheumatol* 2002;29:906–912. [PubMed: 12022348]
15. Thomas E, Symmons DP, Brewster DH, Black RJ, Macfarlane GJ. National study of cause-specific mortality in rheumatoid arthritis, juvenile chronic arthritis, and other rheumatic conditions: a 20 year followup study. *J Rheumatol* 2003;30:958–965. [PubMed: 12734889]

16. Scott DL, Houssien DA. Clinical and laboratory assessments in rheumatoid arthritis and osteoarthritis. *Br J Rheumatol* 1996;35:6–9. [PubMed: 9010081]
17. Manuel DG, Luo W, Ugnat AM, Mao Y. Cause-deleted health-adjusted life expectancy of Canadians with selected chronic conditions. *Chronic Dis Can* 2003;24:108–115. [PubMed: 14733760]
18. Watson DJ, Rhodes T, Guess HA. All-cause mortality and vascular events among patients with rheumatoid arthritis, osteoarthritis, or no arthritis in the UK General Practice Research Database. *J Rheumatol* 2003;30:1196–1202. [PubMed: 12784389]
19. Holbrook TL, Wingard DL, Barrett-Connor E. Self-reported arthritis among men and women in an adult community. *J Community Health* 1990;15:195–208. [PubMed: 2365841]
20. Leigh JP, Fries JF. Arthritis and mortality in the epidemiological follow-up to the National Health and Nutrition Examination Survey I. *Bull N Y Acad Med* 1994;71:69–86. [PubMed: 8069278]
21. Haara MM, Manninen P, Kroger H, Arokoski JP, Karkkainen A, Knekt P, Aromaa A, Heliovaara M. Osteoarthritis of finger joints in Finns aged 30 or over: prevalence, determinants, and association with mortality. *Ann Rheum Dis* 2003;62:151–158. [PubMed: 12525385]
22. Freedman VA, Martin LG, Schoeni RF. Recent trends in disability and functioning among older adults in the United States: a systematic review. *JAMA* 2002;288:3137–3146. [PubMed: 12495394]
23. Fries JF. Measuring and monitoring success in compressing morbidity. *Ann Intern Med* 2003;139:455–459. [PubMed: 12965976]
24. Manton KG, Gu X, Lamb VL. Change in chronic disability from 1982 to 2004/2005 as measured by long-term changes in function and health in the U.S. elderly population. *Proc Natl Acad Sci U S A* 2006;103:18374–18379. [PubMed: 17101963]
25. CDC. Prevalence of Self-Reported Arthritis or Chronic Joint Symptoms Among Adults – United States, 2001. *MMWR Morb Mortal Wkly Rep* 2002;51:948–950. [PubMed: 12437034]
26. Lee TA, Bartle B, Weiss KB. Impact of NSAIDS on mortality and the effect of preexisting coronary artery disease in US veterans. *Am J Med* 2007;120:98. [PubMed: 17208086]e99-16
27. Ukraintseva SV, Arbeev KG, Michalsky AI, Yashin AI. Antiaging treatments have been legally prescribed for approximately thirty years. *Annals of the New York Academy of Sciences* 2004;1019:64–69. [PubMed: 15246996]

Table 1
Descriptive characteristics for males (M) and females (F) participating in each of five NLTCS and for pooled sample of participants of all surveys (ALL)

Sample	Sex	N	ND4, %	ND22, %	Disability, yes, %	Arthritis, yes, %
1982	M	2166	43.8	97.1	87.7	63.5
	F	3921	32.7	95.3	93.1	78.6
1984	M	2038	39.9	95.7	83.3	65.1
	F	3891	31.0	92.4	89.1	77.0
1989	M	1470	39.3	89.7	77.2	64.2
	F	2992	29.2	85.1	85.6	76.3
1994	M	1741	36.3	65.9	57.3	58.0
	F	3347	29.3	60.3	68.9	70.7
1999	M	1805	35.3	35.3	56.8	51.7
	F	3341	30.5	30.6	69.5	67.1
ALL	M	9220	39.1	77.6	73.3	60.4
	F	17492	30.7	73.9	81.8	73.9

ND4(ND22)= number of deaths within 4 (up to 22) years of follow up after the respective interview.

Table 2
Cox proportional hazard models of the relative risks (RR) of deaths for males (M) and females (F) reported on arthritis or rheumatism vs. those who did not report those conditions.

Model	Follow-up, yrs	Sex	ALL		1982		1984		1989		1994		1999	
			RR	95% CI	RR	95% CI	RR	95% CI	RR	95% CI	RR	95% CI	RR	95% CI
M0	4	M	0.91	0.85, 0.97	0.75	0.66, 0.85	0.78	0.67, 0.89	0.84*	0.71, 0.99	1.00 [†]	0.86, 1.18		
		F	0.82	0.77, 0.86	0.68	0.60, 0.77	0.76	0.67, 0.86	0.83*	0.72, 0.97	0.83	0.73, 0.95		
	22	M	0.96 [†]	0.91, 1.00	0.82	0.75, 0.90	0.87	0.80, 0.96	0.94 [†]	0.84, 1.05	1.13 [†]	1.00, 1.27	1.30	1.11, 1.53
		F	0.88	0.84, 0.91	0.82	0.76, 0.88	0.86	0.80, 0.93	0.87	0.79, 0.95	0.93 [†]	0.85, 1.02	0.99 [†]	0.87, 1.13
M1	4	M	0.82	0.77, 0.88	0.76	0.67, 0.87	0.76	0.66, 0.87	0.79	0.67, 0.93	0.84*	0.71, 0.98		
		F	0.74	0.70, 0.79	0.68	0.61, 0.78	0.75	0.66, 0.86	0.80	0.68, 0.92	0.71	0.62, 0.81		
	22	M	0.88	0.84, 0.92	0.82	0.75, 0.90	0.82	0.75, 0.90	0.90 [†]	0.80, 1.01	0.93 [†]	0.83, 1.05	1.01 [†]	0.86, 1.18
		F	0.82	0.79, 0.86	0.82	0.76, 0.88	0.86	0.73, 0.93	0.81	0.74, 0.89	0.77	0.70, 0.85	0.80	0.70, 0.91
M2	4	M	0.81	0.76, 0.87	0.74	0.65, 0.86	0.74	0.63, 0.86	0.84 [†]	0.70, 1.01	0.78	0.66, 0.94		
		F	0.77	0.73, 0.83	0.70	0.61, 0.80	0.80	0.70, 0.92	0.83*	0.70, 0.98	0.77	0.66, 0.89		
	22	M	0.85	0.81, 0.90	0.80	0.73, 0.88	0.79	0.71, 0.87	0.88*	0.78, 1.00	0.88 [†]	0.77, 1.00	1.01 [†]	0.85, 1.21
		F	0.83	0.80, 0.87	0.81	0.74, 0.88	0.88	0.81, 0.96	0.83	0.76, 0.92	0.79	0.71, 0.88	0.81	0.70, 0.93

* 0.01 < p < 0.05
[†] p > 0.05; for other estimates p < 0.01

ALL denotes pooled sample of all five NLTCs waves; CI=Confidence Interval

Note that for 4-year and up to 22-year follow-up periods the relative risks for 1999 NLTCs are the same and, thus, are omitted for the 4-year follow-up

M0: univariate model which includes only arthritis/rheumatism

M1: M0 + age and disability indicator

M2: M1 + demographic factors and self-reported health-related conditions (see Section "Analysis")

Table 3

Multivariate Cox proportional hazard model (model 2 from Table 2) of the relative risks (RR) of deaths for pooled sex-specific samples of individuals reported on arthritis or rheumatism vs. those who did not report those conditions in groups stratified by disability status or age.

Stratification	Males				Females			
	Follow-up period, yrs							
	4	22	4	22	4	22	4	22
Disability	RR	95% CI	RR	95% CI	RR	95% CI	RR	95% CI
	No	0.84 [†]	0.69, 1.02	0.97 [†]	0.86 [†]	0.68, 1.08	0.84	0.74, 0.95
	Yes	0.81	0.75, 0.88	0.82	0.76	0.71, 0.82	0.83	0.79, 0.87
Age yrs	65–74	0.79	0.69, 0.90	0.85	0.74	0.64, 0.85	0.79	0.73, 0.86
	75–84	0.77	0.69, 0.85	0.82	0.79	0.71, 0.87	0.85	0.79, 0.91
	85+	0.96 [†]	0.83, 1.11	0.95 [†]	0.79	0.72, 0.88	0.87	0.80, 0.94

[†] $p > 0.05$

Table 4

Multivariate Cox proportional hazard model (model 2 from Table 2) of the relative risks (RR) of deaths for pooled sex-specific samples of individuals reported on arthritis or rheumatism vs. those who did not report those conditions in groups stratified by disability status and age.

Stratification		Males				Females			
		4		22		4		22	
		RR	95% CI	RR	95% CI	RR	95% CI	RR	95% CI
Disability	Age								
No	65-74	0.85 [†]	0.59, 1.21	0.98 [†]	0.81, 1.19	0.74 [†]	0.48, 1.13	0.74	0.61, 0.90
	75-84	0.87 [†]	0.66, 1.14	0.98 [†]	0.83, 1.16	0.86 [†]	0.62, 1.20	0.85 [†]	0.71, 1.02
	85+	0.68 [†]	0.38, 1.21	0.65 [†]	0.42, 1.02	1.28 [†]	0.73, 2.26	1.32 [†]	0.93, 1.88
Yes	65-74	0.78	0.67, 0.90	0.81	0.73, 0.89	0.73	0.62, 0.85	0.78	0.71, 0.85
	75-84	0.75	0.67, 0.84	0.79	0.72, 0.86	0.78	0.71, 0.87	0.84	0.78, 0.91
	85+	1.0 [†]	0.85, 1.17	0.97 [†]	0.84, 1.11	0.78	0.70, 0.87	0.85	0.78, 0.92

[†] p>0.05