Published in final edited form as:

J Am Chem Soc. 2008 September 10; 130(36): 11852–11853. doi:10.1021/ja8038954.

Ligand Controlled Highly Regio- and Enantio-selective Synthesis of α -Acyloxyketones by Palladium Catalyzed Allylic Alkylation of 1,2-Enediol Carbonates

Barry M. Trost, Jiayi Xu, and Thomas Schmidt

Department of Chemistry, Stanford University, Stanford, California 94305–5080.

bmtrost@stanford.edu

α -Hydroxy carbonyl compounds represent a structural type of both synthetic and biological importance. We previously noted that the enol allyl carbonates of α -siloxy carbonyl compounds underwent smooth palladium catalyzed decarboxylative asymmetric allylic alkylation (AAA) ¹ to allylated α -siloxyaldehydes using a Pd complex bearing the **L_{anth}** ligand regardless of the regioisomeric nature of the starting material (ie. **I** or **II** in eq. 1).² The regioselectivity may be interpreted as a faster equilibration between the Pd enolate **A** and **B** compared to the rate of alkylation which occurs faster via **B** (ie. $k > k_2 > k_1$). However, if **A** and **B** exist as either tight ion-pairs or covalently bonded enolates as we proposed before,^{1c} the Pd catalyst should be involved in **both** R migration and enolate alkylation steps. Thus, by tuning the ligand and the potential migrating group, we envisioned that we could change the reaction pattern in favor of the formation of α -hydroxyketones **III**, which have attracted much attention due to their versatile roles in organic synthesis.^{1k, 3} Herein, we report our success in the highly regio- and enantio-selective synthesis of α -acyloxyketones by such an approach.

Initially we investigated the role of ligands by using carbonate **1a** ($R = \text{tert-butyl dimethylsilyl}$, TBS) as the substrate, which as we reported previously, in the presence of **L_{anth}** decarboxylatively alkylated to the corresponding siloxyaldehyde **4a** with high regioselectivity (Table 1, entry 1). **PHOX** ligands, which, similar to **L_{anth}**, have also been successfully used to catalyze the decarboxylative AAA of enol allyl carbonates, favored the formation of the aldehyde product in a ratio of 4.2 to 1; however, the ee of **3a** was much lower (17%, entry 4). In contrast to these results, varying our ligands to **L_{std}** and **L_{stlb}**, slightly favored the formation of the ketone product (entry 2 and 3). The best selectivity (**3a/4a** = 17/1) was achieved by using

E-mail: bmtrost@stanford.edu.

Supporting Information **Available:** Experimental procedures and characterization data for all new compounds (PDF). This material is available free of charge via the Internet at <http://pubs.acs.org>.

L_{naph} (entry 5). Replacement of OTBS with OAc almost completely suppressed the formation of the aldehyde product (entry 6). Changing solvent from dioxane to 1,2-dimethoxyethane (DME), kept the excellent regioselectivity but also improved the ee of **3b** to 90% (entry 7).⁴ The ligand-dependence of the product distribution of **1b** was similar to that of **1a**, although in all cases the ketone product was the major one (entry 8–11). Besides acetoxy other ester groups were also investigated and **3d** with R = pivaloyl (Piv) had the highest ee value (94%, entry 13). Starting from **2** (R = TBS or Piv), which, after decarboxylation initially generated the more stable Pd enolate **B**, only the aldehyde product was exclusively generated in the presence of **L_{naph}** (entry 14 and 15). This suggests that the equilibrium between **A** and **B** is slower than the alkylation steps ($k < k_1, k_2$), in stark contrast to the reaction catalyzed by **L_{anth}**. The same reaction catalyzed by **PHOX** ligand (entry 16), however, gave a similar amount of aldehyde **4a** (**3a/4a** = 1/7.7) as in the reaction of entry 4 (**3a/4a** = 1/4.2), implying a faster equilibrium and comparably slower alkylation ($k > k_1 \approx k_2$).

The scope of the reaction has been investigated and the results are summarized in Table 2. Besides the aromatic ketones (entry 1–5), enones such as **12b** and **12d** (entry 6 and 7), as well as aliphatic ketones such as **14** and **16** (entry 8 and 9) can be obtained in good yields and high ee's. In general, pivalate protected α -hydroxyketones have moderately higher ee's than the corresponding acetate protected ones; however, acetate is easier to be removed without loss of the enantioselectivity of the α -hydroxyketone. Substrates with a substituted allylic moiety also reacted with full conversions, in some cases at slightly warmer temperature (40 °C) (entry 10–12). The dr's of the corresponding products are over 95/5, and the ee values of the major diastereomers are higher than that of **3b**. These high dr's are reflected in the excellent ee's of **29** and **30** by the hydrogenations of **20** and **22** respectively (eq. 2). More interestingly, OR can be a functionalized group as in **24**, **26**, and **28** (entry 13–15). Such functionality can be useful in further structural elaboration as illustrated by the treatment of **24** with Grubbs II catalyst to afford lactone **31** without any erosion of enantioselectivity (eq. 3).

In summary, the palladium catalyzed decarboxylative AAA of 1,2-enediol carbonates can be precisely controlled by the selection of the ligand to generate either regio-isomer. Interestingly, although acyl migration in sodium enolates is fast even at -78°C ,⁵ such equilibration is much slower with these Pd enolates and shows a ligand dependence. In the case of using **L_{naph}** as ligand it is slower than the alkylation, so that no migration is observed even above room temperature. This supports the concept that the decarboxylative AAA of ketones reacts through a tight ion pair or covalently bonded Pd enolate intermediates.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgement

We thank the National Science Foundation and the National Institutes of Health, General Medical Sciences Grant GM13598, for their generous support of our programs. J. Xu has been supported by Abbott Laboratories Fellowships. We thank Chirotech (now Dow) for their generous gifts of ligands and Johnson Matthey for gifts of palladium salts.

References

1. aBehenna DC, Stoltz BM. *J. Am. Chem. Soc* 2004;126:15044. [PubMed: 15547998]; bTrost BM, Xu J. *J. Am. Chem. Soc* 2005;127:2846. [PubMed: 15740108]; cTrost BM, Xu J. *J. Am. Chem. Soc* 2005;127:17180. [PubMed: 16332054]; For reviews on decarboxylative AAA of simple ketones, see: dYou S-L, Dai L-X. *Angew. Chem. Int. Ed* 2006;45:5246.; eBraun M, Meier T. *Angew. Chem., Int. Ed* 2006;45:6952.; fBraun M, Meier T. *Syn. Lett* 2006;5:661.; gStoltz BM, Mohr JT. *Chem. Asian J* 2007;2:1476. [PubMed: 17935094]. For more examples in metal catalyzed AAA of simple ketones, see: hTrost BM, Schroeder GM. *J. Am. Chem. Soc* 1999;121:6759.; iTrost BM, Schroeder GM. *Chem. Eur. J* 2005;11:174.; jBraun M, Laicher F, Meier T. *Angew. Chem., Int. Ed* 2000;39:3494.; kYan XX, Liang CG, Zhang Y, Hong W, Cao BX, Dai LX, Hou XL. *Angew. Chem., Int. Ed* 2005;44:6544.; lGraening T, Hartwig JF. *J. Am. Chem. Soc* 2005;127:17192. [PubMed: 16332060]; mBélanger É, Cantin K, Messe O, Tremblay M, Paquin J-F. *J. Am. Chem. Soc* 2007;129:1034. [PubMed: 17263376]; nZheng W-H, Zheng B-H; Yan Z, Hou X-L. *J. Am. Chem. Soc* 2007;129:7718. [PubMed: 17539637]; oDoyle AG; Jacobsen EN. *Angew. Chem., Int. Ed* 2007;46:3701.; pHe H, Zheng X-J, Li Y, Dai L-X, You S-L. *Org. Lett* 2007;9:4339. [PubMed: 17854201]; qBraun M, Meier T, Laicher F, Meletis P, Fidan M. *Adv. Synth. Catal* 2008;350:303.
2. Trost BM, Xu J, Markus R. *J. Am. Chem. Soc* 2007;129:282. [PubMed: 17212401]
3. Protocols for the asymmetric synthesis of α -hydroxyketones include: oxidation of enolates, enol ethers or enol esters: aDavis FA, Chen BC. Helmchen G, Hoffmann RW, Mulzer J, Schaumann E. *Houben-Weyl: Methods of Organic Chemistry* 1995;E 21:4497. Georg Thieme Verlag Stuttgart; bDavis FA, Chen BC. *Chem. Rev* 1992;92:919., and references therein cHashiyama T, Morikawa K, Sharpless KB. *J. Org. Chem* 1992;57:5067.; dMorikawa K, Park J, Andersson PG, Hashiyama T, Sharpless KB. *J. Am. Chem. Soc* 1993;115:8463.; eZhu Y, Tu Y, Yu H, Shi Y. *Tetrahedron Lett* 1998;39:7819.; aminoxylation of ketones: fMoriyama N, Yamamoto H. *J. Am. Chem. Soc* 2003;125:6038. [PubMed: 12785817]; gBøgevig A, Sundén H, Córdova A. *Angew. Chem., Int. Ed* 2004;43:1109.; hHayashi Y, Yamaguchi J, Sumiya T, Shoji M. *Angew. Chem., Int. Ed* 2004;43:1112.; benzoin-type reactions: iEnders D, Kallfass U. *Angew. Chem., Int. Ed* 2002;41:1743.; jLinghu X, Potnick JR, Johnson JS. *J. Am. Chem. Soc* 2004;126:3070–3071.3071 [PubMed: 15012135]; reduction of 1,2-diketones: kKoike T, Murata K, Ikariya T. *Org. Lett* 2000;2:3833. [PubMed: 11101431]; oxidation of 1,2-diols: lAdam W, Fell RT, Saha-Moller CR, Zhao C-G. *Tetrahedron: Asymmetry* 1998;9:397.; the rearrangement of α -tertiary hydroxyaldehydes: mOoi T, Saito A, Maruoka K. *J. Am. Chem. Soc* 2003;125:3220. [PubMed: 12630870]; nOoi T, Ohmatsu K, Maruoka K. *J. Am. Chem. Soc* 2007;129:2410. [PubMed: 17290989]; and alkylation: oAndrus MB, Hicken EJ, Stephens JC, Bedke DK. *J. Org. Chem* 2005;70:9470. [PubMed: 16268622]
4. The absolute configuration was assigned by the NMR study of the corresponding O-methylmandelate ester. Trost BM, Belletire JL, Godleski S, McDougal PG, Balkovec JM, Baldwin JJ, Christy ME, Ponticello GS, Varga SL, Springer JP. *J. Org. Chem* 1986;51:2370.
5. Observed in the preparation of substrates. See Supporting Information for details.

Table 2

Reaction Scope.^a

entry

substrate

1

2

5d ($\text{R}_3 = \text{Piv}$)3^b

4

7d ($\text{R}_3 = \text{Piv}$)

entry

substrate

5

9

6

11b

entry

substrate

7

11d ($\text{R}_3 = \text{Piv}$)

8

9

13

entry

substrate

 10^c

entry

substrate

11

12^c

entry

substrate

13

14

entry

substrate

15

^a All reactions were performed on a 0.2 mmol scale at 0.1 M in DME at 23 °C for 16 h, using 2.5 mol% $\text{Pd}_2(\text{dba})_3\text{CHCl}_3$ and 5.5 mol% **Lnaph**; the yields were isolated yields and ee values were determined by chiral HPLC.

^b Reaction was performed at 4 °C.

^c Reaction was performed at 40 °C.

^d Greater than 95/5 dr.