Parasitic infections in HIV-infected patients who visited Seoul National University Hospital during the period 1995-2003

Sang-Mee GUK¹⁾, Min SEO²⁾, Yun-Kyu PARK³⁾, Myoung-Don OH⁴⁾, Kang-Won CHOE⁴⁾, Jae-Lip KIM¹⁾,
Min-Ho CHOI¹⁾, Sung-Tae HONG¹⁾ and Jong-Yil CHAI¹⁾*

¹⁾Department of Parasitology and Tropical Medicine, Seoul National University College of Medicine, and
Institute of Endemic Diseases, Seoul National University Medical Research Center, Seoul 110-799,

²⁾Department of Parasitology, College of Medicine, Dankook University, Cheonan 330-714,

³⁾Department of Parasitology, College of Medicine, Inha University, Incheon 400-712,

⁴⁾Department of Internal Medicine, Seoul National University College of Medicine, Seoul 110-744, Korea

Abstract: The prevalence of parasitic infections was investigated in human immunodeficiency virus (HIV)-infected patients (n = 105) who visited Seoul National University Hospital, Seoul, Korea, during the period from 1995 to 2003. Fecal samples were collected from 67 patients for intestinal parasite examinations, and sputum or bronchoalveolar lavage samples from 60 patients for examination of *Pneumocystis carinii*. Both samples were obtained from 22 patients. Thirty-three (31.4%) of the 105 were found to have parasitic infections; *Cryptosporidium parvum* (10.5%; 7/67), *Isospora belli* (7.5%; 5/67), *Clonorchis sinensis* (3.0%; 2/67), *Giardia lamblia* (1.5%; 1/67), *Gymnophalloides seoi* (1.5%; 1/67), and *Pneumocystis carinii* (28.3%; 17/60). The hospital records of the 11 intestinal parasite-infected patients showed that all suffered from diarrhea. This study shows that parasitic infections are important clinical complications in HIV-infected patients in the Republic of Korea.

Key words: HIV-infected patients, intestinal parasites, *Cryptosporidium parvum*, *Isospora belli*, *Pneumocystis carinii*, opportunistic parasites

INTRODUCTION

Human immunodeficiency virus (HIV) infection is an extremely serious worldwide problem. As of 2004, about three thousand Koreans were reported to be infected with HIV by the National Institute of Health,

Republic of Korea. Since the beginning of the HIV epidemic in the world, opportunistic infections have been recognized as an important clinical complication (Lebbad et al., 2001; Wiwanitkit, 2001; Vajpayee et al., 2003; Zali et al., 2004). In particular, opportunistic protozoans play a major role as etiologic agents of chronic diarrhea in HIV-infected patients. Chronic diarrhea in such patients can profoundly compromise the absorptive function of the small intestine, and cause significant mortality (Escobedo and Nunez, 1999). The protozoan parasites, *Cryptosporidium parvum*, *Isospora belli*, *Cyclospora cayetanensis*, *Entamoeba histolytica*, and *Giardia lamblia* are the causative agents of diarrhea

• Received 7 July 2004, accepted after revision 17 January 2005.

• This study was supported by a grant from the Korea Science and Engineering Foundation [R05-2002-000-00808-0 (2002)] and by the BK21 Human Life Sciences project of the Ministry of Education, Republic of Korea.

*Corresponding author (e-mail: cjy@snu.ac.kr)

among this population (Mohandas et al., 2003). In addition, *Pneumocystis carinii* pneumonia (PCP) remains the most common opportunistic infection associated with acquired immunodeficiency syndrome (AIDS) in developing countries (Wohl et al., 2002; Aderaye et al., 2003).

The detection of specific pathogens that cause severe diarrhea in HIV-infected patients has become a focus of interest, although such attempts have been limited, to date, in the Republic of Korea. Candidiasis and herpes virus infections were common gastrointestinal tract infections among the HIV-infected patients (Choi et al., 1999; Kim et al., 2003), however, no studies have focused specifically on opportunistic parasitic infections. Thus, the present study was undertaken to determine the prevalence of parasitic infections among HIV-infected patients in the Republic of Korea.

MATERIALS AND METHODS

Patients

One hundred and five HIV-infected patients who visited the Seoul National University Hospital, Seoul, Republic of Korea for check-up, and provided fecal, sputum, or bronchoalveolar lavage (BAL) samples, during the period from January 1995 to October 2003, were enrolled in this study. Fecal samples were obtained from 67 patients and sputum or BAL fluid from 60 (22 patients provided both sample types).

Examination of parasitic infections

Fecal samples were immediately fixed in 10% formalin, and processed using the formalin-ether sedimentation technique. The pellet obtained was smeared on a glass slide and observed under a light microscope. In addition, a second smear was made, air-dried, and stained using a modified acid-fast procedure to detect *Cryptosporidium*, *Isospora*, and other coccidian protozoa under a light microscope (x 1,000). Sputum was treated with 0.1 N KOH for 1 hr and washed with phosphate buffered saline. Sputum and BAL samples were centrifuged at 3,000 rpm for 10 min, smeared on a glass slides, and stained with Diff-

Quik (Fisher Scientific, Orangeburg, NY, U.S.A.) to enable *P. carinii* infection diagnosis under a light microscope (x 1,000).

Review of medical records

The medical records of the 11 parasite-positive HIV-infected patients were reviewed; patients' experiences of diarrhea or fever were noted, as were the number of CD4+ cells at the time of the diagnosis of parasitic infections, weight loss, treatment, and effects of the treatment.

RESULTS

The mean age of patients who provided fecal samples was 39.2 ± 10.0 years ($n = 67$; 61 men and 6 women). Sixteen (23.9%) were found to be infected with intestinal (including liver) parasites (Table 1). Of these parasite-positive patients, the majority was men (80.0%) and more than 65% of the 16 were in their thirties and forties. Opportunistic protozoans predominated. The commonest was *C. parvum* (7/67; 10.5%) followed by *I. belli* (5/67; 7.5%) and *G. lamblia* (1/67; 1.5%). Helminthic infections were identified in 3 cases; *Clonorchis sinensis* (2/67; 3.0%) and *Gymnophalloides seoi* (1/67; 1.5%). Fifteen of these 16 positive cases were infected with one parasite species, and the single mixed case was infected with *I. belli* and *G. seoi* (Table 1).

The medical records of 11 parasite-infected patients showed that all or the majority experienced diarrhea (watery, bloody or non-specific) (11/11), weight loss (7/11), and fever (8/11) (Table 2). In 5 patients infected with *C. parvum*, the number of CD4+ lymphocytes was $50/\text{mm}^3$ or lower, whereas in 2 patients infected with *I. belli*, this number was $131/\text{mm}^3$ or higher. One of the two cases with the lowest CD4+ lymphocyte count of $10/\text{mm}^3$, was infected with *G. lamblia*; the other with *C. sinensis*. Three cases of cryptosporidiosis were treated with azithromycin, which provided a temporary effect in 2 cases and no effect in the other.

The mean age of patients subjected to a *P. carinii* examination was 38.9 years ($n = 60$), and 50 were men and 10 women. Of these 60 patients, 17 (28.3%) were

Table 1. Prevalence of parasitic infections in HIV-infected patients

Parasite species	No. posit./ No. exam.	Positive rate (%)
Intestinal or liver parasites		
<i>Cryptosporidium parvum</i>	7/ 67	10.5
<i>Iso spor a belli</i>	5/ 67	7.5
<i>Giardia lamblia</i>	1/ 67	1.5
<i>Clonorchis sinensis</i>	2/ 67	3.0
<i>Gymnophalloides seoi</i> ^{a)}	1/ 67	1.5
Subtotal	16/ 67	23.9
<i>Pneumocystis carinii</i> ^{b)}	17/ 60	28.3
Total	33/105 ^{c)}	31.4

^{a)}Mixed-infected with *I. belli*.

^{b)}Three of 17 were mixed-infections with *C. parvum* or *G. lamblia*.

^{c)}Fecal examination only; 45 patients, *P. carinii* examination only; 38 patients, Both examinations: 22 patients.

infected with *P. carinii* (Table 1), with a mean age of 28.3 years and more than 80% were men. Most of them (12/17) suffered from respiratory difficulties or pneumonia.

DISCUSSION

Opportunistic infection continues to be a ubiquitous complication of advanced HIV infections and the one that exacts a heavy toll in terms of morbidity and

mortality (Vajpayee et al., 2003). Despite recent increases in the number of HIV-infected patients in the Republic of Korea, few studies have been performed on the prevalence of opportunistic parasitic infections among those patients. In the present study, the prevalence of intestinal parasites and *P. carinii* infections among HIV-infected patients who visited Seoul National University Hospital during 1995 and 2003 was found to be high, i.e., 23.9% (16/67) and 28.3% (17/60), respectively. Overall parasite (including *P. carinii*) prevalence was 31.4% (33/105). With regard to intestinal parasites, no prior comparative report has been issued in the Republic of Korea, with the exception of two papers which focused on *C. parvum* and *I. belli* (Choi et al., 1999) or on *Entamoeba histolytica* (Oh et al., 2000). Oocysts of *C. parvum* or *I. belli* were detected in 3.5% (4/115) (Choi et al., 1999) and invasive amebiasis in 3.1% (6/195) of HIV-infected patients (Oh et al., 2000). With regard to PCP, our results were higher than the results previously reported in the Republic of Korea; 10.4% (18/173) (Oh et al., 1999) or 15.9% (28/176) (Kim et al., 2003) of HIV-infected patients.

Cryptosporidium parvum, an intracellular protozoan which parasitizes the intestinal mucosa and elicits transient diarrhea in healthy individuals, is also one of the most important opportunistic pathogens (Chai et al., 2001; Guk et al., 2003). In immunocompromised

Table 2. Clinical records of 11 HIV-infected patients positive for intestinal or liver parasites

Patient code	Parasite	No. CD4+ lymphocytes ^{a)}	Diarrhea	Weight loss	Fever ^{c)}	Tx for parasites ^{d)}	Effect of Tx.	Other infections
1	<i>C. parvum</i>	11/mm ³	watery	17 kg/24 m ^{b)}	+	azithromycin	temporary	TB ^{e)}
2	<i>C. parvum</i>	50/mm ³	watery	17 kg/6 m	+	azithromycin	no effects	PCP ^{f)}
3	<i>C. parvum</i>	NR ^{g)}	watery	10 kg/4 m	+	anti-TB drug	temporary	none
4	<i>C. parvum</i>	50/mm ³	watery	12 kg/10 d	+	not treated	-	PCP/candidiasis
5	<i>C. parvum</i>	50/mm ³	watery	5 kg/5 m	-	azithromycin	temporary	none
6	<i>G. lamblia</i>	10/mm ³	non-specific	NR	+	not treated	-	TB/PCP
7	<i>I. belli</i>	200/mm ³	watery	16 kg/12 m	-	bactrim	effective	candidiasis
8	<i>I. belli</i>	131/mm ³	non-specific	12 kg/24 m	+	SXT ^{h)}	temporary	candidiasis
9	<i>I. belli</i>	NR	non-specific	NR	NR	bactrim	effective	candidiasis
10	<i>C. sinensis</i>	175/mm ³	bloody	none	+	praziquantel	effective	-
11	<i>C. sinensis</i>	10/mm ³	non-specific	NR	+	praziquantel	effective	candidiasis

^{a)}/mm³ blood; counted at the time of fecal examination, ^{b)}m: months, d: days, ^{c)}Most showed mild fever, ^{d)}Tx: treatment,

^{e)}TB: tuberculosis, ^{f)}PCP: *P. carinii* pneumonia, ^{g)}NR: no record, ^{h)}SXT: symptomatic treatment.

patients, this protozoan can cause severe and life-threatening diarrhea, which is often fatal (Chai et al., 2001). In the present study, most HIV-infected patients infected with *C. parvum* experienced severe watery diarrhea and weight loss. The prevalence of cryptosporidiosis is variable (> 0.5%-35% or higher) and depends on geography and the environment (Chai et al., 2001). Previous studies have indicated that CD4⁺ cells are important for host defense mechanisms against cryptosporidiosis (Guk et al., 2003). In 4 of 5 HIV-infected patients infected with *C. parvum*, CD4⁺ cell numbers were lower than in patients infected with *I. belli*. Moreover, no drug therapy has been shown to effectively treat cryptosporidiosis. It is of interest to note that 2 of 3 patients treated with azithromycin showed temporary effects.

Isospora belli had become more prominent in recent years because of its prevalence as an opportunistic pathogen in AIDS patients (Sorvillo et al., 1995). Nevertheless, no report is available on the prevalence of isosporiasis in the Republic of Korea. Moreover, the actual rate of this infection in immunocompetent individuals and HIV-infected patients is likely to be underestimated, because of the asymptomatic shedding of oocysts, treatment with trimethoprim-sulfamethoxazole (TMP-SMX) for other infections, a low number of oocysts excreted, and an inefficiency of feces concentration and oocyst observation (Escobedo and Nunez, 1999). Epidemiological studies on the prevalence of isosporiasis are required in the Republic of Korea.

Pneumocystis carinii, which is now more frequently classified as a fungus, is a ubiquitous organism that can cause interstitial pneumonia, and one of the common opportunistic parasites associated with AIDS in developing countries (Wohl et al., 2002). In the Republic of Korea, it was reported that 10.4% of 173 HIV-infected patients (Oh et al., 1999) and 15.9% of 176 patients (Kim et al., 2003) developed PCP. In our study, samples for the diagnosis of *P. carinii* infection were collected from HIV-infected and suspected to suffer from PCP. Most of them suffered from cough, dyspnea, sputum, and other respiratory symptoms. Therefore, a higher proportion (28.3%) was found to

be infected in the present study. With regard to the treatment of PCP, primary prophylaxis with TMP-SMX was found to induce hypersensitivity in as many as 25% of HIV-infected patients (Fischl et al., 1988). However, no cases of hypersensitivity reactions to TMP-SMX have been reported in 46 Korean HIV-infected patients infected with *P. carinii*. The reason was suggested to be the fast speed of an acetylator status of Koreans (Oh et al., 1999). In the present study, we diagnosed *P. carinii* infection by Diff-Quik staining of sputum or BAL samples. Further studies are required to confirm the prevalence of this fungal infection using serology or other techniques.

The cause of death was checked in 11 HIV-infected patients, but we could find apparent causes of death in only 7 patients. Two HIV-infected patients infected with *P. carinii* died of sepsis and gastrointestinal bleeding, and three patients infected with *C. parvum* (one of the three was co-infected with *P. carinii*) died of pneumonia and liver abscess. Other patients infected with *C. sinensis* or *I. belli* died of pneumonia or disseminated tuberculosis.

The prevalence of helminthic infections in the present study was 4.5% (3/67) which included 2 clonorchiasis and 1 gymnohalloidiasis cases. These two species are important food-borne parasites in the Republic of Korea. In particular, *G. seoi* is transmitted by oysters, which are commonly eaten raw (Chai et al., 2003). This fluke parasitizes the small intestines of man and animals, and can induce gastrointestinal problems and varying degrees of indigestion (Chai et al., 2003).

The results of the present study demonstrate that opportunistic protozoan infections are prevalent among Korean HIV-infected patients and they represent an important source of clinical complications among these patients.

REFERENCES

- Aderaye G, Bruchfeld J, Olsson M, Lindquist L (2003) Occurrence of *Pneumocystis carinii* in HIV-positive patients with suspected pulmonary tuberculosis in Ethiopia. *AIDS* 17: 435-440.

- Chai JY, Choi MH, Yu JR, Lee SH (2003) *Gymnophalloides seoi*: a new human intestinal trematode. *Trends Parasitol* **19**: 109-112.
- Chai JY, Kim NY, Guk SM, Park YK, Seo M, Han ET, Lee SH (2001) High prevalence and seasonality of cryptosporidiosis in a small rural village occupied predominantly by aged people in the Republic of Korea. *Am J Trop Med Hyg* **65**: 518-522.
- Choi HJ, Park SW, Kim HB, Kim NJ, Shin DH, Oh MD, Jung HC, Song IS, Choe KW (1999) The gastrointestinal infections in Korean patients with human immunodeficiency virus infection. *Korean J Gastroenterol* **33**: 482-488.
- Escobedo AA, Nunez FA (1999) Prevalence of intestinal parasites in Cuban acquired immunodeficiency syndrome (AIDS) patients. *Acta Trop* **72**: 125-130.
- Fischl MA, Dickinson GM, La Voie L (1988) Safety and efficacy of sulfamethoxazole and trimethoprim chemoprophylaxis for *Pneumocystis carinii* pneumonia in AIDS. *JAMA* **259**: 1185-1189.
- Guk SM, Yong TS, Chai JY (2003) Role of murine intestinal intraepithelial lymphocytes and lamina propria lymphocytes against primary and challenge infections with *Cryptosporidium parvum*. *J Parasitol* **89**: 270-275.
- Kim JM, Cho GJ, Hong SK, Chang KH, Chung JS, Choi YH, Song YG, Huh A, Yeom JS, Lee KS, Choi JY (2003) Epidemiology and clinical features of HIV infection/AIDS in Korea. *Yonsei Med J* **44**: 363-370.
- Lebbad M, Norrgren H, Naucler A, Dias F, Andersson S, Linder E (2001) Intestinal parasites in HIV-2 associated AIDS cases with chronic diarrhoea in Guinea-Bissau. *Acta Trop* **80**: 45-49.
- Mohandas K, Sehgal R, Sud A, Malla N (2002) Prevalence of intestinal parasitic pathogens in HIV-seropositive individuals in Northern India. *Jpn J Infect Dis* **55**: 83-84.
- Oh MD, Lee K, Kim E, Lee S, Kim N, Choi H, Choi MH, Chai JY, Choe K (2000) Amoebic liver abscess in HIV-infected patients. *AIDS* **14**: 1872-1873.
- Oh MD, Park SW, Kim HB, Kim US, Kim NJ, Choi HJ, Shin DH, Lee JS, Choe K (1999) Spectrum of opportunistic infections and malignancies in patients with human immunodeficiency virus infection in South Korea. *Clin Infect Dis* **29**: 1524-1528.
- Sorvillo FJ, Lieb LE, Seidel J, Kerndt P, Turner J, Ash LR (1995) Epidemiology of isosporiasis among persons with acquired immunodeficiency syndrome in Los Angeles County. *Am J Trop Med Hyg* **53**: 656-659.
- Vajpayee M, Kanswal S, Seth P, Wig N (2003) Spectrum of opportunistic infections and profile of CD4+ counts among AIDS patients in North India. *Infection* **31**: 336-340.
- Wiwanitkit V (2001) Intestinal parasitic infections in Thai HIV-infected patients with different immunity status. *BMC Gastroenterol* **1**: 3.
- Wohl AR, Simon P, Hu YW, Duchin JS (2002) The role of person-to-person transmission in an epidemiologic study of *Pneumocystis carinii* pneumonia. *AIDS* **16**: 1821-1825.
- Zali MR, Mehr AJ, Rezaian M, Meamar AR, Vaziri S, Mohraz M (2004) Prevalence of intestinal parasitic pathogens among HIV-positive individuals in Iran. *Jpn J Infect Dis* **57**: 268-270.