Published in final edited form as:

Glia. 2008 April 15; 56(6): 646–658. doi:10.1002/glia.20643.

Activated Notch1 maintains the phenotype of radial glial cells and promotes their adhesion to laminin by upregulating nidogen

Hedong Li^{1,*}, Yu-Wen Chang¹, Kriti Mohan¹, Hui-Wen Su¹, Christopher L. Ricupero¹, Ajoeb Baridi^{1,2}, Ronald P. Hart¹, and Martin Grumet^{1,*}

¹Department of Cell Biology and Neuroscience, and W. M. Keck Center for Collaborative Neuroscience, Rutgers, State University of New Jersey, 604 Allison Rd, Piscataway, NJ 08854-8082

²Department of Pharmacology & Anatomy, Rudolf Magnus Institute of Neuroscience, Utrecht University, Utrecht, The Netherlands

Abstract

Radial glia are neural stem cells that exist only transiently during CNS development where they serve as scaffolds for neuronal migration. Their instability makes them difficult to study and therefore we have isolated stabilized radial glial clones from E14.5 cortical progenitors (e.g. L2.3) after expression of v-myc. Activated Notch1 intracellular region (actNotch1) promotes radial glia in the embryonic mouse forebrain (Gaiano et al. 2000) and when it was introduced into E14.5 cortical progenitors or radial glial clone L2.3, the cells exhibited enhanced radial morphology and increased expression of the radial glial marker BLBP. A representative clone of L2.3 cells expressing actNotch1 called NL2.3–4 migrated more extensively than L2.3 cells in culture and in white matter of adult rat spinal cord. Microarray and RT-PCR comparisons of mRNAs expressed in these closely related clones showed extensive similarities but differed significantly for certain mRNAs including several cell adhesion molecules. Cell adhesion assays demonstrated significantly enhanced adhesion to laminin of NL2.3–4 by comparison to L2.3 cells. The laminin binding protein nidogen was the most highly induced adhesion molecule in NL2.3–4, and immunological analyses indicated that radial glia synthesize and secrete nidogen. Adhesion of NL2.3–4 cells to laminin was inhibited by anti-nidogen antibodies and required the nidogen binding region in laminin, indicating that nidogen promotes cell adhesion to laminin. The combined results indicate that persistent expression of activated Notch1 maintains the phenotype of radial glial cells, inhibits their differentiation, and promotes their adhesion and migration on a laminin/nidogen complex.

Keywords

brain development; neural stem cell; progenitor; laminin; A2B5

Introduction

Radial glia exhibit unique properties including serving as scaffolds for neuronal migration during neurogenesis (Hatten 1999; Rakic 1990). They lose these properties with differentiation, for example after transplantation of radial glia into adult CNS tissues they differentiate into astrocytes (Cao et al. 2002; Hasegawa et al. 2005). Prolonging radial glial

*Correspondence should be addressed to: Dr. Martin Grumet or Hedong Li, W. M. Keck Center for Collaborative Neuroscience, Rutgers, State University of New Jersey, 604 Allison Rd, Piscataway, NJ 08854-8082, Tel. 732/445-6577, Fax. 732/445-2063, **e-mails:** E-mail: mgrumet@rci.rutgers.edu or E-mail: hedong@rci.rutgers.edu.

phenotypes may be a useful strategy that takes advantage of their beneficial properties to promote functional recovery after spinal cord injury (Hasegawa, 2005). Recent studies indicate that radial glia in the developing CNS are neural stem cells (NSC) but very few persist in the adult brain and spinal cord (Alvarez-Buylla et al. 2002; Anthony et al. 2004; Kriegstein and Gotz 2003).

Radial glia are NSC or NPC that are challenging to maintain and study in vitro because they tend to differentiate. Although progress has been made in isolating NSC/NPC from embryonic stem cells and maintaining them in culture in defined media conditions (Conti et al. 2005), approaches are still needed to control their behavior and fate, particularly after transplantation into the CNS where they are subjected to multiple signals that induce differentiation. Towards that goal, clones of embryonic brain cells have been isolated after introduction of v-myc (Villa et al. 2000). We isolated two clones (L2.3 and RG3.6) from embryonic rat brain cortices that share properties with NSC and radial glia (Hasegawa et al. 2005; Li et al. 2004) and RG3.6 cells promote functional recovery following transplant into the injured rat spinal cord (Hasegawa et al. 2005). Upon extended passage in culture, these cells transition to express immunological markers recognized by monoclonal antibodies A2B5 and 4D4 that are found on glial restricted precursors (Li et al. 2004; Li and Grumet 2007; Noble et al. 2003). Developmental studies showed that radial glia appeared as early as E9-10 in the CNS while the A2B5 and 4D4 markers for glial restricted precursors appeared later, suggesting that glial restricted precursors may be derived from radial glia as they become restricted and develop into neurons and glia (Li et al. 2004). Other cell type specific markers including nestin and vimentin are expressed in NSC and NPC but their expression is not restricted only to these cells. BLBP is restricted to radial glia during development, but it persists in astroglia and adult NPC (Feng et al. 1994; Hartfuss et al. 2001). NSC in culture also expresses BLBP (Conti et al. 2005). Although the function of BLBP is poorly understood, its expression is regulated by Notch1, which is transiently expressed in radial glia (Anthony et al. 2005; Gaiano et al. 2000).

Notch is a transmembrane receptor that mediates lateral inhibition by interacting with ligands on apposing cells including delta (Lai 2004). In response to ligand binding, Notch undergoes proteolysis and the resulting Notch intracellular domain translocates into the nucleus. Notch intracellular domain acts as a transcriptional activator that turns on downstream genes including BLBP, and the transcriptional repressors Hes1 and Hes5 that suppress neurogenesis (Yoon and Gaiano 2005). Thus, Notch signaling plays important roles in cell fate determination during CNS development. ActNotch1 promotes the radial glial phenotype in vivo (Gaiano et al. 2000) and maintains NPC/NSC in an undifferentiated state in the mammalian CNS (Hitoshi et al. 2002).

The morphological integrity of radial glial cells in the developing forebrain has been thought to be influenced by the cortical pial basement membrane where the endfeet of radial glial cells are anchored (Halfter, 2002). Basement membranes are thin layers of extracellular matrix, containing collagen IV and XVIII, laminin and a linker protein nidogen (Yurchenco and Schittny 1990). Nidogen plays an important role in maintaining basement membranes structure. Mice with a targeted deletion of the nidogen-binding site of laminin g1 showed disrupted pial membranes during cortical histogenesis and radial glial cells with retracted endfeet suggesting its importance in cortical development (Halfter, 2002). These mice also exhibited abnormal neuronal migration, which is probably due to the deformed radial glial morphology. The main source of nidogen and laminin proteins in the pial membranes is thought to be the meningeal cells (Sievers et al. 1994).

In the present study, we found that activated Notch1 promoted radial morphology in NPC in culture, maintained BLBP expression, and inhibited cell differentiation. Expression of

activated Notch1 in a subclone of L2.3 called NL2.3-4 yielded cells with an elongated radial morphology in culture and after transplantation into the spinal cord that resisted differentiation. These radial glial-like cells expressed nidogen, which promoted their adhesion to laminin. In addition, primary radial glia expressed and secreted nidogen.

Materials and Methods

Cell culture and differentiation

Generation and cultures of radial glial clone L2.3 have been described previously (Li, 2004). The same culture protocol was used for NL2.3-4 clone and primary cortical radial glia. Briefly, culture medium contained DMEM/F12 (Invitrogen) supplemented with 25 mM glucose (Sigma), 2 mM glutamine (Invitrogen), penicillin/streptomycin (Invitrogen), 10 ng/ml FGF2 (BD Biosciences), 2 µg/ml heparin (Sigma) and 1x B27 (Invitrogen). Cells were propagated as neurospheres and passaged by mild trypsinization (0.025% for 5 min) every 3 days. For differentiation, cells were cultured on laminin-coated coverslips in FGF2 containing serum-free medium for 1 day, then the medium was replaced with culture medium lacking FGF2, including 1% fetal bovine serum (FBS), for 6 days, and the cultures were fixed, and immunostained. For Brefeldin A treatment (5 µg/ml, CalBiochem), cultures were incubated for 90 min at 37°C before fixation.

Retroviral infection and selection of clones

Primary NSC were isolated from E13.5 cortices and maintained in serum free medium as described above. The next day, cells were mechanically dissociated by gently pipeting to yield $1-2 \times 10^6$ cells/ml in serum free medium containing 8 µg/ml polybrene. ActNotch1-GFP or control GFP viruses (Yoon et al. 2004) was added at 1:1000 dilution and incubated with cells for 1 hr on a shaker at 100 rpm. After three washes in serum free medium, the cells were plated in NSC culture medium. GFP signals were observed within 48 hours after infection. Infected cells were passed and plated in culture medium at clonal density (1000 cells / 10-cm dish). After 3 – 4 days, individual neurospheres were picked, expanded and characterized.

Measurement of cellular process lengths

Images of undifferentiated cells stained with nestin were measured using Zeiss Laser Scanning Microscope (LSM) imaging software; cell processes were traced from the beginning to the end using the measurement tool. When there were multiple processes arising from one cell body, only the longest process was measured. Measurements were sorted by range (e.g. 0–20 µm, 20–40 µm, 40–60 µm, etc.) and the number/group were graphed.

Affymetrix Genechip™ analysis and quantitative-RT-PCR (Q-RT-PCR)

Total RNA was purified from neurosphere cultures using RNeasy kit (Qiagen). Samples (n=3 per group) were sent to the Bionomics Research and Technology Center, Rutgers University, for Affymetrix processing. Results were normalized using RMA, fit to a linear model using the *affylnGUI* in Bioconductor (Smyth 2004; Wettenhall et al. 2006; Wettenhall and Smyth 2004) and tested for significance at a 5% false discovery rate (FDR) to control for multiple measurements error and at least 2-fold mean difference between groups (Supplemental Table 1). Gene ontology analysis was performed using GeneSpring (Agilent, Supplemental Table 2).

For PCR analyses, one microgram was reverse-transcribed into cDNA using oligo-dT primer and SuperScript II reverse transcriptase (Invitrogen). Quantitative RT-PCR was performed as described previously (Li et al. 2003) using an ABI PRISM 7900HT Sequence Detection System (Applied Biosystems). Beta-actin was used to normalize the expression levels of each sample. Primers for detecting genes are listed in Table I.

Cell adhesion assay on laminin

Dissociated single cell suspensions (10^6 cells/ml) were incubated on laminin-1 (20 μ g/ml, Invitrogen) coated spots (5000 cells per spot) for 40 min at 37°C, washed with PBS, fixed and stained with 1 μ g/ml propidium iodide (Sigma). Blocking antibody anti-nidogen is a rat IgG from Chemicon. Laminin-2 (Smirnov et al. 2002) and the laminin-2-NS with a deletion of the nidogen-binding site within laminin α 1 chain (Halfter et al. 2002) were generously provided by Dr. Peter Yurchenco and used at 20 μ g/ml. Attached cells per spot were counted and average cell numbers were calculated from triplicate spots for each condition.

Neurosphere spreading assay on laminin

Cultured neurospheres labeled with DAPI were incubated at 37°C on laminin-coated substrates. Pictures were taken soon after the neurosphere attachment (30 min) and when cells migrated out of neurospheres (2 hours) at the same positions. DAPI images were used to quantify the neurosphere spreading area at 30 min (a1) and at 2 hours (a2). Spreading area ratios ((a2-a1)/a1) were calculated.

Transplantation into the spinal cord and cell migration

NL2.3-4 and L2.3 cells were labeled with CellTracker CFDA-SE (Molecular Probes) according to manufacturer's protocol, and 200,000 cells were transplanted into adult Sprague Dawley rat spinal cords at T9 and T13; animals were sacrificed after 3, 14 or 28 days, and spinal cords were sectioned for histological analysis as described (Hasegawa et al. 2005).

Western blot analysis

Cultured cells were harvested in SDS lysis buffer, and heated at 95°C for 5 min. Proteins were separated on 10% SDS-PAGE and transferred onto PVDF membranes and immunoblotted with anti-GFP (1:500, mouse IgG, Chemicon), anti-nidogen (1:200, (McKee et al. 2007)) or anti-NCAM (1:50, rabbit IgG, Grumet lab), followed by horseradish peroxidase (HRP)-conjugated secondary antibodies (1:5000, Jackson lab). The blots were developed using ECL plus detection system (GE Healthcare Amersham). Anti-actin (1:1000, rabbit IgG, Sigma) was used to normalize the sample loading.

Immunocytochemistry and immunohistochemistry

Cultured cells were fixed with 4% paraformaldehyde and immunostained as described (Li and Grumet 2007). The primary antibodies used were monoclonal mouse IgMs: 4D4 (neat, a gift from Dr. Kaprielian's lab) (Liu et al. 2002), A2B5 (1:200, Chemicon), 5A5 (1:1, DSHB) and RC1 (1:5, DSHB); monoclonal mouse IgGs: anti-vimentin (1:10, DSHB), anti-nestin (1:50, DSHB), GalC (1:50, McKinnion's lab), TuJ1 (1:500, Covance); polyclonal rabbit IgGs: anti-BLBP (1:1000, a gift from Dr. Heintz's lab), anti-nidogen (1:200) and anti-GFAP (1:200, DAKO). DAPI (10 μ g/ml, Sigma) was included in the secondary antibody incubations to label nuclei.

In acquiring the fluorescence intensity of BLBP, confocal images were analyzed with the Carl Zeiss LSM 510 program. The appropriate channels were chosen (GFP chnl and BLBP chn2). The images were opened in the 'histogram display mode' where both 'area' and 'colocalization' functions were chosen. The cells colocalising both markers were individually selected by tightly drawing a borderline around the cells with the 'close polyline drawing mode' function. The 'show table' function showed all the parameters that had been measured. The 'area' and 'mean intensity' parameters were chosen and multiplied to finally produce the Fluorescence Intensity of BLBP depicted in the figure (**, $p < 0.01$).

Results

Activated Notch1 (actNotch1) promotes radial glia phenotypes in culture and inhibits differentiation

To determine whether actNotch1 promotes radial glia phenotype in vitro, we used a GFP version of the same viral construct to infect L2.3 cells (Li et al. 2004) and cortical cultures that are derived from E13.5 rat forebrains. The retrovirus encodes for Notch1 intracellular domain fused with GFP, thus the Notch infected cells were easily recognized by their green nuclei. Most actNotch1 infected L2.3 cells exhibited bipolar radial morphology as shown by nestin immunostaining (Fig. 1A, arrow). Notch induces expression of BLBP (Anthony et al. 2005; Gaiano et al. 2000) and we confirmed that actNotch1 induced BLBP expression in actNotch1-GFP cells by comparison to GFP infected controls (Fig. 1A). The percentage of BLBP+ infected cells was ~6 fold higher with actNotch1-GFP than with GFP alone (Fig. 1B). Similarly, in E13.5 cortical cultures, we found that actNotch1-GFP infected cells exhibited strong BLBP expression (Fig. 1C and 1D) and some of these cells had very long bipolar processes resembling radial glia (Fig. 1C, arrow). Other actNotch1-GFP cells had flat shapes (Fig. 1C, arrowheads) suggesting they may already have begun to differentiate and actNotch1 apparently did not revert them into radial glia. The expression of BLBP protein, measured by relative fluorescence intensity, was much higher after actNotch1-GFP infection than in GFP controls (Fig. 1D).

We have shown that the fate of radial glial cells becomes restricted during forebrain development as they acquire the glial restricted precursor markers A2B5 and 4D4 (Li et al. 2004). To determine whether Notch signaling plays a role in radial glial cell fate restriction, we examined 4D4 expression. Compared to GFP infected cultures where 90.7% of cells were positive for 4D4, actNotch1-GFP+ cells showed little or no expression of 4D4 in E13.5 cortical cells (Fig. 2). Similar results were also observed in radial glial clone L2.3 but ~20% of cells continued to express 4D4 typically at very low levels (Fig. 2) while the cells that did not express actNotch1 stained strongly for 4D4. These data suggest that Notch signaling maintains the radial glial phenotype and prevents, or at least delays, differentiation towards glial phenotypes.

Isolation and characterization of a Notch1 expressing radial glial-like clone, NL2.3-4, from NSC/NPC clone L2.3

To study the effect of Notch on radial glia in more detail, we isolated clones from actNotch1 expressing L2.3 cells. Five out of eight GFP+ clones showed radial morphology on laminin substrates. Most of them (7 out of 8) showed strong BLBP expression, an indication of active Notch signaling. One clone, NL2.3-4 was characterized further because of its homogeneous BLBP expression, radial morphology and absence of 4D4. Almost 100% of NL2.3-4 cells exhibited GFP in their nuclei and BLBP staining in their cytoplasm, whereas only ~20% of L2.3 cells expressed BLBP (Fig. 3A, 3B). All cells in both clones showed nestin expression as expected for NPC or NSC (Fig. 3B). However, 4D4 expression was absent in NL2.3-4 cells, while ~25% of L2.3 cells showed 4D4 immunoreactivity. Thus, persistent expression of actNotch1 blocked expression of 4D4.

Further comparison of NL2.3-4 to its parental clone L2.3 revealed differences in cell morphology and differentiation. On laminin substrate and in the presence of FGF2 (10 ng/ml), NL2.3-4 exhibited longer bipolar processes than L2.3 cells that were as much as 20 times as long as the cell body diameter, as revealed by both differential interference microscopy (DIC) and nestin immunostaining (Fig. 3A, 3C). Upon FGF2 withdrawal, L2.3 cells were multipotential giving rise to small numbers of neurons (TuJ1+) and oligodendrocytes (GalC +), and primarily to astrocytes (GFAP+) (Fig. 4 and Li, 2004). In contrast, NL2.3-4 cells responded to express GFAP, but markers for other neural cell types were not detected (Fig.

4A, 4B). Interestingly, the GFAP+ NL2.3-4 cells retained their bipolar elongated morphologies suggesting that their differentiation was incomplete (Fig. 4A), however, both L2.3 and NL2.3-4 expressed S100 β upon differentiation (data not shown). The combined results suggest that actNotch1 inhibits cell differentiation in neural cells.

NL2.3-4 cells migrate more robustly than L2.3 in normal rat spinal cord and resist differentiation

The migratory property of NL2.3-4 cells was evaluated in rat spinal cord (Hasegawa et al. 2005; Hormigo et al. 2001). NL2.3-4 and L2.3 cells labeled with CFDA-SE were transplanted into adult rat normal spinal cord and the migration pattern of the cells was examined after 3 days. NL2.3-4 cells migrated greater distances in spinal cord white matter than L2.3 cells and differences in rostro-caudal migration (X) were measured across serial sagittal sections (Fig. 5A, 5B; 3 mm for NL2.3-4 and 0.8 mm for L2.3). Analysis of fluorescence in serial sections yielded 3 dimensional images of the transplants that had relatively similar volumetric measures for NL2.3-4 ($11.67 \times 10^6 \mu\text{m}^3$) and L2.3 ($11.31 \times 10^6 \mu\text{m}^3$), suggesting at least at 3 days after transplantation there was no dramatic difference in graft survival. In addition, the asymmetry in the migration was confirmed by calculating the ratio of longitudinal/lateral migration in the spinal cord (X/Y, Fig. 5D). The maximal migration of NL2.3-4 cells increased up to 13.8 mm at 4 weeks after transplantation in spinal cord white matter (Fig. 5E). Higher magnification revealed nestin positive bipolar NL2.3-4 cells in spinal cord white matter even 4 weeks after transplantation. Detection of green labeled nuclei in the radial glial-like cells (Fig. 5F, 5G) indicated the persistence of actNotch1-GFP expression but we did not detect GFAP in these cells suggesting they did not exhibit astroglial differentiation. In contrast, NPC without (Cao et al. 2001) or with v-myc (Hasegawa et al. 2005) showed more extensive differentiation (e.g. GFAP expression) 2–4 weeks after transplantation. The results suggest that persistence of actNotch1-GFP expression prevents differentiation of these cells in vivo.

Microarray analysis revealed elevated adhesion molecules in NL2.3-4 vs. L2.3

Considering differences between NL2.3-4 and L2.3 cells in morphology and migration, we compared their patterns of gene expression using Affymetrix rat genome chip, rat 230-2. Among more than 30,000 transcripts detected on the chip, 814 were significant at a 5% FDR with at least a 2-fold mean difference in expression between NL2.3-4 and L2.3 (Supplemental Table 1). Several known down-stream Notch target genes were up-regulated in NL2.3-4 by comparison to L2.3, including BLBP (Fabp7) and Hes5 (but not Hes1) (de la Pompa et al. 1997), confirming that Notch signaling was active in the NL2.3-4 cells. In addition, the radial glial markers BLBP, GLAST, pax6, Tenascin C, nestin and vimentin (Bartsch et al. 1992; Gotz et al. 1998; Hartfuss et al. 2001; Heins et al. 2002) were up-regulated in NL2.3-4 cells (Fig. 6A and Supplemental Table 1). CD-9, which is found in NSC (Klassen et al. 2001) was also up-regulated, but prominin-1 (CD133) (Pfenninger et al. 2007) decreased somewhat (Supplemental Table 1). These in vitro data suggest that actNotch1 signaling promotes Notch-related and radial glial specific genes in vivo (Gaiano et al. 2000).

An unbiased analysis of regulated genes was performed to identify cellular functions affected by actNotch1 expression. The gene ontology (GO) assignments of significantly regulated genes were compared with all genes probed on the array by selecting top-level categories having a hypergeometric p-value of 0.01 or less (Supplemental Table 2). All major GO groupings (biological process, cellular component, and molecular function) presented interpretable functional categories relating to nervous system development. The most significant biological process categories included systems development, nervous system development, cell adhesion, and astrocyte development, supporting our hypothesis that Notch maintains the radial glial phenotype.

We paid particular attention to cell adhesion related genes in this comparison because of the morphological, adhesive and migratory properties of NL2.3-4 cells. In the cell adhesion gene ontology category, 12 genes were found to be at least 2-fold higher in NL2.3-4 while only 2 genes were 2-fold higher in L2.3; some of the genes appeared multiple times including Alcam, CD44 and N-CAM (Fig. 6B). The fact that more adhesion molecules were up-regulated in NL2.3-4 by comparison to L2.3 suggests that NL2.3-4 cells may be more adhesive. The differential expression of several of these genes was confirmed by Q-RT-PCR analysis (Fig. 6C). In addition, up-regulation of nidogen and NCAM proteins was confirmed by western blotting (Fig. 6C insert).

NL2.3-4 cells adhere and spread better on laminin than L2.3

NL2.3-4 cells exhibited robust adhesion to laminin and several of the genes that are most highly upregulated in response to actNotch1 expression bind to laminin including nidogen (Fox et al. 1991) and CD44 (Hibino et al. 2004). In a laminin-mediated cell adhesion assay, we detected a significant difference in cellular adhesion between NL2.3-4 and L2.3 cells within 40 min incubation (Fig. 7A, 7B, 7C). The NL2.3-4 cells adhered better, and they were more migratory on laminin substrates. In a neurosphere spreading assay, NL2.3-4 cell bodies and processes extended longer distances on laminin than L2.3 cells (Fig. 7D, E). Using DAPI fluorescence to measure the spreading area ratio, we found that NL2.3-4 cell bodies migrated significantly greater distances than L2.3 cells on laminin substrate (Fig. 7F).

Nidogen mediates enhanced cell adhesion of NL2.3-4 on laminin

To identify adhesion molecules expressed by NL2.3-4 cells we analyzed their potential involvement in mediating enhanced adhesion of these cells on laminin substrate. Integrins are classic receptors for laminin substrate (Reichardt and Tomaselli 1991; Yurchenco and Schittny 1990). The major laminin receptor, integrin $\beta 1$, did not show significant differences between NL2.3-4 and L2.3 by Q-RT-PCR analysis (Fig. 6C). CD44 is a hyaluronic acid (HA) binding protein that serves as a receptor for laminin (Halfter et al. 2002; Yurchenco and Schittny 1990), however, anti-CD44 blocking antibody (Frank et al. 2005) did not block cell adhesion of NL2.3-4 on laminin at concentrations as high as 200 $\mu\text{g/ml}$ (data not shown). Nidogen is a major component of basement membranes that plays a role in maintaining its integrity by binding to laminin $\alpha 1$ subunit (Halfter et al. 2002; Yurchenco and Schittny 1990). Anti-nidogen inhibited NL2.3-4 cell adhesion on laminin by about 50% compared with no antibody and rat IgG controls (Fig. 8A–D). Adhesion assays using recombinant laminin-2 protein (LM-2) that bears a mutation in its nidogen binding site, called LM-2-NS (Halfter et al. 2002), showed a 5-fold reduction in cell adhesion on LM-2-NS substrates compared to that on wild type LM-2 (Fig. 8E). In addition, NL2.3-4 cells adhered poorly when cultured directly on recombinant nidogen (Fig. 8E), suggesting that nidogen potentiates cell adhesion to laminin rather than acting as a ligand by itself. When the cultures were immunostained for nidogen, there was reactivity on substrates coated with laminin (EHS) and LM-2 but not on those coated with LM-2-NS, which lacks the nidogen-binding region. This effect was more robust after overnight culture (Fig. 8F–H), indicating that the secreted nidogen can bind to laminin on the culture substrate.

E14.5 cortical radial glial cells, but not neurons, secrete nidogen in culture

To explore the biological relevance of elevated nidogen expression by actNotch1 we analyzed nidogen expression in embryonic cortical radial glial cell cultures where Notch signaling is active (Gaiano et al. 2000). Using a polyclonal anti-nidogen antibody, we found that nidogen localized at the pial surface and around blood vessels as reported (Halfter et al. 2002). We also detected some very weak anti-nidogen staining along nestin+ radial glial fibers in E14.5 spinal cord and in primary cultures (data not shown), but it was unconvincing. We considered that it

might be difficult to detect nidogen in the cell because it was synthesized and released. When the cultures were treated with Brefeldin A (5 μ g/ml, a Golgi apparatus blocker) for 90 min to prevent secretion of proteins, nidogen protein was found to accumulate in radial glial cells that were RC1-positive (Fig. 9A). Similarly, nidogen was found in vimentin-positive cells (insert in Fig. 9A). TuJ1-positive neurons did not show nidogen staining (Fig. 9B) suggesting that radial glia but not neuronal precursors express nidogen. Nidogen-positive cells compose a subpopulation of radial glia, defined by expression of the radial glial marker RC1, and after Brefeldin A treatment, nidogen was localized in vesicular structures as expected for secreted proteins that are prevented from being released. In contrast to primary radial glia, strong anti-nidogen immunostaining was detected in the NL2.3-4 cells even without Brefeldin A treatment confirming their robust expression of nidogen induced by actNotch1 (Fig. 9D). The results suggest that radial glia upregulate nidogen expression when Notch signaling is active.

Discussion

Transient expression of Notch1 during neural development promotes survival and maintenance of radial glial NSC and suppresses or at least delays their differentiation (Androutsellis-Theotokis et al. 2006; Gaiano et al. 2000; Hitoshi et al. 2002; Tanigaki et al. 2001). In this study, we have focused on the consequences of persistent expression of actNotch1 in NPC and found that stable expressors could only be obtained when actNotch1 was co-expressed with v-myc. Transient upregulation of actNotch1 induced expression of BLBP, promoted the radial glia phenotype and suppressed cell differentiation (Anthony et al. 2005; Gaiano et al. 2000). Isolation of a NPC clone stably expressing actNotch1 (NL2.3-4) demonstrated that persistent expression maintained the radial glial phenotype both in vitro and in vivo and inhibited differentiation (Hitoshi et al. 2002).

Persistent expression of actNotch1 in NL2.3 inhibited differentiation induced by withdrawal of FGF2 and addition of serum (Li et al. 2004) (Fig. 4). Although GFAP expression was induced in both NL2.3-4 and L2.3, the L2.3 cells changed morphology to yield cells resembling astrocytes while the NL2.3-4 cells retained their bipolar shape indicating incomplete differentiation (Fig. 4A). Expression of GFAP itself is not sufficient criteria for determining differentiation (Bauer et al. 2007) and it is expressed by adult neural stem cells (Alvarez-Buylla et al. 2002). In contrast to L2.3 cells that tend with increasing passage in culture to decrease expression of BLBP and increase expression of the glial restricted precursor markers A2B5 and 4D4 (Li et al. 2004), NL2.3-4 cells maintained robust BLBP expression and did not express these markers, indicating that actNotch1 can suppress neural cell differentiation. Interestingly, the helix-loop-helix protein Id4 was one of the most highly induced genes by expression of actNotch1 (Supplemental Table 1) and these genes may interact genetically (Liu and Harland 2003). Id4 is required for correct timing of neural differentiation (Bedford et al. 2005) and its overexpression in oligodendrocyte progenitors prevents differentiation (Marin-Husstege et al. 2006). These properties of Id4 suggest it may act to suppress differentiation of the NL2.3-4 cells.

Among the genes that differed most significantly between clones that do or do not express actNotch1 were several adhesion molecules with predicted interactions with laminin including nidogen. The function of nidogen is still poorly understood although it has been known for some time that it binds to laminin (Yurchenco and Schittny 1990). We found that radial glia secrete nidogen and that nidogen potentiated adhesion of these cells to laminin. It has been proposed that nidogen at the pial surface is deposited by meningeal cells (Halfter et al. 2002) but the present results suggest that the radial glia themselves may also contribute to its accumulation. Notch1 activation is typically transient during development suggesting that it may induce transient changes in cell adhesion and signalling. Transient nidogen expression very early during development of radial glia acting synergistically with laminin might provide

functional significance to the enigmatic low levels of laminin that have been detected along radial glia (Liesi and Silver 1988). However, in situ hybridization at E16 in the mouse did not show signals for nidogen in radial glia (W. Halfter, personal communication). Thus, additional studies are needed to determine whether nidogen is expressed by radial glia in vivo during early stages of neural development.

Several other proteins related to cell adhesion were up-regulated dramatically in the cells expressing actNotch1 and they may be involved in functions of radial glia. One of these is ALCAM/DM-GRASP/SC-1/BEN, which is a member of the Ig superfamily that has been implicated in axon guidance and is expressed on Bergmann glia (Mothe and Brown 2002); it may play a key role in non-radial cell migration during brain development (Heffron and Golden 2000). N-CAM expression was also upregulated by actNotch1 and it is expressed on radial glia in developing dorsal cortex (Li and Grumet 2007). CD44 is a receptor for hyaluronic acid implicated in axon guidance across the optic chiasm, which is composed of specialized radial glial cells (Marcus and Mason 1995; Sretavan et al. 1994). A subpopulation of astrocyte restricted precursors express CD44 (Liu et al. 2004), consistent with our results that the actNotch1 expressing cells responded to differentiation conditions by expressing the astroglial marker GFAP. Little is known about nELL2 but its mRNA has been detected in E12 ventricular zone where cell bodies of radial glia are located (Kim et al. 2002). Thus, most of the cell adhesion genes that were identified are associated with radial glia including several proteoglycans that contribute to brain extracellular matrix (Morgenstern et al. 2002).

Expression of Notch1 itself was also upregulated in the NL2.3-4 cells expressing actNotch1 by comparison to L2.3 cells (Supplemental Table 1). This is consistent with the known action of Notch in lateral inhibition where transient activation of Notch leads to its upregulation followed by down-regulation and cell differentiation (Lai 2004). The Notch ligands delta-like homolog and jagged1 were down regulated in NL2.3-4 vs. L2.3, consistent with the upregulation of Notch (Supplemental Table 1). Notch1 expression in the CNS is normally transient and is typically followed by differentiation into neurons and glia (Anthony et al. 2005; Gaiano et al. 2000; Temple 2001). The resistance of NL2.3-4 cells to differentiate is also expected with the persistent activation of Notch signaling.

Notch1 behaves as a tumor suppressor in neuroepithelial cells (Lefort and Dotto 2004) although it induces tumors in hematopoietic progenitors (Vilimas et al. 2007). Although Notch1 expression may be required for survival of gliomas, its expression correlates inversely with glioma grade (Purow et al. 2005). We were unable to obtain stable clones from NPC after inducing actNotch1 expression except in cells that stably expressed v-myc (e.g. NL2.3-4). Interestingly, although high levels of actNotch1 expression were observed after transient infection, the stable clones that were obtained only expressed moderate levels, suggesting that high levels of Notch1 may inhibit cell growth and prevent clonal expansion as suggested (Lowell et al. 2006). This is also consistent with the observation that Notch1 increased the fraction of BLBP+/RC2+/BrdU- quiescent radial glia, suggesting that it inhibits cell proliferation (Gaiano et al. 2000). ActNotch1 expression slowed cell cycle progression slightly in NL2.3-4 cells by comparison to the parental L2.3 cells (Li, H., Chang, Y. and Grumet, M., unpublished observations), consistent with a tumor suppressive action (Fan et al. 2004). Transplantation studies using L2.3 indicated that they form masses resembling tumors whereas the NL2.3-4 cells derived from them, which express actNotch1, migrated robustly in vivo and showed no indication of tumorigenicity one month after transplantation (Fig. 5). Moreover, the NL2.3-4 cells continued to express nestin after transplantation into the spinal cord, in contrast to other NPC with or without v-myc that showed significant astroglial differentiation (Cao et al. 2001; Hasegawa et al. 2005). Whereas actNotch1 may be a tumor suppressor in NPC, Notch2 has the opposite effect (Fan et al. 2004).

Prolonged expression of actNotch1 in NPC maintains their radial glial properties, suggesting that they may promote recovery following spinal cord injury by modulating the environment to one that more closely resembles the developing nervous system where regeneration even in mammals is more robust than it is adults. This suppression of differentiation may limit reactive astrogliosis, which is not beneficial in the injured spinal cord (Silver and Miller 2004). Moreover, the tumor suppressive properties of actNotch1 at least in neural cells may provide an extra measure of safety that may make its expression in NSC or NPC favorable for long-term cell transplantation. Additional studies are needed to analyze this in longer-term transplants to verify that tumors do not form.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgements

We thank Dr. Nick Gaiano for viruses, Drs. Nat Heintz, Tod Anthony, Randall McKinnon and Zaven Kaprielian, for antibodies, and Dr. Peter Yurchenco for nidogen proteins. Supported by grants from NIH, the New Jersey Commission on Spinal Cord Research, the New Jersey Commission on Science & Technology, and Invitrogen, Inc. CR is an NSF-IGERT predoctoral fellow.

References

- Alvarez-Buylla A, Seri B, Doetsch F. Identification of neural stem cells in the adult vertebrate brain. *Brain Res Bull* 2002;57(6):751–758. [PubMed: 12031271]
- Androutsellis-Theotokis A, Leker RR, Soldner F, Hoepfner DJ, Ravin R, Poser SW, Rueger MA, Bae SK, Kittappa R, McKay RD. Notch signalling regulates stem cell numbers in vitro and in vivo. *Nature* 2006;442(7104):823–826. [PubMed: 16799564]
- Anthony TE, Klein C, Fishell G, Heintz N. Radial glia serve as neuronal progenitors in all regions of the central nervous system. *Neuron* 2004;41(6):881–890. [PubMed: 15046721]
- Anthony TE, Mason HA, Gridley T, Fishell G, Heintz N. Brain lipid-binding protein is a direct target of Notch signaling in radial glial cells. *Genes Dev* 2005;19(9):1028–1033. [PubMed: 15879553]
- Bartsch S, Bartsch U, Dorries U, Faissner A, Weller A, Ekblom P, Schachner M. Expression of tenascin in the developing and adult cerebellar cortex. *J Neurosci* 1992;12(3):736–749. [PubMed: 1372043]
- Bauer S, Kerr BJ, Patterson PH. The neuropoietic cytokine family in development, plasticity, disease and injury. *Nat Rev Neurosci* 2007;8(3):221–232. [PubMed: 17311007]
- Bedford L, Walker R, Kondo T, van Cruchten I, King ER, Sablitzky F. Id4 is required for the correct timing of neural differentiation. *Dev Biol* 2005;280(2):386–395. [PubMed: 15882580]
- Cao QL, Howard RM, Dennison JB, Whittemore SR. Differentiation of engrafted neuronal-restricted precursor cells is inhibited in the traumatically injured spinal cord. *Exp Neurol* 2002;177(2):349–359. [PubMed: 12429182]
- Cao QL, Zhang YP, Howard RM, Walters WM, Tsoulfas P, Whittemore SR. Pluripotent stem cells engrafted into the normal or lesioned adult rat spinal cord are restricted to a glial lineage. *Exp Neurol* 2001;167(1):48–58. [PubMed: 11161592]
- Conti L, Pollard SM, Gorba T, Reitano E, Toselli M, Biella G, Sun Y, Sanzone S, Ying QL, Cattaneo E, others. Niche-independent symmetrical self-renewal of a Mammalian tissue stem cell. *PLoS Biol* 2005;3(9):e283. [PubMed: 16086633]
- de la Pompa JL, Wakeham A, Correia KM, Samper E, Brown S, Aguilera RJ, Nakano T, Honjo T, Mak TW, Rossant J, others. Conservation of the Notch signalling pathway in mammalian neurogenesis. *Development* 1997;124(6):1139–1148. [PubMed: 9102301]
- Fan X, Mikolaenko I, Elhassan I, Ni X, Wang Y, Ball D, Brat DJ, Perry A, Eberhart CG. Notch1 and notch2 have opposite effects on embryonal brain tumor growth. *Cancer Res* 2004;64(21):7787–7793. [PubMed: 15520184]

- Feng L, Hatten ME, Heintz N. Brain lipid-binding protein (BLBP): a novel signaling system in the developing mammalian CNS. *Neuron* 1994;12(4):895–908. [PubMed: 8161459]
- Fox JW, Mayer U, Nischt R, Aumailley M, Reinhart D, Wiedemann H, Mann K, Timpl R, Krieg T, Chu M-L. Recombinant nidogen consists of three globular domains and mediates binding of laminin to collagen type IV. *European Molecular Biology Organization Journal* 1991;10:3137–3146.
- Frank NY, Margaryan A, Huang Y, Schatton T, Waaga-Gasser AM, Gasser M, Sayegh MH, Sadee W, Frank MH. ABCB5-mediated doxorubicin transport and chemoresistance in human malignant melanoma. *Cancer Res* 2005;65(10):4320–4333. [PubMed: 15899824]
- Gaiano N, Nye JS, Fishell G. Radial glial identity is promoted by Notch1 signaling in the murine forebrain. *Neuron* 2000;26(2):395–404. [PubMed: 10839358]
- Gotz M, Stoykova A, Gruss P. Pax6 controls radial glia differentiation in the cerebral cortex. *Neuron* 1998;21(5):1031–1044. [PubMed: 9856459]
- Halfter W, Dong S, Yip YP, Willem M, Mayer U. A critical function of the pial basement membrane in cortical histogenesis. *J Neurosci* 2002;22(14):6029–6040. [PubMed: 12122064]
- Hartfuss E, Galli R, Heins N, Gotz M. Characterization of CNS precursor subtypes and radial glia. *Dev Biol* 2001;229(1):15–30. [PubMed: 11133151]
- Hasegawa K, Chang Y-W, Berlin HL, Ikeda Y, Kane-Goldsmith O, Grumet M. Embryonic radial glia bridge spinal cord lesions and promote functional recovery following spinal cord injury. *Exp Neurol* 2005;193:394–410. [PubMed: 15869942]
- Hatten ME. Central nervous system neuronal migration. *Annu Rev Neurosci* 1999;22:511–539. [PubMed: 10202547]
- Heffron DS, Golden JA. DM-GRASP is necessary for nonradial cell migration during chick diencephalic development. *J Neurosci* 2000;20(6):2287–2294. [PubMed: 10704504]
- Heins N, Malatesta P, Cecconi F, Nakafuku M, Tucker KL, Hack MA, Chapouton P, Barde YA, Gotz M. Glial cells generate neurons: the role of the transcription factor Pax6. *Nat Neurosci* 2002;5(4):308–315. [PubMed: 11896398]
- Hibino S, Shibuya M, Engbring JA, Mochizuki M, Nomizu M, Kleinman HK. Identification of an active site on the laminin alpha5 chain globular domain that binds to CD44 and inhibits malignancy. *Cancer Res* 2004;64(14):4810–4816. [PubMed: 15256450]
- Hitoshi S, Alexson T, Tropepe V, Donoviel D, Elia AJ, Nye JS, Conlon RA, Mak TW, Bernstein A, van der Kooy D. Notch pathway molecules are essential for the maintenance, but not the generation, of mammalian neural stem cells. *Genes Dev* 2002;16(7):846–858. [PubMed: 11937492]
- Hormigo A, McCarthy M, Nothias JM, Hasegawa K, Huang W, Friedlander DR, Fischer I, Fishell G, Grumet M. Radial glial cell line C6-R integrates preferentially in adult white matter and facilitates migration of coimplanted neurons in vivo. *Exp Neurol* 2001;168(2):310–322. [PubMed: 11259119]
- Kim H, Ha CM, Choi J, Choi EJ, Jeon J, Kim C, Park SK, Kang SS, Kim K, Lee BJ. Ontogeny and the possible function of a novel epidermal growth factor-like repeat domain-containing protein, NELL2, in the rat brain. *J Neurochem* 2002;83(6):1389–1400. [PubMed: 12472893]
- Klassen H, Schwartz MR, Bailey AH, Young MJ. Surface markers expressed by multipotent human and mouse neural progenitor cells include tetraspanins and non-protein epitopes. *Neurosci Lett* 2001;312(3):180–182. [PubMed: 11602340]
- Kriegstein AR, Gotz M. Radial glia diversity: a matter of cell fate. *Glia* 2003;43(1):37–43. [PubMed: 12761864]
- Lai EC. Notch signaling: control of cell communication and cell fate. *Development* 2004;131(5):965–973. [PubMed: 14973298]
- Lefort K, Dotto GP. Notch signaling in the integrated control of keratinocyte growth/differentiation and tumor suppression. *Semin Cancer Biol* 2004;14(5):374–386. [PubMed: 15288263]
- Li H, Babiarz J, Woodbury J, Kane-Goldsmith N, Grumet M. Spatiotemporal heterogeneity of CNS radial glial cells and their transition to restricted precursors. *Dev Biol* 2004;271(2):225–238. [PubMed: 15223331]
- Li H, Berlin Y, Hart RP, Grumet M. Microtubules are critical for radial glial morphology: possible regulation by MAPs and MARKs. *Glia* 2003;44(1):37–46. [PubMed: 12951655]
- Li H, Grumet M. BMP and LIF signaling coordinately regulate lineage restriction of radial glia in the developing forebrain. *Glia* 2007;55(1):24–35. [PubMed: 17001659]

- Liesi P, Silver J. Is astrocyte laminin involved in axon guidance in the mammalian CNS? *Developmental Biology* 1988;130:774–785. [PubMed: 3143614]
- Liu KJ, Harland RM. Cloning and characterization of *Xenopus* Id4 reveals differing roles for Id genes. *Dev Biol* 2003;264(2):339–351. [PubMed: 14651922]
- Liu Y, Han SS, Wu Y, Tuohy TM, Xue H, Cai J, Back SA, Sherman LS, Fischer I, Rao MS. CD44 expression identifies astrocyte-restricted precursor cells. *Dev Biol* 2004;276(1):31–46. [PubMed: 15531362]
- Liu Y, Wu Y, Lee JC, Xue H, Pevny LH, Kaprielian Z, Rao MS. Oligodendrocyte and astrocyte development in rodents: an in situ and immunohistological analysis during embryonic development. *Glia* 2002;40(1):25–43. [PubMed: 12237841]
- Lowell S, Benchoua A, Heavey B, Smith AG. Notch promotes neural lineage entry by pluripotent embryonic stem cells. *PLoS Biol* 2006;4(5):e121. [PubMed: 16594731]
- Marcus RC, Mason CA. The first retinal axon growth in the mouse optic chiasm: axon patterning and the cellular environment. *J Neurosci* 1995;15(10):6389–6402. [PubMed: 7472403]
- Marin-Husstege M, He Y, Li J, Kondo T, Sablitzky F, Casaccia-Bonnet P. Multiple roles of Id4 in developmental myelination: predicted outcomes and unexpected findings. *Glia* 2006;54(4):285–296. [PubMed: 16862533]
- McKee KK, Harrison D, Capizzi S, Yurchenco PD. Role of laminin terminal globular domains in basement membrane assembly. *J Biol Chem* 2007;282(29):21437–21447. [PubMed: 17517882]
- Morgenstern DA, Asher RA, Fawcett JW. Chondroitin sulphate proteoglycans in the CNS injury response. *Prog Brain Res* 2002;137:313–332. [PubMed: 12440375]
- Mothe AJ, Brown IR. Effect of hyperthermia on the transport of mRNA encoding the extracellular matrix glycoprotein SC1 into Bergmann glial cell processes. *Brain Res* 2002;931(2):146–158. [PubMed: 11897100]
- Noble M, Arhin A, Gass D, Mayer-Proschel M. The cortical ancestry of oligodendrocytes: common principles and novel features. *Dev Neurosci* 2003;25(2–4):217–233. [PubMed: 12966219]
- Pfenninger CV, Roschupkina T, Hertwig F, Kottwitz D, Englund E, Bengzon J, Jacobsen SE, Nuber UA. CD133 is not present on neurogenic astrocytes in the adult subventricular zone, but on embryonic neural stem cells, ependymal cells, and glioblastoma cells. *Cancer Res* 2007;67(12):5727–5736. [PubMed: 17575139]
- Purow BW, Haque RM, Noel MW, Su Q, Burdick MJ, Lee J, Sundaresan T, Pastorino S, Park JK, Mikolaenko I, others. Expression of Notch-1 and its ligands, Delta-like-1 and Jagged-1, is critical for glioma cell survival and proliferation. *Cancer Res* 2005;65(6):2353–2363. [PubMed: 15781650]
- Rakic P. Principles of neural cell migration. *Experientia* 1990;46:882–891. [PubMed: 2209797]
- Reichardt LF, Tomaselli KJ. Extracellular matrix molecules and their receptors: Functions in neural development. *Annual Review of Neuroscience* 1991;14:531–570.
- Sievers J, Pehlemann FW, Gude S, Berry M. Meningeal cells organize the superficial glia limitans of the cerebellum and produce components of both the interstitial matrix and the basement membrane. *J Neurocytol* 1994;23(2):135–149. [PubMed: 8195812]
- Silver J, Miller JH. Regeneration beyond the glial scar. *Nat Rev Neurosci* 2004;5(2):146–156. [PubMed: 14735117]
- Smirnov SP, McDearmon EL, Li S, Ervasti JM, Tryggvason K, Yurchenco PD. Contributions of the LG modules and furin processing to laminin-2 functions. *J Biol Chem* 2002;277(21):18928–18937. [PubMed: 11886875]
- Smyth GK. Linear models and empirical bayes methods for assessing differential expression in microarray experiments. *Stat Appl Genet Mol Biol* 2004;3:Article3
- Sretavan DW, Feng L, Pure E, Reichardt LF. Embryonic neurons of the developing optic chiasm express L1 and CD44, cell surface molecules with opposing effects on retinal axon growth. *Neuron* 1994;12(5):957–975. [PubMed: 7514428]
- Tanigaki K, Nogaki F, Takahashi J, Tashiro K, Kurooka H, Honjo T. Notch1 and Notch3 instructively restrict bFGF-responsive multipotent neural progenitor cells to an astroglial fate. *Neuron* 2001;29(1):45–55. [PubMed: 11182080]
- Temple S. Stem cell plasticity--building the brain of our dreams. *Nat Rev Neurosci* 2001;2(7):513–520. [PubMed: 11433376]

- Vilimas T, Mascarenhas J, Palomero T, Mandal M, Buonamici S, Meng F, Thompson B, Spaulding C, Macaroun S, Alegre ML. others. Targeting the NF-kappaB signaling pathway in Notch1-induced T-cell leukemia. *Nat Med* 2007;13(1):70–77. [PubMed: 17173050]
- Villa A, Snyder EY, Vescovi A, Martinez-Serrano A. Establishment and properties of a growth factor-dependent, perpetual neural stem cell line from the human CNS. *Exp Neurol* 2000;161(1):67–84. [PubMed: 10683274]
- Wettenhall JM, Simpson KM, Satterley K, Smyth GK. affyImGUI: a graphical user interface for linear modeling of single channel microarray data. *Bioinformatics* 2006;22(7):897–899. [PubMed: 16455752]
- Wettenhall JM, Smyth GK. limmaGUI: a graphical user interface for linear modeling of microarray data. *Bioinformatics* 2004;20(18):3705–3706. [PubMed: 15297296]
- Yoon K, Gaiano N. Notch signaling in the mammalian central nervous system: insights from mouse mutants. *Nat Neurosci* 2005;8(6):709–715. [PubMed: 15917835]
- Yoon K, Nery S, Rutlin ML, Radtke F, Fishell G, Gaiano N. Fibroblast growth factor receptor signaling promotes radial glial identity and interacts with Notch1 signaling in telencephalic progenitors. *J Neurosci* 2004;24(43):9497–9506. [PubMed: 15509736]
- Yurchenco PD, Schittny JC. Molecular architecture of basement membranes. *Faseb J* 1990;4(6):1577–1590. [PubMed: 2180767]

Fig. 1.

Activated Notch1 promotes radial glial phenotype in cultured L2.3 clone and E13.5 cortical cells. (A) L2.3 cells infected with retrovirus overexpressing actNotch1(actNotch1-GFP), showed enhanced radial morphology and elevated expression of BLBP by comparison to control infection (GFP). Activated Notch1 fused with GFP translocates to the nucleus (arrow). Nestin (red) shows cellular morphology, while DAPI (blue) labels nuclei. The percentage of BLBP (purple) positive cells among infected cells were quantified (B) (n=493 for GFP, n=473 for actNotch1-GFP). (C) E13.5 cortical cells were infected with actNotch1-GFP retrovirus showed long bipolar morphology (arrow), while others were flat (arrowheads). The expression

levels of BLBP protein were quantified by relative fluorescence intensity (D) (n=76 for GFP; n=85 for actNotch1-GFP). Scale bars, 20 μ m.

Fig. 2.

Activated Notch1 suppresses expression of the glial restricted precursors marker 4D4. E13.5 cortical radial glia and L2.3 clone were infected with either actNotch1-GFP (green nucleus) or control GFP retrovirus. The resulting cultures were fixed and stained with the glial restricted precursor marker 4D4 (red) (Li and Grumet 2007), which was dramatically lower in the actNotch1-GFP infected cells compared to controls. Percentages of 4D4+ and 4D4- cells were measured among infected (GFP+) cells shown in panels on right. DAPI (blue) was used to label the nuclei. Scale bars, 10 μ m.

Fig. 3.

Cellular characterization of an actNotch1 expressing clone, NL2.3-4. (A) Expression of cellular markers, BLBP, Nestin and 4D4, were compared between NL2.3-4 and its parental radial glia clone L2.3 by immunostaining. DIC shows that NL2.3-4 cells are more bipolar than the L2.3 cells. ActNotch1-GFP shows green nuclear localization, while DAPI labels nuclei blue. The percentage of marker positive cells is shown in (B). Lengths of cellular processes was measured based on nestin staining and categorized into groups by length (C). Scale bar, 50 μm .

Fig. 4. NL2.3–4 cells differentiate into GFAP+ cells, but not into oligodendrocytes and neurons. (A) L2.3 and NL2.3–4 cells were cultured for 6 days after FGF2 withdrawal on laminin-coated substrate. Cells were fixed and stained with antibodies against GalC (oligodendrocytes), β -III tubulin (neurons), GFAP (astrocytes), nestin (NSC/radial glia) and DAPI (nuclei). All cells were nestin positive and the percentage of cells positive for differentiation markers is shown in (B).

Fig. 5.

ActNotch1 promoted migration of L2.3 in white matter of adult rat spinal cord. NL2.3–4 and L2.3 cells labeled with CellTracker CFDA-SE were transplanted into adult rat spinal cord white matter. After 3 days, rats were killed and histological analysis of spinal cords after sectioning revealed cell migration patterns primarily in white matter (WM). Red arrows indicate the injection sites (A, NL2.3–4 and B, L2.3). Longitudinal migration distances of the transplanted cells were measured in all sections that contained labeled cells. Rostro-caudal (X) and dorso-ventral (Y) cell migration distances are defined by the arrows in (B); results show averages with standard derivations (C) and ratios (X/Y) of migration distances (D). Total rostro-caudal migration distances of NL2.3–4 transplants in spinal cord at day 3, 14, 28 were measured (E). Four weeks after transplantation in spinal cords, NL2.3–4 cells maintained bipolar morphologies with long processes as revealed by nestin staining (red in F). Confocal images showed nestin positive cellular processes (red) with persistent actNotch1-GFP expression in the nucleus (G, arrow) of bipolar cells that lacked GFAP (purple) expression (H). Scale bars: 200 μm in (A) and (B), 100 μm in (F), 50 μm in (G).

Fig. 6. Microarray analysis indicates higher expression of adhesion molecules in NL2.3-4 cells than L2.3 cells. Gene expression comparison was performed in triplicates between NL2.3-4 and L2.3 RNAs using Affymetrix Rat 230-2 microarrays. Among the 31,100 genes on the chip, 720 genes were higher in NL2.3-4 and 349 genes were higher in L2.3 by 5% FDR and two-fold cutoff. Radial glial (RG) and Notch related genes (A) as well as adhesion molecules are listed (B) along with confirmation of candidates by Q-RT-PCR (C) and western blotting (insert in C).

Fig. 7.

NL2.3-4 cells adhered and migrated better on laminin substrate than parental L2.3 cells. Dissociated single cell suspensions of NL2.3-4 (A) and L2.3 (B) were incubated on laminin-coated spots for 40 min and then fixed and stained with propidium iodide (PI). Inverted PI stained images are shown with dotted circles outlining the laminin-coated spots. Average numbers of attached cells per spot and standard errors of triplicates are shown (C). Cultured neurospheres of NL2.3-4 and L2.3 labeled with DAPI were plated on laminin-coated substrates and incubated at 37°C. Phase contrast (D) and DAPI (E) pictures were taken soon after the neurosphere attachment (30 min) and when cells migrated out of neurospheres (2 hours). DAPI images (E) were used to quantify the neurosphere spreading area at 30 min (a1) and at 2 hours (a2). Averages of spreading area ratios from three different experiments (a2-a1)/a1 are shown (F) with standard errors. **, $P < 0.01$ (Student's t-Test). Scale bars, 40 μm in (A and B); 20 μm in (D and E).

Fig. 8.

Nidogen enhanced adhesion of NL2.3-4 cells on laminin. Dissociated cells were incubated on laminin for 40 min in the presence of the indicated proteins (20 μ g/ml). Images of DAPI stained adherent cells are shown in (A-C). The percentages of cells adhered were calculated with the number of adherent cells in controls defined as 100% (D). The percentages of cells adhered to recombinant proteins (20 μ g/ml): LM-2 (laminin-2 isoform), LM-2-NS (laminin-2 containing mutation at nidogen-binding site) and r-Nid (nidogen protein) were calculated with the number of adherent cells on LM-2 defined as 100% (E). Overnight cultures of NL2.3-4 cells on substrates coated with LM (F), LM-2 (G) and LM-2-NS (H) were fixed and stained with anti-nidogen antibody (red). Note that nidogen protein was detected (red) on the cell surface and on the substrates except with LM-2-NS (H). Scale bars, 50 μ m. *, $p < 0.05$ (Student's t-Test).

Fig. 9.

Nidogen expression in primary E14.5 radial glial cells. E14.5 cortical cells were treated with the Golgi blocker, Brefeldin A (5 $\mu\text{g/ml}$) for 90 min followed by fixation with 4% paraformaldehyde (A–C). Immunostaining showed that some RC1+ (A) and vimentin+ (insert in A) radial glial cells express nidogen, but TuJ1+ neurons did not (B). Higher magnification of a radial glial cell showed nidogen-positive secretory vesicles inside the cell body (C) with Brefeldin A treatment. Nidogen protein was detected on the surface of NL2.3–4 cells even without Brefeldin A treatment (D, arrow indicates green nucleus). Scale bars, 10 μm .

Table 1

Primers for Q-RT-PCR

Gene names	Forward	Reverse
B-actin	CGTAGCCATCCAGGCTGTGT	CCAGTGGTACGACCAGAGGC
eGFP	GCAAAGACCCCAACGAGAAG	TCACGAAGTCCAGCAGGACC
Fabp7(BLBP)	GGAAGCTGACAGACAGCCAGA	CGCCCAGAGCCTTCATGTAC
Nestin	CAAGCAGCAGGGTCACTTCC	AGGTTTGTGGCTAAGGAGGTCA
vimentin	GAGCACCTGCAGTCATTCA	CGTGCCAGAGAAGCATTGTC
Hes5	GCACCAGCCCAACTCCAA	ACGGCCATCTCCAGGATGT
Pax6	TCTAACCGAAGGGCCAAGTG	GAGGAGACAGGTGTGGTGGG
Notch1	AGCACTGGAAGGACTCCCA	AAGGACCCAGCTTCCGT
Tnc	CCTTCATTAAGACCCGCTGG	TGCAGCCTCATGAAGCAGAG
Nid	AGCACCTTTCCTGGCTGACTT	TACACATTCCCCAGGCCATC
Cd44	AAGGACACGTGGTAATCCCG	TGGCTGCACAGATAGCGTTG
Alcam	ACCATTGTCATGCCTTGACAG	GCCAAACATGAGGTTCTGAGGT
NCAM	TGATGTGGTCAGCTCTCTGCC	CGGCCTTTGTGTTTCCAGAT
Itgb1	AAGTCCCAAGTGCCATGAGG	CTGCAGGCTCCACACTCAAAT